

Graphene Oxide-Gold Nanorods Nanocomposite-Porphyrin Conjugate as Promising Tool for Cancer Phototherapy Performance

Thabang Calvin Lebepe ^{1,2}, Sundararajan Parani ^{1,2}, Vuyelwa Ncapayi ^{1,2}, Rodney Maluleke ^{1,2}, Grace It Mwad Mbaz ^{1,2}, Olufunto Tolulope Fanoro ², Jose Rajendran Varghese ^{1,2}, Atsuki Komiya ³, Tetsuya Kodama ⁴ and Oluwatobi Samuel Oluwafemi ^{1,2,*}

¹ Department of Chemical Science, University of Johannesburg, Johannesburg 2028, South Africa; calvyn.tl@gmail.com (T.C.L.), sbarani416@gmail.com (S.P.); vuyelwa.tito.ncapayi530@gmail.com (V.N.); rodney.maluleke@gmail.com (R.M.); gmbazitmwad@gmail.com (G.I.M.M.); josv3209@gmail.com (J.R.V.)


² Centre for Nanomaterials Sciences Research, University of Johannesburg, Johannesburg 2028, South Africa; jolufunto@gmail.com

³ Graduate School of Engineering, Institute of Fluid Science, Tohoku University, Sendai 980-8577, Japan; komiya@tohoku.ac.jp



⁴ Graduate School of Biomedical Engineering, Tohoku University, Sendai 980-8575, Japan; kodama@tohoku.ac.jp



* Correspondence: oluwafemi.oluwatobi@gmail.com

SUPPORTING DOCUMENTS



Scheme S1. Schematic diagram for the synthesis of TMePyP.

Figure S1. ^1H -NMR of TPyP.Figure S2. ^1H -NMR of TMePyP.

Figure S3. ^{13}C -NMR of TPyP.Figure S4. ^{13}C -NMR of TMePyP.