

Supplementary Material

Development of Controlled-Release Carbamide Peroxide Loaded Nanoemulgel for Tooth Bleaching: In Vitro and Ex Vivo Studies

Siriporn Okonogi ^{1,2,*}, Adchareeya Kaewpinta ³, Sakornrat Khongkhunthian ^{2,4} and Pisaisit Chaijareenont ^{2,5}

¹ Department of Pharmaceutical Sciences, Faculty of Pharmacy, Chiang Mai University, Chiang Mai 50200, Thailand

² Research Center of Pharmaceutical Nanotechnology, Chiang Mai University, Chiang Mai 50200, Thailand;); sakornratk@hotmail.com (S.K.); yodent@hotmail.com (P.C.)

³ Interdisciplinary Program in Nanoscience and Nanotechnology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand; akaewpinta@gmail.com

⁴ Department of Restorative Dentistry and Periodontology, Faculty of Dentistry, Chiang Mai University, Chiang Mai 50200, Thailand

⁵ Department of Prosthodontics, Faculty of Dentistry, Chiang Mai University, Chiang Mai 50200, Thailand

* Correspondence: okng2000@hotmail.com; Tel.: +66-53-944-311

Figure S1: Release kinetics of CP from gel formulations according to (a) zero-order kinetics, (b) first-order kinetics, (c) Higuchi model, and (d) Korsmeyer–Peppas model.

