

Supplementary Materials

Screening and selection of a new medium for diosgenin production via microbial biocatalysis of *Fusarium* sp.

Wancang Liu ¹, Haibo Xiang ², Tao Zhang ¹, Xu Pang ¹, Jing Su ¹, Hongyu Liu ¹, Baiping Ma ³ and Liyan Yu ^{1,*}

¹ Division for Medicinal Microorganisms Related Strains CAMS Collection Center of Pathogenic Microorganisms, Institute of Medicinal Biotechnology, Chinese Academy of Medical Sciences & Peking Union Medical College, 100050 Beijing, China; liuwancang@imb.pumc.edu.cn (W.L.); zhangt@cpcc.ac.cn (T.Z.) pangxu@imb.pumc.edu.cn (X.P.); sujing@cpcc.ac.cn (J.S.); hylu@cpcc.ac.cn (H.L.)

² State Key Laboratory of Biocatalysis and Enzyme Engineering, School of Life Sciences, Hubei University, 430011 Wuhan, China; xhb2086@hubu.edu.cn

³ Beijing Institute of Radiation Medicine, 100850 Beijing, China; mabaiping@sina.com

* Correspondence: yly@cpcc.ac.cn; Tel.: +86 10 63187118

Figure S1: Diagnostic plots of the BBD model adequacy for diosgenin yield. (a) The plot of predicted values versus experimental values. (b) The plot of normal % probability. (c) The plot of internally studentized residuals.




Figure S2: The overlay plot of perturbation for diosgenin yield. A, wheat bran. B, phosphate. C, Tween-80.

