

Supporting information for
Study of cytotoxic and photodynamic activities of dyads composed of a zinc
phthalocyanine appended to an organotin

Isabelle Toubia,^{ac#} Christophe Nguyen,^{b#} Stéphane Diring,^a Marine Pays,^b Elodie Mattana,^b Philippe Arnoux^c, Céline Frochot,^{*c} Magali Gary-Bobo,^{*b} Marwan Kobeissi,^{*d} Fabrice Odobel^{*a}

^aCEISAM, Chimie Et Interdisciplinarité, Synthèse, Analyse, Modélisation, CNRS, UMR CNRS 6230, UFR des Sciences et des Techniques ; 2, rue de la Houssinière - BP 92208; 44322 NANTES Cedex 3 (France). E-mail: Fabrice.Odobel@univ-nantes.fr

^bIBMM, Univ Montpellier, CNRS, ENSCM, Montpellier, France. E-mail: magali.gary-bobo@inserm.fr

^cLRGP, Laboratoire Réactions et Génie des Procédés, UMR 7274 CNRS-Université de Lorraine, 1 rue Grandville, 54000 Nancy, France. E-mail: celine.frochot@univ-lorraine.fr

^dLaboratoire RammalRammal, Equipe de Synthèse Organique Appliquée SOA, Université Libanaise, Faculté des Sciences 5, Nabatieh, Liban. E-mail: mkobeissi@ul.edu.lb

[#]These authors contributed equally to this study.




Figure S1. Overlay of the UV-Vis. absorption spectra of compounds **SnPh₂** and **Sn₂Ph₆** recorded in dichloromethane.




Figure S2. Mass spectrum of phthalocyanine 3.




Figure S3. Mass spectrum of phthalocyanine 4.




Figure S4. Mass spectrum of compound ZnPcSnPh_2




Figure S5. Mass spectrum of compound $\text{ZnPcSn}_2\text{Ph}_6$




Figure S6. ¹H NMR spectrum of phthalocyanine **3** recorded in CDCl₃.




Figure S7. ¹H NMR spectrum of phthalocyanine **4** recorded in DMSO-d₆ with a trace of pyridine.




Figure S8. ^1H NMR spectrum of compound **ZnPc-SnPh₂** recorded in THF- d_8 .




Figure S9. ^1H NMR spectrum of compound **ZnPc-Sn₂Ph₆** recorded in DMSO- d_6 with a trace of pyridine.