

Neuroprotective Effect of Artichoke-Based Nanoformulation in Sporadic Alzheimer's Disease Mouse Model: Focus on Antioxidant, Anti-Inflammatory, and Amyloidogenic Pathways

Heba A. S. El-Nashar ¹, Haidy Abbas ², Mariam Zewail ², Mohamed H. Noureldin ³, Mai M. Ali ⁴, Marium M. Shamaa ³, Mohamed A. Khattab ⁵ and Nehal Ibrahim ^{1,*}

¹ Pharmacognosy Department, Faculty of Pharmacy, Ain Shams University, Cairo 11566, Egypt; heba_pharma@pharma.asu.edu.eg (H.A.S.E.-N.); nehal.sabry@pharma.asu.edu.eg (N.I.)

² Department of Pharmaceutics, Faculty of Pharmacy, Damanhour University, Damanhour 22514, Egypt; haidy.abass@pharm.dmu.edu.eg (H.A.); mariamzewail@pharm.dmu.edu.eg (M.Z.)

³ Department of Biochemistry, Division of Clinical and Biological Sciences, College of Pharmacy, Arab Academy for Science, Technology and Maritime Transport, Alexandria P.O. Box 1029, Egypt; mohamed.noureldin@aast.edu (M.H.N.); marium.muhammed@aast.edu (M.M.S.)

⁴ Department of Pharmaceutics, Division of Pharmaceutical Sciences, College of Pharmacy, Arab Academy for Science, Technology and Maritime Transport, Alexandria P.O. Box 1029, Egypt; drmaiali@aast.edu (M.M.A.)

⁵ Department of Cytology and Histology, Faculty of Veterinary Medicine, Cairo University, Cairo 12211, Egypt; mabdelrazik@cu.edu.eg

* Correspondence: nehal.sabry@pharma.asu.edu.eg




Figure S1. Total ion chromatogram (TIC) of the methanolic extract of *C. cardunculus* L. bracts. Peaks numbers follow those listed in Table 1.