



Supplementary Materials

Evidence of Increased Resistance in Neuroblastoma Cells after Administration of Drug Atorvastatin

Ana Salomé Correia ^{1,2,3}, Lara Marques ^{1,3} and Nuno Vale ^{1,3,4,*}

¹OncoPharma Research Group, Center for Health Technology and Services Research (CINTESIS), Rua Doutor Plácido da Costa, 4200-450 Porto, Portugal

²Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, Rua de Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal

³CINTESIS@RISE, Faculty of Medicine, University of Porto, Alameda Professor Hernâni Monteiro, 4200-319 Porto, Portugal

⁴Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine, University of Porto, Alameda Professor Hernâni Monteiro, 4200-319 Porto, Portugal

*Correspondence: nunovale@med.up.pt; Tel.: +351-220-426-537




Figure S1. Effect of 48h-incubation of echinomycin 0.1 nM-5 nM, on the viability of SH-SY5Y cells, determined by MTT assay. The results represent the mean \pm SEM of three independent experiments, expressed as the percentage of the vehicle (100%). Statistically significant * $p < 0.05$ vs. vehicle.