

SUPPLEMENTARY DATA**Cucumarioside A₂-2 Causes Macrophage Activation
In Mouse Spleen**

Pislyagin E.A.,^{1*} Manzhulo I.V.,^{2,3} Gorpenchenko T.Y.,⁴ Dmitrenok P.S.,¹ Avilov S.A.,¹ Silchenko A.S.,¹ Wang Y-M.,⁵ Aminin D. L.¹

¹ G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, Russia


² National Scientific Center of Marine Biology, Far Eastern Branch of the Russian Academy of Science, Vladivostok, Russia

³ School of Biomedicine, Far Eastern Federal University, Vladivostok, Russia

⁴ Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, Russia

⁵ Institute of Molecular Medicine and Bioengineering, Department of Biological Science and Technology, National Chiao Tung University, Taiwan

* Correspondence: pislyagin@hotmail.com; Tel.: 7(4232)31-99-32; Fax: 7(4232)31-40-50



Supplementary Fig. S2. The assignment of signals in the ^{13}C NMR spectrum of cucumarioside $\text{A}_2\text{-2}$.