
Supplementary Materials

Efficient Preparation of Bafilomycin A1 from Marine *Streptomyces lohii* Fermentation Using Three-Phase Extraction and High-Speed Counter-Current Chromatography

Ye Yuan ^{1,2}, Xiaoping He ¹, Tingting Wang ¹, Xingwang Zhang ³, Zhong Li ^{4,5}, Xiaoqing Xu ⁶, Weiyang Zhang ¹, Xiaojun Yan ^{2,*}, Shengying Li ^{3,*}, Shan He ^{1,*}

¹ Li Dak Sum Yip Yio Chin Kenneth Li Marine Biopharmaceutical Research Center, College of Food and Pharmaceutical Sciences, Ningbo University, Ningbo, Zhejiang 315800, China; 23yuanye@163.com (Y.Y.); 18892628896@163.com (X.H.); 18251965896@163.com (T.W.); zhangweiyang13@126.com (W.Z.)

² School of Marine Science, Ningbo University, Ningbo, Zhejiang 315211, China

³ State Key Laboratory of Microbial Technology, Shandong University, Qingdao, Shandong 266237, China; zhangxingwang@sdu.edu.cn (X.Z.)

⁴ Shandong Provincial Key Laboratory of Synthetic Biology, and CAS Key Laboratory of Biofuels at Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao, Shandong 266101, China; lizhong@qibebt.ac.cn (Z.L.)

⁵ University of Chinese Academy of Sciences, Beijing 100049, China

⁶ Rushan Hanwei Biological Science and Technology Co., Ltd., Rushan, Shandong 264502, China; xiaoqingxu@163.com (X.X.)

* Correspondence: yanxiaojun@nbu.edu.cn (X.Y.); lishengying@sdu.edu.cn (S.L.); heshan@nbu.edu.cn (S.H.); Fax: +86-574-87600458 (X.Y.); +86-574-87604388 (S.H.).




Figure S1. ¹H NMR (500 MHz, CDCl₃) spectrum of bafilomycin A1




Figure S2. ¹³C NMR (125 MHz, CDCl₃) spectrum of bafilomycin A1