

A concise route for the synthesis of tetracyclic meroterpenoids: (\pm)-Aureol preparation and mechanistic interpretation.

Antonio Rosales Martínez,^{1*} Lourdes Enríquez,² Martín Jaraíz,² Laura Pozo Morales,¹ Ignacio Rodríguez-García,³ Emilio Díaz Ojeda.¹

¹ Department of Chemical Engineering, Escuela Politécnica Superior, University of Sevilla, 41011 Sevilla, Spain

² Department of Electricity and Electronics, University of Valladolid, Valladolid, Spain

³ Organic Chemistry, ceiA3, University of Almería, 04120 Almería, Spain


Correspondence: arosales@us.es

SUPPORTING INFORMATION


Table of contents

¹ H NMR of (\pm)-albicanol (5).....	S-2
¹ H NMR of (\pm)-albicanal (6).....	S-3
¹ H NMR of 7	S-4
¹³ C NMR of 7	S-5
¹ H NMR of 8	S-6
¹³ C NMR of 8	S-7
¹ H NMR of 9	S-8
¹³ C NMR of 9	S-9
¹ H NMR of 10	S-10
¹³ C NMR of 10	S-11
¹ H NMR of (\pm)-aureol ((\pm)- 1).....	S-12
¹³ C NMR of (\pm)-aureol ((\pm)- 1).....	S-13


¹H NMR of (±)-albicanol (5)




^1H NMR of (\pm)-albicanal (6)




¹H NMR of 7




¹³C NMR of 7




¹H NMR of 8




¹³C NMR of 8




¹H NMR of 9




¹³C NMR of 9




¹H NMR of 10



¹³C NMR of 10



^1H NMR of (\pm) -aureol (\pm) -1



^{13}C NMR of (\pm)-aureol ((\pm)-1)

