

Sdy-1 executes antitumor activity in HepG2 and Hela cancer cells by inhibiting the Wnt/ β -catenin signaling pathway

Mengyu Sun ^{1,‡}, Dongdong Zhou ^{1,‡}, Jingwan Wu¹, Jing Zhou ² and Jing Xu ^{1,*}

¹ One Health Institute, School of Chemical Engineering and Technology, Hainan University, Haikou 570228, China; summer@hainanu.edu.cn (M. S.); dongdongchoy@hainanu.edu.cn (D. Z.); 2021110817000019@hainanu.edu.cn (J. W.)

² School of Life Sciences, Hainan University, Haikou 570228, China; 993725@hainanu.edu.cn (J. Z.)

* Correspondence: happyjing3@hainanu.edu.cn (J. X.)

‡ These authors contributed equally to this work.

Contents

Figure S1. ^1H -NMR of Sdy-1	3
Figure S2. ^{13}C -NMR of Sdy-1	3
Figure S3. DEPT of Sdy-1	4
Figure S4. ^1H - ^1H COSY of Sdy-1	4
Figure S5. HMQC of Sdy-1	5
Figure S6. HMBC of Sdy-1	5
Figure S7. NOESY of Sdy-1	6
Figure S8. ESI-MS of Sdy-1.....	6




Figure S1. ¹H-NMR of Sdy-1




Figure S2. ¹³C-NMR of Sdy-1




Figure S3. DEPT of Sdy-1




Figure S4. ^1H - ^1H COSY of Sdy-1




Figure S5. HMQC of Sdy-1




Figure S6. HMBC of Sdy-1




Figure S7. NOESY of Sdy-1




Figure S8. ESI-MS of Sdy-1