

Supplementary Materials

Identification of Axinellamines A and B as Anti-Tubercular Agents

Emily J. Strong ^{1,#}, Lendl Tan ^{1,#}, Sasha Hayes ^{2,3}, Hayden Whyte ¹, Rohan A. Davis ^{2,3} and Nicholas P. West ^{1,*}

¹ School of Chemistry and Molecular Biosciences, and the Australian Infectious Diseases Research Centre, The University of Queensland, Brisbane QLD 4072, Australia

² Griffith Institute for Drug Discovery, School of Environment and Science, Griffith University, Brisbane QLD 4111, Australia

³ NatureBank, Griffith University, Brisbane QLD 4111, Australia

* Correspondence: n.west@uq.edu.au

These authors contributed equally to this work

Contents:

- Figure S1** *M. tuberculosis* inhibition of bioassay-guided fractions from NatureBank extracts
Figure S2 ^1H NMR (800 MHz) Spectrum of TFA Salt of Axinellamine A (1) in $\text{DMSO-}d_6$ at 25 °C
Figure S3 UHPLC-MS of TFA Salt of Axinellamine A (1)
Figure S4 ^1H NMR (800 MHz) Spectrum of TFA Salt of Axinellamine B (2) in $\text{DMSO-}d_6$ at 25 °C
Figure S5 UHPLC-MS of TFA Salt of Axinellamine B (2)

Figure S1: *M. tuberculosis* inhibition of bioassay-guided fractions from NatureBank extracts




Figure S2: ^1H NMR (800 MHz) Spectrum of TFA Salt of Axinellamine A (**1**) in $\text{DMSO-}d_6$ at 25 $^\circ\text{C}$




Figure S3: UHPLC-MS of TFA Salt of Axinellamine A (1)




Figure S4: ^1H NMR (800 MHz) Spectrum of TFA Salt of Axinellamine B (**2**) in $\text{DMSO-}d_6$ at 25°C




Figure S5: UHPLC-MS of TFA Salt of Axinellamine B (2)

