



Article

# Association between the First Occurrence of Allergic Rhinitis in Preschool Children and Air Pollution in Taiwan

Hui-Ying Chung <sup>1,2</sup>, Chia-Jung Hsieh <sup>1</sup>, Chun-Chieh Tseng <sup>1</sup> and Lih-Ming Yiin <sup>1,\*</sup>

<sup>1</sup> Department of Public Health, Tzu Chi University, 701, Sec. 3, Zhongyang Rd., Hualien City 97062, Taiwan; fatty99@gms.tcu.edu.tw (H.-Y.C.); cjhsieh@mail.tcu.edu.tw (C.-J.H.); tsengcc@mail.tcu.edu.tw (C.-C.T.)

<sup>2</sup> TCU Center for Value-Added Health Data Analysis and Application, 701, Sec. 3, Zhongyang Rd., Hualien City 97062, Taiwan

\* Correspondence: lmyiin@mail.tcu.edu.tw; Tel.: +886-3-8565301 (ext. 2273)

Academic Editor: Kim Natasha Dirks

Received: 18 December 2015; Accepted: 23 February 2016; Published: 27 February 2016

**Abstract:** The occurrence of allergic rhinitis (AR) may be significantly influenced by air pollution. This study examined the association between the first occurrence of AR in preschool children and the pre-incident levels of air pollutants in Taiwan. We identified 9960 eligible subjects from a systematic sampling cohort database containing 400,000 insureds of the National Health Insurance from 2007 to 2011 and matched them with the environmental monitoring data from 2006 to 2011 according to the locations of their clinics. Pre-incident levels were determined using the average concentrations of air pollutants one or two weeks prior to the AR diagnoses. Logistic regression analyses were performed to determine any significant relationships between AR and specific air pollutants. The first AR incidence for Taiwanese preschool children, which increased with age, was 10.9% on average; boys appeared to have a higher percentage (14.2%) than girls (8.27%). Among the air pollutants, carbon monoxide (CO) and nitrogen oxides (NO<sub>x</sub>) were significantly related to AR after adjusting for age and gender ( $p < 0.05$ ). Because both pollutants are considered to be traffic emissions, this study suggests that traffic emissions in Taiwan need to be controlled to lower the prevalence of children's AR.

**Keywords:** air pollution; allergic rhinitis; nitrogen oxides; preschool children; traffic emission

## 1. Introduction

Allergic rhinitis (AR) is a common respiratory disease occurring in individuals who breathe air with allergens that cause nose inflammation and swelling. A variety of allergens can trigger AR, including molds, pollens, animal dander, and dust mites [1]. Seasonal changes, such as temperature and humidity variations, may also play a role in exacerbating AR [2–4]. When the nose suffers from AR, it becomes blocked, which likely also blocks the sinuses (sinusitis); both are irritating and cause discomfort and headaches, all of which may interfere with daily life. Although AR is usually not considered a severe disease, its potential to develop into asthma is of great concern and warrants investigation [5].

A number of studies have indicated that the increasing prevalence of AR is associated with air pollution severity [2,6–12]. Of the commonly monitored air pollutants, which include carbon monoxide (CO), nitrogen oxides (NO<sub>x</sub>), ozone (O<sub>3</sub>), particulate matter (PM), and sulfur dioxide (SO<sub>2</sub>), several have been found to be related to AR; however, the relationships are not consistent. For example, NO<sub>2</sub>, PM<sub>10</sub> (aerodynamic diameter  $\leq 10 \mu\text{m}$ ) and SO<sub>2</sub> have been associated with AR in China [11,13,14], whereas in the United States, O<sub>3</sub> and PM<sub>2.5</sub> (aerodynamic diameter  $\leq 2.5 \mu\text{m}$ ) are

the related environmental factors [7]. Despite the differences in species, traffic-related air pollution or residence in an urban area appears to be an important factor for the increasing prevalence of AR in many countries [4,7,15–20].

Many studies have focused on the relationships between AR and the environment; however, not all age groups are studied evenly. There are relatively few data for preschool children, likely due to their low prevalence of AR or their limited ability to independently fill out research questionnaires. Nevertheless, preschool children's health warrants additional attention and care, and this issue is relevant for discussion and investigations. Thus, we conducted this study in preschool children using Taiwan's National Health Insurance Data, from which the AR subjects were identified by doctor's diagnoses, and combining the environmental data from the subjects' adjacent monitoring sites to determine the association between the first occurrence of AR and environmental pollutants.

## 2. Materials and Methods

### 2.1. Inclusion of Subjects

We used the systematic sampling cohort database of 400,000 insureds of the National Health Insurance of Taiwan from 2007 to 2011 and found 23,103 preschool children aged 0–6 years. After excluding those who did not have sufficient information or adjacent environmental monitoring data, there were 9960 children, of which 4456 were male and 5504 were female. The diagnoses were made following the International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM), and the codes 477.0, 477.1, 477.2, 477.8 and 477.9 were categorized as AR. Those who were assigned to any of the ICD-9CM codes were selected as study subjects, and only the first diagnoses were used to relate to the air pollution at one and two weeks prior to them. Each eligible subject must have had more than one AR-related visit and been confirmed to be AR-free for at least one year prior to the first AR diagnosis during the period; that is, the year of 2007 was considered a clearance period from which no AR records were retrieved. There were 7326 AR-related visits, of which 1088 children, 632 male and 456 female, were eligible and included in the study. Because of the secondary data with the removal of personal identification, the study was granted for exempt review by the Research Ethics Committee of Tzu Chi General Hospital (No: IRB102-57, approved on 23 April 2013).

### 2.2. Environmental Monitoring Data

Environmental monitoring sites adjacent to or sharing the same district codes with the subjects' clinics were used for the sources of environmental data, which came from 47 of the available 66 environmental monitoring sites in Taiwan. Data for the monitored air pollutants (CO, NO<sub>x</sub>, O<sub>3</sub>, PM<sub>10</sub>, and SO<sub>2</sub>) were available for download (in daily and monthly averages) from the Taiwan Air Quality Network (<http://taqm.epa.gov.tw/taqm/en/default.aspx>). To address possible latency in health effects for the cohort database covering 2007–2011, we retrieved the environmental monitoring data from one year ahead of the period (2006) to 2011.

### 2.3. Data Management and Analysis

Data were divided by age or gender to assess potential effects on AR incidence. The cohort database without personal identification provided the subjects' clinical locations at the county or district level, which was sufficient for linkage with adjacent environmental monitoring sites.

A logistic regression analysis was conducted to compare the rates between ages or genders, whereas Pearson correlation was performed to determine the relationship between the five air pollutants. We used a dummy variable for each subject's AR occurrence (1 = first occurrence; 0 = one of two randomly selected pre-AR time points) in a logistic regression analysis to calculate the odds ratios (ORs) for AR occurrence and the levels of air pollution when adjusting variables that may have been effective in incidence (e.g., age or gender). Because air pollution-induced respiratory symptoms were confirmed to be related to exposures for several days to weeks [21,22] and the diagnoses may be

registered days after the true occurrences, we used the environmental data from one and two weeks prior to the diagnoses for the pre-incident levels of air pollution in the regression analysis. The levels, which were determined by categorizing the average concentrations with quartiles of the descriptive statistics of the 2007–2011 environmental monitoring data, were Level 1 from the lowest to Q1, Level 2 from Q1 to Q3, and Level 3 from Q3 to the highest. All statistical analyses were performed using SAS 9.2.

### 3. Results

The first AR incidence in Taiwanese preschool children for 2007–2011 was 10.9%, with the boys' outcome significantly higher than the girls' (14.2% > 8.28%, OR = 1.83, 95% CI = 1.62–2.09) (Table 1). The rates significantly increased with age from 4.21% to 18.6% as age increased from 2 to 6. This increasing trend should be of concern to parents and caregivers because AR could lead to chronic respiratory diseases or asthma without proper care or treatment.

**Table 1.** Incidence rates of the first occurrence of allergic rhinitis in preschool children by gender and age for 2007–2011.

Population	Rhinitis Cases (%)	Odds Ratio (95% CI)
Total ( <i>n</i> = 9960)	1088 (10.9%)	
Gender		
Male ( <i>n</i> = 4456)	633 (14.2%)	1.84 (1.62, 2.09) *
Female ( <i>n</i> = 5504)	455 (8.27%)	Referent
Age at first diagnosis (years)		
0–2 ( <i>n</i> = 3342)	141 (4.21%)	Referent
3 ( <i>n</i> = 1732)	171 (9.87%)	2.49 (1.97, 3.13) *
4 ( <i>n</i> = 1803)	210 (11.6%)	2.99 (2.40, 3.74) *
5 ( <i>n</i> = 1574)	286 (18.2%)	5.04 (4.08, 6.23) *
6 ( <i>n</i> = 1509)	280 (18.6%)	5.18 (4.19, 6.41) *

\* Statistically significant.

The statistical results of the daily concentrations of the monitored air pollutants corresponding to the subjects are presented in Table 2. The maximum daily concentrations of the pollutants were below the ambient standards, indicating that air quality during the period in Taiwan was generally acceptable. The mean concentrations, which are considered to be the averages of annual concentrations from 2006 to 2011, were comparable with the annual ambient standards, which were available for PM<sub>10</sub> and SO<sub>2</sub>. The mean concentration of SO<sub>2</sub> (4.81 ± 1.85 ppb) was far below its annual standard (30 ppb), showing the efficacy in emission control, whereas that of PM<sub>10</sub> (56.8 ± 14.4 µg/m<sup>3</sup>) was close to the standard (65 µg/m<sup>3</sup>), suggesting a need for control efforts.

**Table 2.** Descriptive statistics of daily concentrations of air pollutants for 2006–2011 in Taiwan.

Air Pollutant	Mean ± SD	Minimum	25th Percentile (Q1)	Median (Q2)	75th Percentile (Q3)	Maximum	Taiwan's Ambient Standards
CO (ppb)	561 ± 172	150	448	553	618	1,070	9000 <sup>a</sup>
NO <sub>x</sub> (ppb)	19.6 ± 5.72	2.33	16.1	20.5	22.9	29.7	50 <sup>b</sup>
O <sub>3</sub> (ppb)	27.9 ± 3.85	20.5	25.9	27.7	29.9	41.7	60 <sup>a</sup>
PM <sub>10</sub> (µg/m <sup>3</sup> )	56.8 ± 14.4	21.7	47.9	56.7	65.0	88.1	65 <sup>b</sup> ; 125 <sup>c</sup>
SO <sub>2</sub> (ppb)	4.81 ± 1.85	1.90	3.51	4.32	6.11	11.7	30 <sup>b</sup> ; 100 <sup>c</sup>

CO, carbon monoxide; NO<sub>x</sub>, nitrogen oxides; O<sub>3</sub>, ozone; PM<sub>10</sub>, particulate matter with aerodynamic diameter ≤ 10 µm; SO<sub>2</sub>, sulfur dioxide; SD, standard deviation. <sup>a</sup> 8-h average; <sup>b</sup> annual average; <sup>c</sup> daily average.

The correlation between the air pollutants is given in Table 3. NO<sub>x</sub>, a byproduct of combustion, was highly correlated with CO and SO<sub>2</sub> ( $r = 0.851$  and  $0.809$ ), indicating emission sources of motor vehicles and industry/power plants, respectively, because CO was primarily produced by motor vehicles and SO<sub>2</sub> was a major combustion product of fuel [23]. NO<sub>x</sub> was also highly correlated with O<sub>3</sub> but in an opposite direction ( $r = -0.801$ ), and a similar correlation was found between SO<sub>2</sub> and O<sub>3</sub> ( $r = -0.485$ ). The negative association between NO<sub>x</sub> or SO<sub>2</sub> and O<sub>3</sub> suggests the consumption of either precursor of the photochemical smog reaction that generates O<sub>3</sub>. PM<sub>10</sub> was not significantly associated with any other variables with the exception of SO<sub>2</sub> ( $r = 0.531$ ), indicating that a certain quantity of PM<sub>10</sub> in Taiwan was generated from fuel combustion.

**Table 3.** Correlation between daily concentrations of the monitored air pollutants for 2016–2011 in Taiwan.

( <i>n</i> = 2191)	CO	NO <sub>x</sub>	O <sub>3</sub>	PM <sub>10</sub>	SO <sub>2</sub>
CO	1.000				
NO <sub>x</sub>	0.851 **	1.000			
O <sub>3</sub>	−0.744 **	−0.801 **	1.000		
PM <sub>10</sub>	0.195	0.320	−0.109	1.000	
SO <sub>2</sub>	0.630 *	0.809 **	−0.485 *	0.531 *	1.000

\*  $p < 0.05$ ; \*\*  $p < 0.01$ .

Age and gender both affected the AR incidence (as shown in Table 1); thus, we adjusted for those factors when conducting logistic regression analyses. The univariate analysis shows significant ORs for AR occurrence when comparing the higher pre-incident levels of CO or NO<sub>x</sub> (e.g., Levels 2 and 3) with the low level (Level 1) either one or two weeks prior to the occurrence (Table 4). Regarding other pollutants (O<sub>3</sub>, PM<sub>10</sub> and SO<sub>2</sub>), the level differences did not significantly affect AR occurrence. We also performed multivariate logistic regression analyses to determine the association between AR and all air pollutants together but excluded CO from the analyses because of its collinearity with NO<sub>x</sub> ( $r = 0.851$ ). NO<sub>x</sub> is the only variable that was significantly related to AR with ORs as high as 1.94 (95% CI = 1.41–2.71) for one week before and 1.50 (95% CI = 1.19–2.03) for two weeks before. The pre-incident levels one week before occurrence resulted in higher ORs than that two weeks before, suggesting that AR induced by air pollution should occur within a week. In summary, the relatively high pre-incident levels of traffic-related pollutants (NO<sub>x</sub> coupled with CO) were likely to lead to AR within the following week.

**Table 4.** Uni- and multivariate logistic regression analyses between AR occurrence and different pre-incident levels of air pollutants after adjustment for age and gender.

Air Pollutant Level	One Week before AR		Two Weeks before AR	
	Univariate	Multivariate	Univariate	Multivariate
( <i>n</i> = 3264)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
CO				
Level 1	1.00		1.00	
Level 2	1.69 (1.05–1.95) *		1.40 (1.15–1.91) *	
Level 3	1.75 (1.21–2.29) *		1.14 (1.02–1.86) *	
NO <sub>x</sub>				
Level 1	1.00	1.00	1.00	1.00
Level 2	1.67 (1.12–1.94) *	1.83 (1.38–2.53) *	1.33 (1.03–1.75) *	1.50 (1.19–2.03) *
Level 3	1.78 (1.24–2.45) *	1.94 (1.41–2.71) *	1.47 (1.14–1.84) *	1.41 (1.11–1.89) *
O <sub>3</sub>				
Level 1	1.00	1.00	1.00	1.00
Level 2	0.93 (0.64–1.32)	0.95 (0.69–1.16)	1.35 (0.81–1.61)	1.22 (0.81–1.64)
Level 3	0.97 (0.68–1.37)	1.02 (0.70–1.22)	1.27 (0.76–1.70)	0.97 (0.62–1.20)
PM <sub>10</sub>				
Level 1	1.00	1.00	1.00	1.00
Level 2	1.08 (0.75–1.58)	1.14 (0.64–1.32)	0.86 (0.58–1.23)	1.05 (0.72–1.23)
Level 3	1.15 (0.56–1.59)	1.09 (0.42–1.47)	1.12 (0.79–1.45)	1.08 (0.81–1.35)
SO <sub>2</sub>				
Level 1	1.00	1.00	1.00	1.00
Level 2	0.52 (0.36–1.26)	0.74 (0.49–1.16)	1.00 (0.63–1.29)	0.79 (0.61–1.18)
Level 3	0.56 (0.33–1.31)	0.85 (0.68–1.27)	1.05 (0.67–1.33)	0.87 (0.69–1.26)

\* *p* < 0.05; OR, odds ratio; ppm, parts per million; ppb, parts per billion.

#### 4. Discussion

Our finding that the incidence of first AR was higher for boys than girls is consistent with other previous reports [4,10,11]. As indicated by other studies, more severe bronchial responsiveness and/or airway inflammation for boys than girls could explain this finding [24,25]. Lu *et al.* [11] reported an 8.4% prevalence in Chinese preschool children, which was less than our finding (10.9%); however, the ascending trend with age did not differ from ours. They also noted that age-related accumulative personal exposure to air pollutants was significantly associated with AR, suggesting that prevention from exposure to air pollution helps minimize the likelihood of AR occurrence in children.

Our finding of the association between AR and two air pollutants, CO and NO<sub>x</sub>, in preschool children was consistent with the results from previous Taiwanese studies assessing different ages [4,10], suggesting that AR occurrence in children at all ages in Taiwan should be related to outdoor concentrations of CO and NO<sub>x</sub>. Lee *et al.* [4] noted that outdoor NO<sub>x</sub> rather than personal NO<sub>x</sub> was better predictive of AR [26], indicating that NO<sub>x</sub> itself was likely not the direct cause of AR. Hwang *et al.* [10] noted no biological plausibility for CO to influence the airways, suggesting a role for a CO surrogate. Thus, both previous studies considered that these two pollutants served as surrogates for traffic emissions that may have altered the susceptibility of children to allergens. Several studies focused on the effects of traffic emissions by examining the odds ratios for AR or respiratory symptoms with the residential proximity to major roads. A Korean study found that living less than 75 m from a major road was significantly associated with AR for children [15]; a Polish study indicated that seven-year-old children living within 200 m of a major roadway had a significant OR of 1.4 for nasal symptoms compared with those living within 200–500 m, and an even higher OR compared with those living beyond 500 m [17]. Regarding adults, a Swedish survey showed that living within 100 m of a road with a traffic intensity of >10 cars/min was significantly associated with AR (OR = 1.30) [27]. Based on these results, exposure to more traffic emissions may be more likely to lead to AR or other respiratory symptoms at all ages.

One of the traffic emissions of great concern is diesel exhaust particulate (DEP) [28], which has been found to induce the production of Immunoglobulin E (IgE) antibody and oxidative stress in the airways; that is, DEP may enhance allergic inflammation, sensitize the airways to subsequent allergen exposure, and exacerbate allergic symptoms [18,22]. A recent study indicated a significantly positive association between DEP and aeroallergen sensitization for young children less than four years [29], supporting the role of DEP in respiratory allergies. Additionally, DEP may act as an adjuvant to allergens to enhance the allergic response [30]. A Japanese study demonstrated that no IgE antibody response to an AR-causing allergen in Japan was observed from the allergen immunized mice; however, positive results were shown for those immunized with the allergen together with DEP [31].

Other than the association between AR and traffic-related air pollutants, our results also resembled those of previous Taiwanese studies by demonstrating a negative association between O<sub>3</sub> and NO<sub>x</sub> [4,10]. This is not surprising because O<sub>3</sub> is a secondary product derived from the photochemical smog reaction, which consumes NO<sub>x</sub> and hydrocarbons in the presence of sunlight to form O<sub>3</sub> [32]; additionally, Taiwan's location in the sub-tropic and tropic regions means that it is a favorable place for the photochemical smog reaction to take place to produce O<sub>3</sub>. By integrating these three Taiwanese studies, we conclude that the correlation between air pollutants has been consistent since 1994 and the association between AR and outdoor concentrations of CO and NO<sub>x</sub> has been identically significant for preschool children, middle school children and adolescents.

In comparing our data with the environmental data of both previous Taiwanese studies [4,10], we found that the annual concentration of O<sub>3</sub> in Taiwan had gradually increased over the years (reflecting climate change), which resulted in higher temperatures in favor of O<sub>3</sub> production [32]. Despite no significant relationships between AR and O<sub>3</sub> in this study, O<sub>3</sub> is a threat to respiratory health due to its reactivity and irritating effect. However, studies have found a significant association between O<sub>3</sub> and respiratory allergies [7,9,12], suggesting that the impairing effect of O<sub>3</sub> should not be overlooked.

There were several limitations to this study. First, we did not have access to personal information and thus could not assess direct AR-triggering effects; using the identification-free cohort database, however, we were able to determine the impacts of outdoor air pollution on AR. Secondly, environmental monitoring data were derived based on the locations of the clinics and not where the patients actually lived because of a lack of access to personal information. Fortunately, this did not appear to present an issue because most Taiwanese who lived in urban areas were believed to go to nearby clinics for minor respiratory diseases, such as colds and AR. Thirdly, PM<sub>2.5</sub>, one of the major pollutants of traffic emissions, was not included in this study because such data were not measured at all environmental monitoring sites. We expect associations between PM<sub>2.5</sub> and respiratory symptoms to be assessed in the future in Taiwan when PM<sub>2.5</sub> becomes a regular measure for all environmental monitoring sites.

## 5. Conclusions

The first AR incidence for preschool children in Taiwan was higher in males than females for 2007–2011. CO and NO<sub>x</sub>, which are considered to be traffic-related pollutants, were found to be significantly associated with AR in preschool children during the assessed period in Taiwan. The association is consistent with those derived from previous Taiwanese studies focusing on middle school children, suggesting that traffic emissions in Taiwan need to be more effectively controlled to lower the likelihood of AR occurrence.

**Acknowledgments:** This work was supported by Tzu Chi University under Supplement Grant (TCU100184D).

**Author Contributions:** Hui-Ying Chung and Lih-Ming Yiin conceived of the study concept. Chia-Jung Hsieh and Chun-Chieh Tseng contributed to the study design and provided professional recommendations to the manuscript. Hui-Ying Chung also performed the data analysis and wrote the initial draft. Chia-Jung Hsieh, Chun-Chieh Tseng and Lih-Ming Yiin contributed to manuscript revision.

**Conflicts of Interest:** The authors declare no conflicts of interest

## References

1. World Health Organization. Allergic Rhinitis and Sinusitis. Available online: [http://www.who.int/respiratory/other/Rhinitis\\_sinusitis/en/](http://www.who.int/respiratory/other/Rhinitis_sinusitis/en/) (assessed on 25 June 2015).
2. Wang, K.-Y.; Chau, T.-T. An association between air pollution and daily outpatient visits for respiratory disease in a heavy industry area. *PLoS ONE* **2013**, *8*, e75220. [[CrossRef](#)] [[PubMed](#)]
3. Yalcin, A.D.; Basaran, S.; Bisgin, A.; Polat, H.H.; Gorczynski, R.M. Pollen aero allergens and the climate in Mediterranean region and allergen sensitivity in allergic rhinoconjunctivitis and allergic asthma patients. *Med. Sci. Monit.* **2013**, *19*, 102. [[PubMed](#)]
4. Lee, Y.; Shaw, C.; Su, H.; Lai, J.; Ko, Y.; Huang, S.-L.; Sung, F.-C.; Guo, Y. Climate, traffic-related air pollutants and allergic rhinitis prevalence in middle-school children in Taiwan. *Eur. Respir. J.* **2003**, *21*, 964–970. [[CrossRef](#)] [[PubMed](#)]
5. Bousquet, J.; Khaltayev, N.; Cruz, A.; Denburg, J.; Fokkens, W.; Togias, A.; Zuberbier, T.; Baena-Cagnani, C.; Canonica, G.; Van Weel, C. Allergic rhinitis and its impact on asthma (ARIA) 2008. *Allergy* **2008**, *63*, 158–160. [[CrossRef](#)] [[PubMed](#)]
6. Bhattacharyya, N. Air quality influences the prevalence of hay fever and sinusitis. *Laryngoscope* **2009**, *119*, 429–433. [[CrossRef](#)] [[PubMed](#)]
7. Parker, J.D.; Akinbami, L.J.; Woodruff, T.J. Air pollution and childhood respiratory allergies in the United States. *Environ. Health Perspect.* **2009**, *117*, 140–147. [[CrossRef](#)] [[PubMed](#)]
8. Arnedo-Pena, A.; García-Marcos, L.; Urueña, I.C.; Monge, R.B.; Suárez-Varela, M.M.; Canflanca, I.M.; Garrido, J.B.; Quirós, A.B.; Varela, Á.L.-S.; Hernández, G.G. Air pollution and recent symptoms of asthma, allergic rhinitis, and atopic eczema in schoolchildren aged between 6 and 7 years. *Arch. Bronconeumol.* **2009**, *45*, 224–229. [[CrossRef](#)] [[PubMed](#)]
9. Hajat, S.; Haines, A.; Atkinson, R.; Bremner, S.; Anderson, H.; Emberlin, J. Association between air pollution and daily consultations with general practitioners for allergic rhinitis in London, United Kingdom. *Am. J. Epidemiol.* **2001**, *153*, 704–714. [[CrossRef](#)] [[PubMed](#)]

10. Hwang, B.-F.; Jaakkola, J.; Lee, Y.-L.; Lin, Y.-C.; Guo, Y.-L.L. Relation between air pollution and allergic rhinitis in Taiwanese schoolchildren. *Respir. Res.* **2006**, *7*, 23. [[CrossRef](#)] [[PubMed](#)]
11. Lu, C.; Deng, Q.; Ou, C.; Liu, W.; Sundell, J. Effects of ambient air pollution on allergic rhinitis among preschool children in Changsha, China. *Chin. Sci. Bull.* **2013**, *58*, 4252–4258. [[CrossRef](#)]
12. Wong, T.W.; Tam, W.; Yu, I.T.S.; Wun, Y.T.; Wong, A.H.; Wong, C.M. Association between air pollution and general practitioner visits for respiratory diseases in Hong Kong. *Thorax* **2006**, *61*, 585–591. [[CrossRef](#)] [[PubMed](#)]
13. Zhang, F.; Wang, W.; Lv, J.; Krafft, T.; Xu, J. Time-series studies on air pollution and daily outpatient visits for allergic rhinitis in Beijing, China. *Sci. Total Environ.* **2011**, *409*, 2486–2492. [[CrossRef](#)] [[PubMed](#)]
14. Liu, M.-M.; Wang, D.; Zhao, Y.; Liu, Y.-Q.; Huang, M.-M.; Liu, Y.; Sun, J.; Ren, W.-H.; Zhao, Y.-D.; He, Q.-C. Effects of outdoor and indoor air pollution on respiratory health of Chinese children from 50 kindergartens. *J. Epidemiol.* **2013**, *23*, 280–287. [[CrossRef](#)] [[PubMed](#)]
15. Jung, D.-Y.; Leem, J.-H.; Kim, H.-C.; Kim, J.-H.; Hwang, S.-S.; Lee, J.-Y.; Kim, B.-J.; Hong, Y.-C.; Hong, S.-J.; Kwon, H.-J. Effect of traffic-related air pollution on allergic disease: Results of the children's health and environmental research. *Allergy Asthma Immunol. Res.* **2015**, *7*, 359–366. [[CrossRef](#)] [[PubMed](#)]
16. Hansell, A.L.; Rose, N.; Cowie, C.T.; Belousova, E.G.; Bakolis, I.; Ng, K.; Toelle, B.G.; Marks, G.B.; Childhood Asthma Prevention Study Team. Weighted road density and allergic disease in children at high risk of developing asthma. *PLoS ONE* **2014**, *9*, e98978. [[CrossRef](#)] [[PubMed](#)]
17. Porebski, G.; Woźniak, M.; Czarnobilska, E. Residential proximity to major roadways is associated with increased prevalence of allergic respiratory symptoms in children. *Ann. Agric. Environ. Med.* **2014**, *21*, 760–766. [[CrossRef](#)] [[PubMed](#)]
18. Brown, J.; Frew, A. Diesel exhaust particles and respiratory allergy. *Eur. Respir. Mon.* **2002**, *7*, 180–192.
19. Cesaroni, G.; Badaloni, C.; Porta, D.; Forastiere, F.; Perucci, C.A. Comparison between various indices of exposure to traffic-related air pollution and their impact on respiratory health in adults. *Occup. Environ. Med.* **2008**, *65*, 683–690. [[CrossRef](#)] [[PubMed](#)]
20. Myong, J.-P.; Kim, H.; Lee, K.; Chang, S. Time trends of allergic rhinitis and effects of residence on allergic rhinitis in Korea from 1998 through 2007–2009. *Asian Nurs. Res.* **2012**, *6*, 102–106. [[CrossRef](#)] [[PubMed](#)]
21. Braun-Fahrlander, C.; Ackermann-Lieblich, U.; Schwartz, J.; Gnehm, H.P.; Rutishauser, M.; Wanner, H.U. Air pollution and respiratory symptoms in preschool children. *Am. Rev. Respir. Dis.* **1992**, *145*, 42–47. [[CrossRef](#)] [[PubMed](#)]
22. Konishi, S.; Ng, C.F.S.; Stickley, A.; Nishihata, S.; Shinsugi, C.; Ueda, K.; Takami, A.; Watanabe, C. Particulate matter modifies the association between airborne pollen and daily medical consultations for pollinosis in Tokyo. *Sci. Total Environ.* **2014**, *499*, 125–132. [[CrossRef](#)] [[PubMed](#)]
23. Cunningham, W.P.; Cunningham, M.A. *Environmental Science: A Global Concern*; McGraw-Hill Education: New York, NY, USA, 2015; pp. 347–373.
24. Hovland, V.; Riiser, A.; Mowinckel, P.; Carlsen, K.H.; Carlsen, K.C.L. Asthma with allergic comorbidities in adolescence is associated with bronchial responsiveness and airways inflammation. *Pediatr. Allergy Immunol.* **2014**, *25*, 351–359. [[CrossRef](#)] [[PubMed](#)]
25. Zhao, Z.; Huang, C.; Zhang, X.; Xu, F.; Kan, H.; Song, W.; Wieslander, G.; Norback, D. Fractional exhaled nitric oxide in Chinese children with asthma and allergies—A two-city study. *Respir. Med.* **2013**, *107*, 161–171. [[CrossRef](#)] [[PubMed](#)]
26. Krämer, U.; Koch, T.; Ranft, U.; Ring, J.; Behrendt, H. Traffic-related air pollution is associated with atopy in children living in urban areas. *Epidemiology* **2000**, *11*, 64–70. [[CrossRef](#)] [[PubMed](#)]
27. Lindgren, A.; Stroh, E.; Montnémy, P.; Nihlén, U.; Jakobsson, K.; Axmon, A. Traffic-related air pollution associated with prevalence of asthma and COPD/chronic bronchitis. A cross-sectional study in Southern Sweden. *Int. J. Health Geogr.* **2009**, *8*, 2. [[CrossRef](#)] [[PubMed](#)]
28. Peterson, B.; Saxon, A. Global increases in allergic respiratory disease: The possible role of diesel exhaust particles. *Ann. Allergy Asthma Immunol.* **1996**, *77*, 263–270. [[CrossRef](#)]
29. Codispoti, C.D.; LeMasters, G.K.; Levin, L.; Reponen, T.; Ryan, P.H.; Myers, J.M.B.; Villareal, M.; Burkle, J.; Evans, S.; Lockey, J.E. Traffic pollution is associated with early childhood aeroallergen sensitization. *Ann. Allergy Asthma Immunol.* **2015**, *114*, 126–133. [[CrossRef](#)] [[PubMed](#)]
30. Sydbom, A.; Blomberg, A.; Parnia, S.; Stenfors, N.; Sandström, T.; Dahlen, S. Health effects of diesel exhaust emissions. *Eur. Respir. J.* **2001**, *17*, 733–746. [[CrossRef](#)] [[PubMed](#)]



31. Muranaka, M.; Suzuki, S.; Koizumi, K.; Takafuji, S.; Miyamoto, T.; Ikemori, R.; Tokiwa, H. Adjuvant activity of diesel-exhaust particulates for the production of IgE antibody in mice. *J. Allergy Clin. Immunol.* **1986**, *77*, 616–623. [[CrossRef](#)]
32. Kleinman, L.; Lee, Y.-N.; Springston, S.R.; Lee, J.H.; Nunnermacker, L.; Weinstein-Lloyd, J.; Zhou, X. Peroxy radical concentration and ozone formation rate at a rural site in the Southeastern United States. *J. Geophys. Res.* **1995**, *100*, 7263–7273. [[CrossRef](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).