

Dissipation Behavior of Three Fungicides during the Industrial Processing of *Paeoniae Radix Alba* and Associated Processing Factors

Sheng-Nan Li ^{1,†}, Ming-Na Sun ^{3,†}, Fan Wang ¹, Xing Xu ², Xin-Hong Zhang ¹, Jin-Juan Ma ¹, Jin-Jing Xiao ^{1,2}, Min Liao ^{1,2} and Hai-Qun Cao ^{2,3*}

¹ School of Plant Protection, Anhui Agricultural University, Hefei 230036, China;
13210615585@163.com (S.-N.L.); xiaojj187012@163.com (J.-J.X.); liaomin3119@126.com (M.L.);
fxhlrh@163.com(F.W.); majinjuan1024@163.com(J.-J.M.); zxh1060559211@163.com(X.-H.Z.)

² Provincial Key Laboratory for Agri-Food Safety, Anhui Agricultural University, Hefei 230036, China; haiquncao@163.com (H.-Q.C.); xu99xing@163.com (X.X.)

³ Institute of Plant Protection and Agro-Product Safety, Anhui Academy of Agricultural Sciences, Key Laboratory of Agro-Product Safety Risk Evaluation (Hefei), Ministry of Agriculture, Hefei 230031, China; summingna@126.com (M.-N.S.).

* Correspondence: haiquncao@163.com; Tel.: +86-0551-6578-5730; Fax: +86-0551-6578-5730

† These authors contributed equally to this work.

Received: date; Accepted: date; Published: date




Figure S1. Typical chromatograms of pesticides in control (a) and spiked sample (b) with a standard mixture at 50 µg/kg.




Figure S2. Residues of epoxiconazole, difenoconazole, and azoxystrobin in PRA during cooling