



**Table 1.** Rates of health problems among women with hair loss compared to women in the general population in Israel.

	Hair loss	General population: ages 45-54			General population: ages 55-64		
		% (n)	$\chi^2$	RR (95% CI)	% (n)	$\chi^2$	RR (95% CI)
Migraines	60.4% (1501)	11.5% (128) [1]	5847.20***	5.25 (4.45 6.20)	9.6% (81)	7400.73***	6.29 (5.10 7.75)
Diabetes	17.8% (436)	8.1% (39) [2]	311.45***	2.21 (1.61 3.01)	14.2% (63)	26.57***	1.26 (0.98 1.61)
Hypertension	22.3% (541)	15.8% (76) [2]	91.34***	1.42 (1.14 1.76)	31.2% (139)	89.14***	0.72 (0.61 0.84)
Cancer	40.8% (1006)	2.2% (11) [2]	7172.85***	17.90 (9.96 32.17)	8.8% (39)	3253.90***	4.66 (3.44 6.32)

	General population: ages 65+		
	% (n)	$\chi^2$	RR (95% CI)
Migraines	8.6% (49) [1]	8488.18***	6.80 (5.33 9.14)
Diabetes	20.5% (109) [2]	10.61*	0.87 (0.72 1.05)
Hypertension	52.6% (280) [2]	888.78***	0.42 (0.38 0.47)
Cancer	13.6% (72) [2]	1557.12***	3.01 (2.42 3.76)

Note. . \*  $p < .05$ , \*\*\*  $P < .001$ . RR = relative risk; 95% CI = 95% confidence interval. Superscript numbers <sup>1,2</sup> represent the references in the References list from which the data on the general population of women in Israel was extracted: [1] Israeli National Health Interview Survey (INHIS-1), 2003-2004: selected findings. (2006). Israel Center for Disease Control, Ministry of Health; [2] Israel National Health Interview Survey (INHIS-3), 2013-2015: selected findings. (2017). Israel Center for Disease Control, Ministry of Health.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).