




Article

Effect of Different Pollution Parameters and Chemical Components of PM_{2.5} on Health of Residents of Xinxiang City, China

Shuang Wang ¹, Mandeep Kaur ^{2,3}, Tengfei Li ^{2,3}  and Feng Pan ^{1,*}

¹ School of Environment, Henan Normal University, Xinxiang 453007, China; Wshuang730@gmail.com
² Henan Key Laboratory of Earth System Science Observation and Modeling, Henan University, Jinming Campus, Kaifeng 475004, China; mk9041985@gmail.com (M.K.); flyerlee1024@gmail.com (T.L.)
³ College of Environment and Planning, Henan University, Kaifeng 475004, China
* Correspondence: fengpan@htu.edu.cn; Tel.: +86-130-7269-8639

Abstract: The present study was planned to explore the pollution characteristics, health risks, and influence of atmospheric fine particulate matter (PM_{2.5}) and its components on blood routine parameters in a typical industrial city (Xinxiang City) in China. In this study, 102 effective samples 28 (April–May), 19 (July–August), 27 (September–October), 28 (December–January) of PM_{2.5} were collected during different seasons from 2017 to 2018. The water-soluble ions and metal elements in PM_{2.5} were analyzed via ion chromatography and inductively coupled plasma–mass spectrometry. The blood routine physical examination parameters under different polluted weather conditions from January to December 2017 and 2018, the corresponding PM_{2.5} concentration, temperature, and relative humidity during the same period were collected from Second People’s Hospital of Xinxiang during 2017–2018. Risk assessment was carried out using the generalized additive time series model (GAM). It was used to analyze the influence of PM_{2.5} concentration and its components on blood routine indicators of the physical examination population. The “mgcv” package in R.3.5.3 statistical software was used for modeling and analysis and used to perform nonparametric smoothing on meteorological indicators such as temperature and humidity. When Akaike’s information criterion (AIC) value is the smallest, the goodness of fit of the model is the highest. Additionally, the US EPA exposure model was used to evaluate the health risks caused by different heavy metals in PM_{2.5} to the human body through the respiratory pathway, including carcinogenic risk and non-carcinogenic risk. The result showed that the air particulate matter and its chemical components in Xinxiang City were higher in winter as compared to other seasons with an overall trend of winter > spring > autumn > summer. The content of nitrate (NO₃⁻) and sulfate (SO₄²⁻) ions in the atmosphere were higher in winter, which, together with ammonium, constitute the main components of water-soluble ions in PM_{2.5} in Xinxiang City. Source analysis reported that mobile pollution sources (coal combustion emissions, automobile exhaust emissions, and industrial emissions) in Xinxiang City during the winter season contributed more to atmospheric pollution as compared to fixed sources. The results of the risk assessment showed that the non-carcinogenic health risk of heavy metals in fine particulate matter is acceptable to the human body, while among the carcinogenic elements, the order of lifetime carcinogenic risk is arsenic (As) > chromium (Cr) > cadmium (Cd) > cobalt (Co) > nickel (Ni). During periods of haze pollution, the exposure concentration of PM_{2.5} has a certain lag effect on blood routine parameters. On the day when haze pollution occurs, when the daily average concentration of PM_{2.5} rises by 10 µg·m⁻³, hemoglobin (HGB) and platelet count (PLT) increase, respectively, by 9.923% (95% CI, 8.741–11.264) and 0.068% (95% CI, 0.067–0.069). GAM model analysis predicted the maximum effect of PM_{2.5} exposure concentration on red blood cell count (RBC) and PLT was reached when the hysteresis accumulates for 1d (Lag0). The maximum effect of exposure concentration of PM_{2.5} on MONO is reached when the lag accumulation is 3d (Lag2). When the hysteresis accumulates for 6d (Lag5), the exposure concentration of PM_{2.5} has the greatest effect on HGB. The maximum cumulative effect of PM_{2.5} on neutrophil count (NEUT) and lymphocyte (LMY) was strongest when the lag was 2d (Lag1). During periods of moderate to



Citation: Wang, S.; Kaur, M.; Li, T.; Pan, F. Effect of Different Pollution Parameters and Chemical Components of PM_{2.5} on Health of Residents of Xinxiang City, China. *Int. J. Environ. Res. Public Health* **2021**, *18*, 6821. <https://doi.org/10.3390/ijerph18136821>

Academic Editor: Tiziana Schilirò

Received: 6 February 2021

Accepted: 2 June 2021

Published: 25 June 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

severe pollution, the concentration of water-soluble ions and heavy metal elements in PM_{2.5} increases significantly and has a significant correlation with some blood routine indicators.

Keywords: air pollution; PM_{2.5}; water-soluble ions; blood routine parameters; urban air quality; heavy metal

1. Introduction

In recent years, air pollution during haze weather characterized by the presence of high fine particulate matter (PM_{2.5}) has continued to appear in some cities in China, which poses a great threat to the ecological balance and human health [1,2]. It has become one of the primary ecological and social problems in China, and also the focus of heated discussions among many scientists [3]. Worldwide, China is reported to be one of the countries with the highest levels of fine particulate matter (PM_{2.5}) pollution and it is still the primary air pollutant in many cities [4–6]. Many studies have reported heavy metal elements and water-soluble ions as the most crucial components of PM_{2.5} [7,8]. The sources of water-soluble ions are complex and can have a significant impact on weather and visibility, while heavy metal elements are easily adsorbed by PM_{2.5} due to their stable chemical properties. When these chemical components enter the human body through breathing or skin contact, they lead to various functional disorders and alterations that directly affect human health [9].

In China, the Beijing-Tianjin-Hebei region and Yangtze River Delta, among others, have become areas of concern due to the high content of PM_{2.5} in their surroundings. Scientists have become more concerned about the air quality in China and are paying more and more attention to the changing environment. With the emergence of industrialization and rapid development of social economy, the components of air pollution have changed, and unfavorable weather (haze) changes are occurring more frequently. There are very few continuous and periodic studies on PM_{2.5} in the Central Plains urban areas and related cities in China due to a lack of relevant data. Xinxiang City, one of the core cities in the construction of the Central Plains economic zone in China, has become the area of concern for many researchers, but still data on air pollution is mainly based on studies during the heating period and around the Spring Festival in China [10]. Xinxiang City is located at the eastern foot of the Taihang Mountains. It is one of the “transmission channel cities” and is known as the “battery industry capital”. It is of great significance to explore the sources of atmospheric particulate matter and its chemical components and conduct risk assessments. Therefore, more and more research must be conducted and control of air pollution in Xinxiang City needs to be strengthened.

Worldwide, epidemiological studies on particulate matter and their effect on human health began in the late 1880s. These studies focused on long-term and short-term exposure to particulate matter and health indicators, such as visiting rate, vital capacity, incidence rate of respiratory system, exacerbation of disease, and mortality [11,12]. Reports on the health risks and damage value during heavy haze pollution showed that acute bronchitis and asthma are the main diseases caused by PM_{2.5}, accounting for more than 90% [13]. According to Zhu Yidan and others’ investigations on the relationship between respiratory diseases and symptoms of school students in different areas of Beijing who experience varying air pollution levels, it was found that the incidence of children’s persistent cough being more serious in haze areas, and children in different air-pollution-affected areas suffered from respiratory illness [14]. However, the current data on smog and human health are limited to epidemiological surveys (diagnosis rate, incidence, disease types, etc.), there is no specific clinical diagnosis reference data for the response of various human organs to haze, which can be used as clinical diagnosis and treatment data, such as the blood routine and C reactive protein of the human body exposed to PM_{2.5}. The physiological index is one of the commonly used standards in different pathology and toxicology studies

to indicate the health status of any organism [11,14–16]. Medium- and long-term exposure to air pollution may lead to changes in some indexes. A review of the literature revealed that $PM_{2.5}$ has a toxic effect mainly in terms of oxidative damage, to a certain extent, which can cause inflammatory and oxidative stress reactions in the lungs of rats and increase the susceptibility of lung tissue to bacteria [17]. Therefore, it can be inferred that the corresponding examination data will change when the human body responds to such inflammatory responses. The evidence of the influence of atmospheric pollutants on peripheral blood indicators in the population is still limited. In this study, the blood routine indexes of people exposed to different environmental conditions were analyzed; warning people that increased $PM_{2.5}$ individual exposure can cause changes in multiple blood routine indexes with lag and cumulative effects, which can provide a reference for health management and disease prevention of people under haze pollution. Based on this research background, it will be of practical significance to understand the characteristics of air pollutants and evaluate their health risks.

Considering all aspects in mind, the present study area, Xinxiang City, was selected as an investigation area due to a lack of information on such studies. The specific goals of the present research were to:

- I. explore the pollution characteristics and pollution sources of different chemical components in fine particles and evaluate the environmental ecological risk of heavy metals in $PM_{2.5}$ based on the health risk assessment model,
- II. investigate the capability of a generalized additive model (GAM) to explore the effect of $PM_{2.5}$ exposure concentration on the blood routine parameters of the population during periods of haze pollution.

2. Material and Methods

2.1. Sample Collection

Data Sources

(1) Atmospheric fine particulate matter ($PM_{2.5}$) data, synchronized hourly values, and meteorological data during the study period were obtained from the national air quality network monitoring and management platform.

(2) The monitoring and analysis of $PM_{2.5}$ and its chemical components were carried out in strict accordance with the “technical specification for manual monitoring method (gravimetric method) of ambient air particulate matter ($PM_{2.5}$)” and “technical method guide for source apportionment monitoring of ambient air particulate matter (Trial)”. The time of each sampling was started in morning at 10 a.m. and samples were collected the next day at 9 a.m. The data of metal elements and water-soluble ions in $PM_{2.5}$ were determined via inductively coupled plasma–mass spectrometry (ICP-MS) and ion chromatography (IC).

(3) The medical examination data of a normal population in Xinxiang was collected from the Second People’s Hospital of Xinxiang in 2017–18. The routine blood examination parameters of healthy people were obtained from January to December 2017, during different polluted weather conditions, including white blood cell count (WBC), red blood cell count (RBC), hemoglobin concentration (HGB), platelet count (PLT), neutral ratio (NEUT%), lymphatic ratio (LMY%), and monocyte ratio (MONO%).

2.2. Experimental Instruments

The main instruments used in the study were an intelligent medium flow sampler (TH-150) for collection of $PM_{2.5}$, an ion chromatograph (ICS-1100) for determination of water-soluble ions, and an inductively coupled plasma–mass spectrometer (7500a, Agilent) for heavy metal determination.

2.3. Sampling Time and Location

The sampling site was set on the top of the Xinxiang City Environmental Protection Monitoring Station (E 113°53′01″, N 35°18′11″), about 15 m above the ground mainly surrounded by the intersection of life, scientific research culture, and commercial areas

without industrial sources of pollution (Figure 1). The sampling was done for 23 h a day (10 a.m. to 9 a.m.). During the whole sampling period, the sampling time, atmospheric pressure, average temperature, weather conditions, sample volume and other related parameters were recorded, and bad weather such as rain and snow was avoided as much as possible. After sampling, the sample film was placed in a refrigerator frozen layer and stored away from light for analysis. After removing the external factors such as rain and snow, weather, and machine failure, 102 effective valid samples were actually collected. Among them, 28 samples were collected in the spring season from April 2017 to May 2017, 19 samples in the summer season from July 2017 to August 2017, 27 effective samples were collected in autumn from September 2017 to October 2017, and 28 valid samples were collected in the winter season from December 2017 to January 2018. The collection, transportation and storage of samples was done strictly according to the technical specification for the manual monitoring method (gravimetric method) of ambient air particulate matter [18]. A Wuhan Tianhong TH-150c series intelligent flow sampler was used for sampling and the sampling flow was 100 L/min. The fine particulate particles were collected on a quartz filter (7201-03, PALL) with a diameter of 90 mm (the effective diameter of the sampler was 80 mm) and stored for further analysis.



Figure 1. Location of the sampling site in Xinxiang, Henan, China.

2.4. Statistical Methods

2.4.1. Principal Component Analysis (PCA)

PCA (principal component analysis) is a source analysis method advocated and approved by the U.S. Environmental Protection Agency. Its main purpose is to reduce the dimension of a group of variables. This method can synthesize and simplify high-dimensional variables on the basis of keeping the original data information to the utmost extent, so it has certain advantages and has been widely used in the environmental field. The absolute principal component score is a quantitative estimation of the source's share of

each pollutant through the principal component factor score. Specifically, it is divided into the following steps: first, standardize the data, as shown in Formula (1).

$$Z_{ik} = \frac{C_{ik} - C_i}{\sigma_i} (i = 1, 2, \dots, m; k = 1, 2, \dots, n) \quad (1)$$

In the formula, k is the standardized concentration value (dimensionless), C_{ik} is the concentration of compound i in the k -th observation, and C_i is the average value of the formula of C_{ik} ; σ_i is the standard deviation. The basic formula of PCA is as follows:

$$Z_{ik} = \sum_{j=1}^p W_{ij} P_{jk} (i = 1, 2, \dots, m; j = 1, 2, \dots, p; k = 1, 2, \dots, n) \quad (2)$$

In the formula, W_{ij} is the factor load (dimensionless), which represents the correlation coefficient between compound i and the factor j obtained via principal component analysis; P_{jk} is the factor score (dimensionless) [19,20].

The principal component analysis of metal elements and water-soluble ions in PM_{2.5} collected from Xinxiang City was carried out by using IBM SPSS Statistics (SPSS Inc.; Chicago, IL, USA) (September 2006) statistical software. The orthogonal rotation method with Kaiser Standardization was adopted, and the rotation converged after four iterations.

2.4.2. Health Risk Assessment Model

Heavy metal elements enter the human body mainly through the mouth, respiratory pathway, and skin exposure pathway, thus causing harm to the human body. Of these three routes, inhalation through the respiratory system poses the greatest health risk to the human body [9]. The US EPA exposure model was used to evaluate the health risks caused by heavy metals in PM_{2.5} to the human body through the respiratory pathway, which mainly included carcinogenic risk and non-carcinogenic risk.

The formula used for calculating the daily average exposure dose for non-carcinogens (average daily dose, ADD, mg/(kg × d)) and the lifetime average exposure dose for carcinogens (life average daily dose, LADD, mg/(kg × d)) is as follows:

$$\text{ADD (LADD)} = C \times \text{IR} \times \text{EF} \times \text{ED} / (\text{AT} \times \text{BW}) \quad (3)$$

where, C is the concentration of heavy metal elements in the air, (mg/m³); IR is respiratory rate, (m³/d); EF is the exposure frequency, (d/a); ED is the duration of exposure (a); BW is body weight (kg); and AT is the average exposure time (d). Among them, the exposure parameters were selected from the relevant parameters in the Chinese population exposure parameters manual (adult volume) released by the Ministry of Environmental Protection [21] to make the health risk assessment more in line with the characteristics of Chinese people.

The non-carcinogenic risk of a single heavy metal element was indicated by HQ (hazard quotient), which was used to evaluate the health risk. The total non-carcinogenic risk is expressed as HI (hazard index), representing the cumulative risk of multiple contaminants. The lifetime carcinogenic risk of carcinogenic heavy metals was characterized by ILCR (incremental lifetime cancer risk). The formula used for calculation is as follows:

$$\text{HI} = \sum_{i=1}^n \text{HQ}_i = \sum_{i=1}^n (\text{ADD}_i / \text{RfD}_i) \quad (4)$$

$$\text{ILCR} = \text{LADD} \times \text{SF} \quad (5)$$

where, RfD is the reference dose (mg/(kg × d)), and SF is the carcinogenic slope factor ((kg × d)/mg). When HI or HQ are ≤ 1 , the risk of non-carcinogenic health is low and can be ignored, when HQ is > 1 , there is a non-carcinogenic risk, when ILCR is $< 10^{-6}$, the carcinogenic risk is negligible; if the ILCR is between 10^{-6} and 10^{-4} , the heavy metal

element is considered to have no carcinogenic risk, whereas when ILCR is $>10^{-4}$, there is a higher carcinogenic risk.

2.4.3. Risk Assessment

A semi-parametric generalized additive model (GAM) was used. The model was validated before use. The “mgcv” package in R.3.5.3 statistical software was used for modeling and analysis and a smooth spline function was used to perform nonparametric smoothing on meteorological indicators such as temperature and humidity. The PM_{2.5} concentration value during periods of haze pollution from 1 January 2017 to 31 December 2017 was introduced into the model as a linear variable to observe its impact on different blood routine indicators. The formula is as follows:

$$\text{“Log E” (“Yi”) = } \beta \text{ “PM}_{2.5} \text{” + s (“time, df”) + DOW (week) + s (“temperature, df”) + s (humidity, df) + } \alpha \text{ (6)}$$

where E(Y_i) is the expected value of a certain blood routine index in the physical examination population on the first day, β is the regression coefficient, s is the non-parametric spline smoothing function, df is degree of freedom, time is the time series, DOW is used to adjust the day of the week effects, (temperature, df) is the temperature spline smoothing function, s (humidity, df) is the relative humidity spline smoothing function, and α is the intercept. Time, temperature and humidity are adjusted by the natural smoothing (S) function. According to Akaike’s information criterion (AIC) [17], the optimal model is selected. When the AIC value is the smallest, the goodness of fit of the model is the highest, and the df is the best degree of freedom. Test level $\alpha = 0.05$.

The regression coefficient β was estimated according to the model, and the corresponding percentage change and 95% confidence interval (CI) of each blood routine index was calculated when the PM_{2.5} concentration increased by $10 \mu\text{g}\cdot\text{m}^{-3}$. In the aspect of lag effect analysis, the air pollution effect with a lag of 0–7d was selected. The regression coefficient β estimated by the model was used to calculate the RR of different lag days, and the best lag period was determined according to the maximum relative risk value.

3. Results and Discussion

3.1. Pollution Characteristics and Source Analysis of Chemical Components in PM_{2.5}

3.1.1. Concentration of PM_{2.5} in Xinxiang City

The change in mass concentration of atmospheric fine particulate matter (PM_{2.5}) during different sampling periods is shown in Figure 2.




Figure 2. Regularity of mass concentration of PM_{2.5} during different sampling periods in Xinxiang city, China.

The mass concentration of atmospheric PM_{2.5} during the sampling period was ranged from $8.92 \mu\text{g}\cdot\text{m}^{-3}$ to $242.37 \mu\text{g}\cdot\text{m}^{-3}$, with a daily average mass concentration of $63.26 \mu\text{g}\cdot\text{m}^{-3}$.

Based on the concentration limits of various air pollutants specified in the latest “ambient air quality standards” (GB 3085–2012), it does not exceed the national secondary concentration limit ($75 \mu\text{g}\cdot\text{m}^{-3}$) but it was 1.66 times the national first-level concentration standard ($35 \mu\text{g}\cdot\text{m}^{-3}$).

During the sampling period, the maximum daily average concentration of $\text{PM}_{2.5}$ was reported in December while the minimum concentration was observed in July. The seasonal variation regular pattern was winter ($96.77 \mu\text{g}\cdot\text{m}^{-3}$) > spring ($52.77 \mu\text{g}\cdot\text{m}^{-3}$) > autumn ($51.30 \mu\text{g}\cdot\text{m}^{-3}$) > summer ($46.32 \mu\text{g}\cdot\text{m}^{-3}$) (Figure 3). According to the regulation of ambient air quality index (AQI), the analysis results show that the $\text{PM}_{2.5}$ concentration level in winter (December) was the highest, and heavy pollution weather accounted for 17.9% of the sampling days in winter; the $\text{PM}_{2.5}$ concentration level in summer (July) was the lowest, and fine weather accounted for 89.47% of the total sampling days in summer. Combined with the geographical location and climate characteristics of city, this situation may be due to low temperature and dry weather during winter and the atmospheric boundary layer tends to be stable which was not conducive to the diffusion of air pollutants [22]. The trend of the change of summer low, winter high is consistent with the annual monitoring results of atmospheric particles. Poor diffusion conditions in winter and high pollutant emissions in the heating season are the reasons for the high concentration of PM in winter [23–26], while meteorological factors in summer (large precipitation, high temperature, strong radiation forcing and high atmospheric boundary layer, the atmospheric vertical diffusion is the most favorable) are the important factors for the low concentration of PM [27].




Figure 3. Metal element concentrations in $\text{PM}_{2.5}$ samples collected from Xinxiang city. Al, aluminum; Na, sodium; Ca, calcium; Mg, magnesium; K, potassium; As, arsenic; Fe, iron; Zn, zinc; Cr, chromium; Cu, copper; Pb, lead; Mn, manganese; V, vanadium; Ni, nickel; Co, cobalt; Ag, silver; Cd, cadmium.

Globally, $\text{PM}_{2.5}$ emissions have slowly declined since 1996, but the $\text{PM}_{2.5}$ concentration of weighted population exposure has continued to increase slowly since 1990 [28,29]. As of 2016, the living environment of only 27% of the population in the United States, Canada, Australia, and a few other countries in the world meets the IT-3 standard of the WHO’s air quality guidelines (annual average $\text{PM}_{2.5} < 10 \mu\text{g}\cdot\text{m}^{-3}$), while there is still 44% of the population whose living environment cannot meet the IT-1 standard of the lowest particulate matter concentration (annual average $\text{PM}_{2.5} < 35 \mu\text{g}\cdot\text{m}^{-3}$) [30]. In many Asian countries, haze weather with low visibility and a high concentration of $\text{PM}_{2.5}$ as the main feature frequently occurs [30]. The weighted average annual $\text{PM}_{2.5}$ exposure concentrations of the populations of Bangladesh, Nepal, India, and Pakistan has increased rapidly since 2010, reaching $101 \mu\text{g}\cdot\text{m}^{-3}$, $78 \mu\text{g}\cdot\text{m}^{-3}$, $76 \mu\text{g}\cdot\text{m}^{-3}$, and $76 \mu\text{g}\cdot\text{m}^{-3}$, respectively, in 2017, which are far higher than China’s level ($56 \mu\text{g}\cdot\text{m}^{-3}$) [30]. Thailand, Indonesia, and Malaysia are also facing haze due to large-scale biomass combustion [31,32]. With the promulgation of the Air Pollution Prevention and Control Action Plan (2013–2017), the air quality in most

areas in China has improved considerably, but the average annual $PM_{2.5}$ concentration in 190 cities in China from 2014 to 2015 is still as high as $61 \mu\text{g}\cdot\text{m}^{-3}$, which is six times the WHO IT-3 standard [33]. The concentration of $PM_{2.5}$ in Beijing during haze in January 2013 was as high as $786 \mu\text{g}\cdot\text{m}^{-3}$, affecting nearly 600 million people in the Beijing-Tianjin-Hebei region, Yangtze River Delta, Pearl River Delta, and other places. During the haze event in December 2013, the average daily $PM_{2.5}$ in Shanghai was as high as $416 \mu\text{g}\cdot\text{m}^{-3}$, and the instantaneous maximum $PM_{2.5}$ concentration in Nanjing exceeded $900 \mu\text{g}\cdot\text{m}^{-3}$ [34].

3.1.2. Pollution Characteristics

The metal elements in $PM_{2.5}$ samples collected during the sampling period were determined and the results are shown in the Figure 3. The concentration of 18 metal elements reported in $PM_{2.5}$ collected from Xinxiang city follow an order (high to low) as $\text{Al} > \text{Ca} > \text{Mg} > \text{K} > \text{Na} > \text{As} > \text{Fe} > \text{Zn} > \text{Cr} > \text{Mn} > \text{Pb} > \text{Cu} > \text{V} > \text{Ag} > \text{Ni} > \text{Cd} > \text{Co} > \text{Be}$. Metallic elements, including Al, Na, Ca, Mg, K, As, Fe, and Zn, showed high concentrations while Cd, Ag, V, Co, Ni, and Pb were found in low concentrations. In addition to crustal elements, the concentrations of heavy metals such as Zn, As, Cr, and Mn were higher in the measured metal elements.

The difference in weather conditions and human activities in different seasons will lead to a change of trace metal elements in atmospheric fine particles. The contents of Al and Ca were the highest in winter (heating period), were $1.63 \mu\text{g}\cdot\text{m}^{-3}$ and $0.72 \mu\text{g}\cdot\text{m}^{-3}$, respectively. This is consistent with the comparison of metal element content in winter $PM_{2.5}$ in Beijing [35], Guangzhou [36], and Henan [37]. Qiao Baowen et al. [38] stated that Al and Ca are the identification elements of crustal source and construction dust, and the ratio of Ca/Al is used to judge the degree of influence of construction activities in a certain place. The ratio of Ca/Al in Xinxiang City was about 0.44, which is greater than the soil background value of Henan Province (0.38), indicating that construction is still frequent in winter, and construction dust has a certain contribution to the formation of smog. Seasonal analysis showed that Ca, Mg, K, and Fe in $PM_{2.5}$ were higher in winter and spring than in summer and autumn. This may be due to the higher wind speed in winter and spring in Xinxiang City, which makes the mass concentration of crustal elements in $PM_{2.5}$ increase. In addition, most of the metal elements were low in summer which may be due to the higher air humidity in summer and autumn and more rainfall than in spring and winter, which plays a role in the deposition of metal elements. Compared with Beijing [38], Tianjin [39], Nanjing [40], and Handan [41], the concentration of As in Xinxiang City was higher, and the content of As in the main cities of our country exceeds the concentration limit specified by the WHO ($6 \text{ ng}\cdot\text{m}^{-3}$), which reflects the phenomenon that the air pollution of As in our country generally exceeds the standard, which should be paid attention to.

Water soluble ions are an important component of atmospheric fine particles ($PM_{2.5}$), which can reduce the visibility of the atmosphere and also have an important impact on the pH of air particles [42,43]. The total concentration levels of eight water-soluble ions (from high to low) were observed as $\text{SO}_4^{2-} > \text{NO}_3^- > \text{NH}_4^+ > \text{Ca}^{2+} > \text{Cl}^- > \text{K}^+ > \text{Na}^+ > \text{Mg}^{2+}$ and the proportion of water-soluble ions concentration to total ion concentration was 34.46%, 31.32%, 16.4%, 7.73%, 6.01%, 2.6%, 0.95%, and 0.53% (Figure 4).

The sulphates (SO_4^{2-}), nitrates (NO_3^-), and ammonia (NH_4^+) ions are also called SNA, where S denotes sulphates, N-nitrates, and A-ammonia. Many studies have shown that SNA is an important part of atmospheric $PM_{2.5}$ (accounting for more than 70%) in some urban areas in China. SNA also acts as an important factor that can contribute to declining atmospheric visibility of urban areas. The concentration level of water-soluble ions in $PM_{2.5}$ in Xinxiang City was compared with some of domestic and foreign cities (Figure 5).




Figure 4. Concentration and proportion of some water-soluble ions in PM_{2.5}. Na⁺, sodium ion; NH₄⁺, ammonium ion; K⁺, potassium ion; Mg²⁺, magnesium ion; Ca²⁺, calcium ion; Cl⁻, chloride ion; NO₃⁻, nitrate; SO₄²⁻, sulfate.




Figure 5. Concentrations of water-soluble ions in PM_{2.5} in different cities and seasons.

It can be observed that compared with some European cities, such as Poland and Greece [44–46], the concentrations of water-soluble ions in PM_{2.5} are generally higher in some Asian cities, including Beijing [47], Shijiazhuang [48], Nanjing [49], and Seoul [50].

This phenomenon shows that the air pollution level in Asia is more serious than that in Europe. When compared with other Chinese cities, the concentration of total water-soluble ions in PM_{2.5} in Xinxiang City is lower but with higher concentration levels of NO₃⁻ and SO₄²⁻, which is close to Shijiazhuang, Taiyuan, and other cities. This indicates that the air condition may be greatly affected by human activities, and the concentration of secondary ions in particulate matter is high. The ratio of NO₃⁻ and SO₄²⁻ in atmospheric particles can evaluate the impact of fixed and mobile sources on the atmospheric environment [51]. When (NO₃⁻)/(SO₄²⁻) > 1, it indicates that the influence of mobile pollution sources is higher than that of fixed pollution sources; when (NO₃⁻)/(SO₄²⁻) < 1, it indicates that the contribution of fixed pollution sources is greater than that of mobile pollution sources. During the sampling period, the ratio of NO₃⁻ to SO₄²⁻ varied from 0.10 to 2.56, with the lowest ratio in summer and the highest ratio in winter, with an average of 1.81, indicating that the contribution rate of mobile pollution sources in winter in Xinxiang City is higher than that of fixed pollution sources.

3.1.3. Source Resolution

The source inventory method, diffusion model, and receptor model are the three mainstream methods currently used in source analysis research. Since the subject of this study is air pollutants, the PMF source analysis receptor model is greatly affected by meteorological conditions, the receptor data needs are large, and the calculation is large and complicated. Therefore, due to the small number of samples in this study, after comprehensive consideration and comparison of multiple source analysis models (PCA, PMF, CMB), the principal component analysis method was selected to analyze the emission sources of $PM_{2.5}$ in Xinxiang City in order to explore the main emission sources of air pollution.

The main factor analysis method (Figure 6) is a method to reduce the dimension of variables to facilitate description, understanding, and analysis. It is a common multivariate statistical analysis method for qualitatively identifying the main pollution sources of atmospheric particulate matter. According to the analysis result of the maximum variance rotation factors, factor 1 depicted the elements with a higher load coefficient, which included Cu, Cr, Ni, and V. One of the earlier studies by Pulles et al. [52] reported a high content of Cu in the exhaust gas of gasoline vehicles while high Ni content came from sources like coal-burning, fuel oil, and nickel ore smelting. Thus, factor 1 (16.64%) is considered as the mixing source of motor vehicles and metallurgy and chemical industry. The high load coefficient for factor 2 (14.34%) are mainly secondary water-soluble ions, including NH_4^+ , NO_3^- , and SO_4^{2-} . These secondary inorganic ions in the atmosphere are mainly derived from gaseous precursors generated by human activities. The water-soluble ions, including K^+ and Cl^- , showed a high load coefficient of factor 3 (13.41%). In addition to this, the metal elements Mn and Cd also showed a high load coefficient, among which Mn is a trace element produced in the combustion of coal, while Cd is derived mainly from garbage incineration [53], indicating that the source of pollution may be biomass combustion. The elements with a high load coefficient of factor 4 (12.36%) are Na, Mg, Ca, Ca^{2+} , Mg^{2+} , and As. Since Ca and Mg are crustal elements mostly from ground dust [54,55], while As is a major component in the coking production process often high in metallurgical and chemical dust. Therefore, factor 4 is considered to be the industrial source and crustal source. Since Zn is mainly derived from metal smelting and industrial emissions, coal combustion is the main source besides the non-ferrous metallurgical industry, which is consistent with the environmental conditions of large-scale coal-fired heating in Xinxiang in winter. The main anthropogenic source of Co is coal combustion, so the pollution source for factor 5 (4.83%) may represent coal combustion.

3.2. Health Risk Assessment of Heavy Metal Elements

According to the EPA Integrated Risk Information Database (IRIS) and the International Cancer Institute (IARC), pollutants can be divided into two categories, carcinogens and non-carcinogens. At present, the domestic health risk assessment is mainly based on the soil health risk assessment model proposed by the US Environmental Protection Agency [56] and some of the parameters have been modified for use [57] (Table 1). The present study mainly considers the respiratory pathway to evaluate the risk of exposure of heavy metals in $PM_{2.5}$ to human health. Based on the available data and the results of source analysis, the carcinogenic metal elements studied include Cr, Co, Ni, As, and Cd and non-carcinogenic elements included Pb, Cu, V, Mn, and Zn.




Figure 6. Diagram of maximum variance rotation factor analysis. Na⁺, sodium ion; NH₄⁺, ammonium ion; K⁺, potassium ion; Mg²⁺, magnesium ion; Ca²⁺, calcium ion; Cl⁻, chloride ion; NO₃⁻, nitrate; SO₄²⁻, sulfate; Cu, copper; Pb, lead; Cr, chromium; Ni, nickel; Cd, cadmium; Be, beryllium; Na, sodium; Mg, magnesium; Al, aluminum; K, potassium; Ca, calcium; V, vanadium; Mn, manganese; Fe, iron; Co, cobalt; Zn, zinc; As, arsenic; Ag, silver.

Table 1. Related parameters of population exposure evaluation model.

| Parameter | Different People | | |
|-------------------------|------------------|--------------|----------|
| | Adult Male | Adult Female | Children |
| IR (m ³ /d) | 17.7 | 14.3 | 8.7 |
| EF (d/a) | 365 | 365 | 365 |
| ED (a) | 30 | 30 | 10 |
| BW (kg) | 64 | 54.4 | 16 |
| AT (carcinogenic)/d | 70 × 365 | 70 × 365 | 70 × 365 |
| AT (non-carcinogenic)/d | 30 × 365 | 30 × 365 | 10 × 365 |

IR; respiratory rate; EF; exposure frequency; ED; duration of exposure; BW; body weight; AT; average exposure time.

SPSS statistical software was used to calculate the 95% UCL of the mass concentration of ten heavy metal elements in PM_{2.5}, and then the non-carcinogenic risk and lifetime carcinogenic risk were calculated. The health risk evaluation results are shown in Table 2 and the relevant data for SF and RfD values of each heavy metal element are shown in Table 3.

As can be seen from the table, the non-carcinogenic risk (HQ) produced by 10 heavy metals in PM_{2.5} to different groups of people was less than 1.0 and the total non-carcinogenic risk (HI) of adults and children was also less than 1.0, which indicated that the non-carcinogenic health risk of heavy metals in PM_{2.5} in Xinxiang was low and within the acceptable range of human body. Among all metals, the non-carcinogenic health risks of Mn, Co, and Cr are higher and the risk in adults and children follows a trend of Mn > Cr > Co with more occurrence in children followed by adult males and adult females. The long-term exposure to Mn has been reported to cause corresponding neurological symptoms [58] while Cd and Mn mostly originate from transportation and industrial exhaust emissions.

Table 2. Health risk assessment table for 10 heavy metal elements in PM_{2.5}.

| Element | HQ | | | ILCR | | |
|---------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| | Adult Male | Adult Female | Children | Adult Male | Adult Female | Children |
| Cu | 3.5×10^{-5} | 3.3×10^{-5} | 6.9×10^{-5} | | | |
| Pb | 6.9×10^{-4} | 6.6×10^{-4} | 1.4×10^{-3} | | | |
| Zn | 4.6×10^{-5} | 4.3×10^{-5} | 8.9×10^{-5} | | | |
| V | 1.2×10^{-4} | 1.1×10^{-4} | 2.3×10^{-4} | | | |
| Mn | 1.7×10^{-1} | 1.6×10^{-1} | 3.3×10^{-1} | | | |
| Co | 0.3×10^{-1} | 0.1×10^{-1} | 0.1×10^{-1} | 1.5×10^{-6} | 6.8×10^{-7} | 4.7×10^{-7} |
| As | 7.6×10^{-5} | 3.4×10^{-5} | 2.3×10^{-5} | 3.4×10^{-4} | 1.5×10^{-4} | 1.1×10^{-4} |
| Cr | 1.6×10^{-1} | 7.2×10^{-2} | 0.5×10^{-1} | 2.0×10^{-4} | 8.7×10^{-5} | 6.0×10^{-5} |
| Ni | 1.3×10^{-5} | 0.6×10^{-5} | 0.4×10^{-5} | 2.3×10^{-7} | 1.0×10^{-7} | 0.7×10^{-7} |
| Cd | 2.7×10^{-4} | 1.2×10^{-4} | 0.8×10^{-4} | 1.7×10^{-6} | 7.6×10^{-7} | 5.2×10^{-7} |
| HI | <1 | <1 | <1 | | | |

HQ, non-carcinogenic risk (hazard quotient); HI, total non-carcinogenic risk (hazard index); ILCR, lifetime carcinogenic risk values of carcinogenic heavy metals (incremental lifetime cancer risk); Cu, copper; Pb, lead; Zn, zinc; V, vanadium; Mn, manganese; Co, cobalt; As, arsenic; Cr, chromium; Ni, nickel; Cd, cadmium.

Table 3. Dose-response parameters for heavy metal elements entering the body through respiration.

| Classification | Element | RfD | SF |
|-------------------------------|---------|-----------------------|-------|
| Carcinogenic heavy metals | Cr | 2.86×10^{-5} | 42.0 |
| | Co | 5.71×10^{-6} | 9.80 |
| | Ni | 2.06×10^{-2} | 0.84 |
| | As | 0.3 | 15.10 |
| | Cd | 1.0×10^{-3} | 6.30 |
| Non-carcinogenic heavy metals | Pb | 3.52×10^{-3} | |
| | Cu | 4.02×10^{-2} | |
| | V | 7.0×10^{-3} | |
| | Mn | 1.43×10^{-5} | |
| | Zn | 0.3 | |

RfD; reference dose (mg/(kg × d)); SF; carcinogenic slope factor ((kg × d)/mg); Cr, chromium; Co, cobalt; Ni, nickel; As, arsenic; Cd, cadmium; Pb, lead; Cu, copper; V, vanadium; Mn, manganese; Zn, zinc.

Among carcinogenic elements, the sequence of lifetime carcinogenic risk follows a trend of As > Cr > Cd > Co > Ni, in which As and Cr were two orders of magnitude higher than Cd and Co, and three orders of magnitude higher than Ni. The carcinogenic health risk value of As in adults and children was higher than 10^{-4} , indicating that there was a relatively obvious carcinogenic risk to humans. The health risk value of Cr was higher than 10^{-4} in adult males and between 10^{-6} and 10^{-4} in adult females and children, indicating that there was a significant carcinogenic risk for adult males, while there may be carcinogenic risk for adult females and children. Co and Cd showed a health risk value ranged between 10^{-6} and 10^{-4} in adult men, indicating that there may be a certain risk of carcinogenesis, but in adult women and children it was less than the risk threshold of 10^{-6} , indicating that there was no carcinogenicity risk. The health risk value of Ni in adults and children was below the threshold of 10^{-6} , indicating that there was no health risk. In addition, we must also consider the fact that metal exposure also occurs through ingestion and skin contact. If these pathways are considered, the risk may be higher [59].

3.3. Correlation Analysis of PM_{2.5} Exposure Level and Blood Routine Parameters of Healthy People

3.3.1. Pollution Lag Risk Assessment

According to the preliminary collection of the physical examination data of the normal physical examination population in the health management department of a hospital in Xinxiang City, 357 physical examination population data were collected during severe pollution periods (AQI ≥ 150; PM_{2.5} ≥ 115 μg/m³) in winter (December), and 354 physical examination population data were collected during fine weather periods (AQI ≤ 100) in summer (July). The measurement data were expressed as mean ± standard deviation (x ± s),

and a t test was performed; $p < 0.05$ indicated that the difference was statistically significant. Preliminary comparative analysis found that during moderate and severe pollution periods, the number of white blood cells, red blood cells, hemoglobin concentration, and neutrophil ratios decreased significantly, while lymphatic ratios showed an increase (Table 4).

Table 4. Comparison of blood routine indicators between non-haze periods and haze periods.

| Blood Routine Index | Haze Group (n = 357) | Non-Haze Group (n = 354) | p (sig.) |
|---------------------|-------------------------|-----------------------------|----------|
| WBC | 5.80 ± 1.52 | 6.25 ± 1.55 | 0.001 |
| MONO (%) | 3.77 ± 1.16 | 3.84 ± 1.03 | 0.393 |
| RBC | 4.60 ± 0.47 | 4.72 ± 0.46 | 0.001 |
| LMY (%) | 35.31 ± 8.27 | 33.68 ± 7.77 | 0.006 |
| HGB | 134.11 ± 16.44 | 140.01 ± 16.05 | 0.001 |
| PLT | 222.08 ± 5.98 | 222.68 ± 48.07 | 0.513 |
| NEUT (%) | 58.82 ± 8.45 | 60.56 ± 8.29 | 0.007 |

Measurement data is expressed as mean ± standard deviation ($x \pm s$). $p < 0.05$ indicates that the difference is statistically significant using t test. WBC, white blood cell count; MONO (%), monocyte ratio; RBC, red blood cell count; LMY (%), lymphatic ratio; HGB (g/L), hemoglobin; PLT, platelet count; NEUT (%), neutrophil ratio.

The changes in all these indicators predicted that under different weather conditions, the body's specific cell population would have a different reactive response to foreign stimuli, indicating that the human body is at risk of infection when haze pollution occurs. Among all parameters, decrease in hemoglobin concentration was the most obvious, indicating decreased immunity in the population was significant in the polluted weather.

The comparative analysis of the study found that when polluted weather occurs, the routine indicators of human blood would not produce a fixed change trend on the day of clean weather compared to the previous day. Therefore, it is speculated that the blood routine parameters would not change immediately in a day when haze pollution occurs. In order to further explore the hysteresis effect of polluted weather on the blood routine parameters of the population, one must retrospectively collect blood routine physical examination parameters of healthy people under different polluted weather conditions from January to December in 2017, including white blood cell count (WBC), red blood cell count (RBC), hemoglobin concentration (hemoglobin, HGB), platelet count (platelet, PLT), neutral ratio (neutrophil ratio, NEUT%), lymphocyte ratio (LMY%), mononuclear ratio (monocyte ratio, MONO%). After adjusting the temperature, humidity, day of week effects, and other confounding factors, the modeling analysis found that on the day of haze, the average daily exposure concentration of $PM_{2.5}$ was significantly correlated with RBC and PLT ($p < 0.05$), but no significant correlation was observed between $PM_{2.5}$ concentration and the other five indicators (Table 5). For every $10 \mu\text{g}\cdot\text{m}^{-3}$ increase in $PM_{2.5}$, HGB and PLT increased by 9.923% (95% CI, 8.741–11.264) and 0.068% (95% CI, 0.067–0.069), respectively.

Lag effect analysis found that with every increase in $PM_{2.5}$ exposure concentration by $10 \mu\text{g}\cdot\text{m}^{-3}$, NEUT and LMY increased by 0.006% and 0.080%, respectively, after a 1-day delay; PLT and MONO increased by 0.025% and 0.033%, respectively, with a 2-day lag; RBC and HGB increased by 8.713% and 0.012%, respectively, after a 5-days delay, while RBC increased by 7.728% with a lag of 6 days. When the lag days were 7 days, HGB and PLT increased by 0.011% and 0.016%, respectively. The correlation was not statistically significant between $PM_{2.5}$ and increased WBC concentrations with the change of lag time in the initial stage (Table 6).

Combined with Figure 7, the research showed that when the time lag accumulated for 1d (Lag0), the exposure concentration of $PM_{2.5}$ reached the maximum effect on RBC and PLT, which was statistically significant at $p < 0.05$.

Table 5. Correlation between PM_{2.5} concentration and blood routine parameters during polluted weather.

| Blood Indicators | PM _{2.5} (µg/m ³) | | |
|------------------|--|---------------------|--------------|
| | Change * (%) | 95% CI | p (sig) |
| WBC | 2.170 | 1.901~2.478 | 0.370 |
| RBC | 9.923 | 8.741~11.264 | 0.027 |
| HGB | 0.008 | 0.0078~0.0090 | 0.731 |
| PLT | 0.068 | 0.067~0.069 | 0.012 |
| NEUT | 0.006 | 0.0061~0.007 | 0.371 |
| MONO | 0.048 | 0.041~0.054 | 0.505 |
| LMY | 0.009 | 0.008~0.01 | 0.260 |

* Bold represent significant correlation between PM_{2.5} concentration and blood routine parameters at *p* < 0.05. CI, confidence interval; *p*, *p* value; WBC, white blood cell count; MONO (%), monocyte rate; RBC, red blood cell count; LMY (%), lymphatic ratio; HGB (g/L), hemoglobin; PLT, platelet count; NEUT (%), neutrophil ratio.

Table 6. Correlation between PM_{2.5} and blood routine parameters after different lag times.

| Lag Time | WBC | | RBC | | HGB | | PLT | | NEUT | | MONO | | LMY | |
|----------|----------|--------|--------------|--------|--------------|--------|--------------|--------|--------------|--------|--------------|--------|-------------|--------------|
| | <i>p</i> | RR (%) | <i>p</i> | RR (%) | <i>p</i> | RR (%) | <i>p</i> | RR (%) | <i>p</i> | RR (%) | <i>p</i> | RR (%) | <i>p</i> | RR (%) |
| Lag1 | 0.437 | 2.170 | 0.769 | 9.923 | 0.287 | 0.008 | 0.191 | 0.068 | 0.026 | 0.006 | 0.219 | 0.048 | 0.07 | 0.008 |
| Lag2 | 0.645 | 2.328 | 0.761 | 8.043 | 0.381 | 0.065 | 0.000 | 0.025 | 0.827 | 0.005 | 0.029 | 0.033 | 0.777 | 0.006 |
| Lag3 | 0.139 | 2.328 | 0.221 | 7.425 | 0.345 | 0.166 | 0.376 | 0.288 | 0.539 | 0.008 | 0.899 | 0.094 | 0.129 | 0.007 |
| Lag4 | 0.505 | 2.547 | 0.264 | 7.963 | 0.189 | 0.010 | 0.739 | 0.012 | 0.582 | 0.005 | 0.279 | 0.059 | 0.786 | 0.012 |
| Lag5 | 0.805 | 2.843 | 0.048 | 8.713 | 0.015 | 0.012 | 0.685 | 0.011 | 0.774 | 0.006 | 0.193 | 0.036 | 0.241 | 0.005 |
| Lag6 | 0.924 | 1.983 | 0.068 | 7.728 | 0.262 | 0.036 | 0.387 | 0.022 | 0.710 | 0.005 | 0.289 | 0.043 | 0.201 | 0.018 |
| Lag7 | 0.233 | 1.673 | 0.146 | 8.371 | 0.095 | 0.011 | 0.084 | 0.016 | 0.298 | 0.006 | 0.255 | 0.049 | 0.202 | 0.018 |

Bold represent significant correlation between PM_{2.5} concentration and blood routine parameters at different lag times at *p* < 0.05. *P*, level of significance; RR, relative risk; WBC, white blood cell count; MONO (%), monocyte ratio; RBC, red blood cell count; LMY (%), lymphatic ratio; HGB (g/L), hemoglobin; PLT, platelet count; NEUT (%), neutrophil ratio.




Figure 7. Effects of PM_{2.5} exposure concentration on blood routine parameters in different lag conditions. RBC, red blood cell count; HGB (g/L), hemoglobin; PLT, platelet count; NEUT (%), neutrophil ratio; MONO, monocyte ratio; LMY (%), lymphatic ratio.

When the time lag accumulates for 3d (Lag2), the exposure concentration of PM_{2.5} showed the maximum effect on MONO (statistically significant at $p < 0.05$). When the hysteresis accumulation was 6d (Lag5), the effect of PM_{2.5} exposure concentration on HGB reached the maximum, which is statistically significant at $p < 0.05$. The maximum cumulative effect of PM_{2.5} on NEUT and LMY was strongest when lagged 2d (Lag1), which was 0.006% and 0.008%, respectively. The effect value increased slightly when lagged 5 days, but the increase was no longer statistically significant at $p < 0.05$. Many earlier studies have shown that effect of PM_{2.5} on red blood cell width and RBC count is risk factors for cardiovascular disease [60], while MCHC (mean hemoglobin concentration) is a strong predictor of cardiovascular disease.

Pope III et al. [61] reported that with the increase in PM_{2.5} concentration, cardiovascular mortality rate increased by 0.84%. The another study by Pope III et al. [62] on patients in the United States with ischemic heart disease reported that with the increase of PM_{2.5} concentration, the risk of the occurrence of acute ischemic coronary artery events increased. In our study, it was found that the exposure concentration of PM_{2.5} had a significant effect on RBC and HGB under different hysteresis conditions, suggesting that PM_{2.5} may cause some damage to the cardiovascular system. Several studies have shown that an increase in PM_{2.5} concentration can increase blood indicators like NEUT and LMY [63,64]. A group study in the population of the United States found that fine particles (PM_{2.5}) can increase helper and killer T cells abruptly. The analysis results of the present study showed that the exposure concentration of PM_{2.5} would increase the neutral ratio, lymphatic ratio, and monocyte ratio under a certain lag condition, which is consistent with previous research studies [63,64].

3.3.2. Correlation Analysis of Chemical Components in Atmospheric Particles and Routine Blood Parameters

A large number of studies have shown that the contents of atmospheric particulates and their toxic chemical components are different under differentially polluted weather conditions, and their harm to animal and human health is also different [65–67]. The mass concentration of PM_{2.5} is generally high in moderate and severe pollution weather. Eight water-soluble ions and heavy metal elements have shown strong correlation with PM_{2.5} in moderate and severe pollution periods, and their concentrations fluctuate greatly with changes in PM_{2.5} concentrations. The concentrations of most water-soluble ions and heavy metal elements were higher in winter polluted weather. The total concentrations of water-soluble ions were $70.40 \mu\text{g}\cdot\text{m}^{-3}$ and $30.59 \mu\text{g}\cdot\text{m}^{-3}$ in moderate and severe pollution periods, respectively, and $30.59 \mu\text{g}\cdot\text{m}^{-3}$ in clean weather. The plasma concentrations of NO_3^- , NH_4^+ , and SO_4^{2-} were significantly increased, which indicated that NH_4^+ contributed greatly to the increase of secondary ions in winter. The total concentrations of heavy metals were $184.38 \text{ ng}\cdot\text{m}^{-3}$ and $147.22 \text{ ng}\cdot\text{m}^{-3}$ under the two weather conditions, and most of the heavy metal concentrations were increased, especially Pb and Cd concentrations. In general, the concentrations of PM_{2.5} and its chemical components in polluted weather are generally higher than those in clean weather, which is consistent with the research conclusions in international cities [68,69]. Pearson's correlation analysis results between chemical components in atmospheric particles and routine blood parameters are shown in Table 7.

The results showed that there was a significantly positive correlation between Ni and WBC, Pb and Cd and lymphoid ratio; Ca^{2+} and Mg^{2+} ions and neutrophil ratio. Furthermore, there was a significant negative correlation between As and RBC count; Ca^{2+} , Mg^{2+} , Cu, Ni and lymphoid ratio; Na^+ and Mn and hemoglobin concentration; Na^+ and platelet count; Pb and Cd and neutrophil ratio.

Table 7. Correlation between chemical constituents in fine particulate matter and blood routine parameters.

| Elements | WBC | MONO (%) | RBC | LMY (%) | HGB | PLT | NEUT (%) |
|-------------------------------|---------------|----------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Na ⁺ | −0.300 | −0.118 | −0.277 | 0.209 | −0.445 * | −0.372 * | −0.175 |
| NH ₄ ⁺ | −0.307 | 0.010 | −0.195 | 0.314 | −0.336 | −0.327 | −0.304 |
| K ⁺ | −0.323 | −0.100 | −0.154 | 0.286 | −0.345 | −0.279 | −0.281 |
| Mg ²⁺ | 0.210 | −0.341 | 0.079 | −0.484 * | −0.051 | 0.105 | 0.521 * |
| Ca ²⁺ | 0.253 | −0.236 | 0.126 | −0.585 * | 0.098 | 0.155 | 0.609 * |
| Cl [−] | −0.311 | −0.161 | −0.120 | 0.327 | −0.312 | −0.222 | −0.325 |
| NO ₃ [−] | −0.231 | 0.020 | −0.164 | 0.282 | −0.299 | −0.232 | −0.292 |
| SO ₄ ^{2−} | −0.249 | 0.064 | −0.214 | 0.130 | −0.301 | −0.327 | −0.097 |
| Cu | 0.284 | 0.151 | 0.139 | −0.409 * | 0.264 | 0.138 | 0.321 |
| Pb | −0.166 | −0.061 | −0.202 | 0.424 * | −0.352 | −0.059 | −0.440 * |
| Cr | 0.212 | 0.140 | −0.201 | −0.055 | −0.044 | −0.021 | 0.028 |
| Ni | 0.414* | 0.205 | 0.109 | −0.408 * | 0.322 | 0.085 | 0.357 |
| Cd | −0.049 | −0.048 | −0.016 | 0.428 * | −0.199 | −0.119 | −0.452 * |
| Mn | −0.080 | −0.079 | −0.312 | 0.260 | −0.388 * | −0.260 | −0.260 |
| Zn | 0.326 | 0.130 | 0.175 | −0.248 | 0.205 | 0.348 | 0.091 |
| As | 0.023 | 0.079 | −0.367 * | 0.115 | −0.360 * | −0.02 | −0.095 |

* Bold represent significant correlation at the 0.01 level (both sides). WBC, white blood cell count; MONO (%), monocyte ratio; RBC, red blood cell count; LMY (%), lymphatic ratio; HGB (g/L), hemoglobin; PLT, platelet count; NEUT (%), neutrophil ratio; Na⁺, sodium ion; NH₄⁺, ammonium ion; K⁺, potassium ion; Mg²⁺, magnesium ion; Ca²⁺, calcium ion; Cl[−], chloride ion; NO₃[−], nitrate ion; SO₄^{2−}, sulfate ion; Cu, copper; Pb, lead; Cr, chromium; Ni, nickel; Cd, cadmium; Mn, manganese; Zn, zinc; As, arsenic.

Regarding cadmium, due to its long biological half-life (approximately 20–30 years in the human body), its excretion rate in the human body is low, and it mainly exists in soft tissues (mainly liver and kidney). Therefore, cadmium metal has a variety of toxic effects, including nephrotoxicity, carcinogenicity, teratogenicity, and endocrine and reproductive toxicity. In addition, with regard to the mechanism of cadmium metal to the human body, the current evidence shows that after the human body is exposed to cadmium, it will induce genome instability through a complex multifactor mechanism. Most importantly, cadmium will interact with the DNA mismatch repair mechanism, thereby inducing apoptosis [70,71]. Previous studies have shown that increased levels of heavy metal elements Pb and Cr in particulate matter can cause decreased average daily peak expiratory flow in school-age children, and it has potential toxicity or carcinogenic risks to human health [72,73]. Small particles containing Mn, Zn, Pb, and Cr showed more significant toxic effects on human pulmonary bronchial epithelial cells [74]. The lymphatic ratio and the neutral ratio have an important indicator role in the inflammatory response [75]. The correlation analysis results of the present study revealed significant positive correlation between Pb and Cd and the lymphatic ratio and negative correlation with the neutral ratio. The changes of the lymphatic ratio and neutral ratio are more closely related to some water-soluble ions and heavy metal elements like Mg²⁺. Although the proportion of metal elements was small due to the characteristics of easy enrichment and non-degradability. Mg²⁺ can cause a stimulating effect on the skin and respiratory tract, which can cause a series of symptoms like cough and chest pain in the human body. It can be speculated, based on the results of the present study, that the heavy metal elements and water-soluble ions in atmospheric fine particles might be the main cause of changes in blood routine indicators during periods of polluted weather in the studied area.

4. Conclusions

The present study revealed the air particulate matter and its chemical components in Xinxiang City are highly polluted in winter and the overall change pattern was winter > spring > autumn > summer. The secondary air pollution in Xinxiang City is serious in winter and SO₄^{2−} NO₃[−] NH₄⁺ were the main components of water-soluble ions in PM_{2.5}. The results of the risk assessment analysis showed that the non-carcinogenic health risk of heavy metals in the fine particulate matter was acceptable to human body, while among the carcinogenic elements, the order of lifetime carcinogenic risk was arsenic(As) > chromium (Cr) > cadmium (Cd) > cobalt (Co) > nickel (Ni). This study also confirmed that some

metal elements, such as arsenic (As) and chromium (Cr), have significant negative effects on human health. The main sources of air pollution in Xinxiang City were coal combustion, vehicle exhaust emissions, and industrial emissions. Therefore, more time should be taken in the future to strengthen the control and supervision of traffic pollution sources and industrial pollution sources in and around Xinxiang City.

The research results showed that the exposure concentration of PM_{2.5} in the atmosphere would have a certain hysteresis effect on the blood routine parameters of the population, and the effect on different blood routine indexes was different under different hysteresis conditions. The present study suggested that there is a need for time to strengthen the control and supervision of traffic pollution sources and industrial pollution sources in and around the studied area. On the day of the haze, when the daily average concentration of PM_{2.5} rose by 10 µg·m⁻³, HGB and PLT increased, respectively, by 9.923% (95% CI, 8.741–11.264) and 0.068% (95% CI, 0.067–0.069). The maximum effect of exposure concentration of PM_{2.5} on MONO was reached when the lag accumulation was 3d (Lag2). When the hysteresis accumulated for 6d (Lag5), the exposure concentration of PM_{2.5} had the greatest effect on HGB. The maximum cumulative effect of PM_{2.5} on NEUT and LMY reached the strongest when the lag was 2d (Lag1). During periods of moderate to severe pollution, the concentration of water-soluble ions and heavy metal elements in PM_{2.5} increased significantly and had a significant correlation with some blood routine indicators.

The data of this study preliminarily showed that the blood routine indicators between the non-haze group and the haze group had indeed changed. The exposure concentration of PM_{2.5} had a certain lag effect on the blood routine parameters of the population, and the effect on different blood routine indexes was different under different lag conditions. This result can provide a reference for disease prevention and health management of people in haze-polluted weather in Xinxiang City. However, this study also had certain limitations. For example, the current study only included physical examination data from one hospital, and the data sampling was not comprehensive enough and was underrepresented. In addition, the air pollution exposure concentration used was regional air pollution monitoring data, which could accurately reflect the exposure of each patient. Therefore, subsequent research should focus on the correlation between the exposure concentration of each individual air pollutant and the biological indicators in order to obtain more accurate conclusions.

Author Contributions: Conceptualization, S.W. and F.P.; funding acquisition, S.W. and F.P.; investigation, S.W.; methodology, S.W. and F.P.; writing—original draft, S.W.; writing—review & editing, M.K. and T.L. All authors have read and agreed to the published version of the manuscript.

Funding: Financial support for this project was provided by the Natural Science Foundation of Henan Province, China (Project No.182300410131).

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of College of Environment, Henan Normal University, China.

Informed Consent Statement: Patient consent was waived for this study because the data was collected directly from the hospital and no direct human interaction was there.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: The authors would like to acknowledge the College of Environment, Henan Normal University, China, for providing laboratory facilities for experimental analysis.

Conflicts of Interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Li, H.; Qian, X.; Hu, W.; Wang, Y.; Gao, H. Chemical speciation and human health risk of trace metals in urban street dusts from a metropolitan city, Nanjing, SE China. *Sci. Total Environ.* **2013**, *456–457*, 212–221. [[CrossRef](#)] [[PubMed](#)]
2. Li, H.; Wang, J.; Wang, Q.; Qian, X.; Qian, Y.; Yang, M.; Li, F.; Lu, H.; Wang, C. Chemical fractionation of arsenic and heavy metals in fine particle matter and its implications for risk assessment: A case study in Nanjing, China. *Atmos. Environ.* **2015**, *103*, 339–346. [[CrossRef](#)]
3. Cao, Q.; Rui, G.; Liang, Y. Study on PM_{2.5} pollution and the mortality due to lung cancer in China based on geographic weighted regression model. *BMC Public Health* **2018**, *18*, 925. [[CrossRef](#)]
4. Duan, F.K.; He, K.B.; Ma, Y.L.; Yang, F.M.; Yu, X.C.; Cadle, S.H.; Chan, T.; Mulawa, P.A. Concentration and chemical characteristics of PM_{2.5} in Beijing, China: 2001–2002. *Sci. Total Environ.* **2006**, *355*, 264–275. [[CrossRef](#)] [[PubMed](#)]
5. Decarlo, F.P.; Ulbrich, I.M.; Crounse, J.; de Foy, B.; Dunlea, E.J.; Aiken, A.C.; Knapp, D.; Weinheimer, A.J.; Campos, T.; Wennberg, P.O. Investigation of the sources and processing of organic aerosol over the central Mexican plateau from aircraft measurements during MILAGRO. *Atmos. Chem. Phys.* **2010**, *10*, 5257–5280. [[CrossRef](#)]
6. Atkinson, R.W.; Mills, I.C.; Walton, H.A.; Anderson, H.R. Fine particle components and health—A systematic review and meta-analysis of epidemiological time series studies of daily mortality and hospital admissions. *J. Exp. Sci. Environ. Epidemiol.* **2015**, *25*, 208–214. [[CrossRef](#)] [[PubMed](#)]
7. Zhang, H.H.; Li, Z.; Liu, Y.; Xinag, P.; Cui, X.Y.; Ye, H.; Hu, B.L.; Lou, L.P. Physical and chemical characteristics of PM_{2.5} and its toxicity to human bronchial cells BEAS-2B in the winter and summer. *J. Zhejiang Univ. Sci. B* **2018**, *19*, 317–326. [[CrossRef](#)]
8. Wang, K.; Wang, W.; Li, L.; Li, J.; Wei, L.; Chi, W.; Hong, L.; Zhao, Q.; Jiang, J. Seasonal concentration distribution of PM_{1.0} and PM_{2.5} and a risk assessment of bound trace metals in Harbin, China: Effect of the species distribution of heavy metals and heat supply. *Sci. Rep.* **2020**, *10*, 8160. [[CrossRef](#)]
9. Xing, Y.F.; Xu, Y.H.; Shi, M.H.; Lian, Y.X. The impact of PM_{2.5} on the human respiratory system. *J. Thorac. Dis.* **2016**, *8*, E69–E74. [[PubMed](#)]
10. Feng, J.; Yu, H.; Su, X.; Liu, S.; Li, Y.; Pan, Y.; Sun, J.-H. Chemical composition and source apportionment of PM_{2.5} during Chinese spring festival at Xinxiang, a heavily polluted city in north China: Fireworks and health risks. *Atmos. Res.* **2016**, *182*, 176–188. [[CrossRef](#)]
11. D’Amato, G.; Baena-Cagnani, C.E.; Cecchi, L.; Annesi-Maesano, I.; Nunes, C.; Ansotegui, I.; D’Amato, M.; Liccardi, G.; Sofia, M.; Canonica, W.G. Climate change, air pollution and extreme events leading to increasing prevalence of allergic respiratory diseases. *Multidiscip. Respir. Med.* **2013**, *8*, 12. [[CrossRef](#)]
12. Naddafi, K.; Hassanvand, M.S.; Yunesian, M.; Momeniha, F.; Nabizadeh, R.; Faridi, S.; Gholampour, A. Health impact assessment of air pollution in megacity of Tehran, Iran. *Iran. J. Environ. Health Sci. Eng.* **2012**, *9*, 28. [[CrossRef](#)] [[PubMed](#)]
13. Xie, Y.B.; Chen, J.; Li, W. Health risks and damage value assessment of Beijing residents’ continuous exposure to high concentrations of PM_{2.5} during heavy haze pollution. *Environ. Sci.* **2014**, *35*, 1–8. (In Chinese)
14. Zhu, Y.D.; Wei, J.R.; Huang, L.; Wang, S.H.; Tian, H.M.; Guo, X.B. Comparison of the occurrence of respiratory diseases and symptoms among school-age children in areas with different air pollution levels. *J. Peking Univ. Med. Ed.* **2015**, *47*, 395–399.
15. Rich, D.Q.; Kipen, H.M.; Huang, W.; Wang, G.; Wang, Y.; Zhu, P.; Ohman-Strickland, P.; Hu, M.; Philipp, C.; Diehl, S.R.; et al. Association between changes in air pollution levels during the Beijing Olympics and biomarkers of inflammation and thrombosis in healthy young adults. *JAMA* **2012**, *307*, 2068–2078. [[CrossRef](#)] [[PubMed](#)]
16. Fajersztajn, L.; Veras, M.; Barrozo, L.V.; Saldiva, P. Air pollution: A potentially modifiable risk factor for lung cancer. *Nat. Rev. Cancer* **2013**, *13*, 674–678. [[CrossRef](#)] [[PubMed](#)]
17. Hu, J.R.; Xu, H.J.; Li, Q.Y.; Li, C.L.; Sun, X.W.; Zhang, X.J. Effects of PM_{2.5} on chronic airway inflammation and oxidative stress in passive smoking rats. *J. Shanghai Jiaotong Univ. Med. Ed.* **2014**, *34*, 619–625. (In Chinese)
18. Wang, Z.; Calderón, L.; Patton, A.P.; Allacci, M.S.; Senick, J.; Wener, R.; Andrews, C.J.; Mainelis, G. Comparison of real-time instruments and gravimetric method when measuring particulate matter in a residential building. *J. Air Waste Manag. Assoc.* **2016**, *66*, 1109–1120. [[CrossRef](#)] [[PubMed](#)]
19. Thurston, G.D.; Spengler, J.D. A quantitative assessment of source contributions to inhalable particulate matter pollution in metropolitan Boston. *Atmos. Environ.* **1985**, *19*, 9–25. [[CrossRef](#)]
20. Miller, S.L.; Anderson, M.J.; Daly, E.P.; Milford, J.B. Source apportionment of exposures to volatile organic compounds. I. Evaluation of receptor models using simulated exposure data. *Atmos. Environ.* **2002**, *36*, 3629–3641. [[CrossRef](#)]
21. Ministry of Environmental Protection of China. *Exposure Factors Handbook of Chinese Population*; China Environmental Science Press: Beijing, China, 2013.
22. Zhang, H.; Wang, Y.; Hu, J.; Ying, Q.; Hu, X.M. Relationships between meteorological parameters and criteria air pollutants in three megacities in China. *Environ. Res.* **2015**, *140*, 242–254. [[CrossRef](#)] [[PubMed](#)]
23. Peng, R.D.; Dominici, F.; Louis, T.A. Model choice in time series studies of air pollution and mortality. *J. R. Stat. Soc. A Stat. Soc.* **2006**, *169*, 179–203. [[CrossRef](#)]
24. Dai, Q.; Liu, B.; Bi, X.; Wu, J.; Liang, D.; Zhang, Y.; Feng, Y.; Hopke, P.K. Dispersion Normalized PMF Provides Insights into the Significant Changes in Source Contributions to PM_{2.5} after the COVID-19 Outbreak. *Environ. Sci. Technol.* **2020**, *54*, 9917–9927. [[CrossRef](#)] [[PubMed](#)]

25. Liu, H.; Tian, H.; Zhang, K.; Liu, S.; Cheng, K.; Yin, S.; Liu, Y.; Liu, X.; Wu, Y.; Liu, W.; et al. Seasonal variation, formation mechanisms and potential sources of PM_{2.5} in two typical cities in the Central Plains Urban Agglomeration, China. *Sci. Total Environ.* **2019**, *657*, 657–670. [CrossRef] [PubMed]
26. Liu, T.; Gong, S.; He, J.; Yu, M.; Wang, Q.; Li, H.; Liu, W.; Zhang, J.; Li, L.; Wang, X.; et al. Attributions of meteorological and emission factors to the 2015 winter severe haze pollution episodes in china's Jing-Jin-Ji area. *Atmos. Chem. Phys.* **2017**, *17*, 2971–2980. [CrossRef]
27. An, Z.; Huang, R.J.; Zhang, R.; Tie, X.; Li, G.; Cao, J.; Zhou, W.; Shi, Z.; Han, Y.; Gu, Z.; et al. Severe haze in northern China: A synergy of anthropogenic emissions and atmospheric processes. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 8657–8666. [CrossRef]
28. Huang, R.J.; Zhang, Y.; Bozzetti, C.; Ho, K.F.; Cao, J.J.; Han, Y.; Daellenbach, K.R.; Slowik, J.G.; Platt, S.M.; Canonaco, F.; et al. High secondary aerosol contribution to particulate pollution during haze events in China. *Nature* **2014**, *514*, 218–222. [CrossRef]
29. Cohen, A.J.; Brauer, M.; Burnett, R.; Anderson, H.R.; Frostad, J.; Estep, K.; Balakrishnan, K.; Brunekreef, B.; Dandona, L.; Dandona, R.; et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: An analysis of data from the Global Burden of Diseases Study 2015. *Lancet* **2017**, *389*, 1907–1918. [CrossRef]
30. HEI (Health Effects Institute). *State of Global Air 2019*; Special Report; Health Effects Institute: Boston, MA, USA, 2019.
31. Lee, J.S.H.; Jaafar, Z.; Tan, A.K.J.; Carrasco, L.R.; Ewing, J.J.; Bickford, D.P. Toward clearer skies: Challenges in regulating transboundary haze in southeast asia. *Environ. Sci. Policy* **2016**, *55*, 87–95. [CrossRef]
32. Tan, S.H.; Roth, M.; Velasco, E. Particle exposure and inhaled dose during commuting in Singapore. *Atmos. Environ.* **2017**, *170*, 245–258. [CrossRef]
33. Zhang, Y.L.; Cao, F. Fine particulate matter (PM_{2.5}) in China at a city level. *Sci. Rep.* **2015**, *5*, 14884. [CrossRef]
34. Leng, C.; Duan, J.; Xu, C.; Zhang, H.; Wang, Y.; Wang, Y. Insights into a historic severe haze event in Shanghai: Synoptic situation, boundary layer and pollutants. *Atmos. Chem. Phys.* **2016**, *16*, 9221–9234. [CrossRef]
35. Yu, L.D.; Wang, G.F.; Zhang, R.J.; Zhang, L.M.; Yu, S.; Wu, B.B.; Li, X.F.; An, K.; Chu, J.H. Characterization and Source Apportionment of PM_{2.5} in an Urban Environment in Beijing. *Aerosol Air Qual. Res.* **2013**, *13*, 574–583. [CrossRef]
36. Tao, J.; Zhang, L.; Cao, J.; Zhong, L.; Chen, D.; Yang, Y.; Chen, D.; Chen, L.; Zhang, Z.; Wu, Y. Source apportionment of PM_{2.5} at urban and suburban areas of the Pearl River Delta region, south China—With emphasis on ship emissions. *Sci. Total Environ.* **2017**, *574*, 1559–1570. [CrossRef]
37. Liu, Z.X.; Wang, G.Z.; Guo, R.; Huang, K.; Zhang, Y. Chemical characteristics and source apportionment of atmospheric dustfall in Pingdingshan City, Henan Province. *Environ. Chem.* **2011**, *30*, 825–831.
38. Qiao, B.W.; Liu, Z.Y.; Hu, B.; Liu, J.Y.; Feng, N.N.; Wu, F.K.; Xu, Z.J.; Wang, Y.S. Analysis of the concentration characteristics and sources of metal elements in PM_{2.5} in winter in Beijing. *Environ. Sci.* **2017**, *38*, 876–883. (In Chinese)
39. Yao, Q.; Han, S.Q.; Cai, Z.Y. Pollution and ecological risk assessment of heavy metals in PM_{2.5} during heating period in Tianjin. *China Environ. Sci.* **2013**, *9*, 1596–1600.
40. Liu, F.L.; Lu, X.; Wu, M.L.; Liu, J.; Ren, Y.R.; Guo, Z.B. Pollution characteristics and source apportionment of heavy metals in atmospheric fine particles in Nanjing. *J. Environ. Eng.* **2014**, *8*, 652–658.
41. Qi, M.Y.; Lu, X.H.; Zhang, C.Y.; Zhao, Y.; Ji, S.P.; Wang, Y.; Wang, L.T.; Ma, X. Study on PM_{2.5} Concentration Variation Characteristics and Pollution Source Areas in Handan City from 2013 to 2017. Available online: kns.cnki.net/kcms/detail/detail.aspx?DBCode=CJFD&DBName=CJFDLAST2019&fileName=HJXU201804018 (accessed on 6 February 2021).
42. Hu, G.; Zhang, Y.; Sun, J.; Zhang, L.; Shen, X.; Lin, W.; Yang, Y. Variability, formation and acidity of water-soluble ions in PM_{2.5} in Beijing based on the semi-continuous observations. *Atmos. Res.* **2014**, *145–146*, 1–11. [CrossRef]
43. Zheng, S.; Pozzer, A.; Cao, C.X.; Lelieveld, J. Long-term (2001–2012) fine particulate matter (PM_{2.5}) and the impact on human health in Beijing, China. *Atmos. Chem. Phys.* **2015**, *15*, 5715–5725. [CrossRef]
44. Gao, J.; Tian, H.; Cheng, K.; Lu, L.; Zheng, M.; Wang, S.; Hao, J.; Wang, K.; Hua, S.; Zhu, S.; et al. The variation of chemical characteristics of PM_{2.5} and PM₁₀ and formation causes during two haze pollution events in urban Beijing, China. *Atmos. Environ.* **2015**, *107*, 1–8. [CrossRef]
45. Zhou, X.; Cao, Z.; Ma, Y.; Wang, L.; Wu, R.; Wang, W. Concentrations, correlations and chemical species of PM_{2.5}/PM₁₀ based on published data in China: Potential implications for the revised particulate standard. *Chemosphere* **2016**, *144*, 518–526. [CrossRef] [PubMed]
46. Rogula-Kozłowska, W.; Klejnowski, K.; Rogula-Kopiec, P. Spatial and seasonal variability of the mass concentration and chemical composition of PM_{2.5} in Poland. *Air Qual. Atmos. Health* **2014**, *7*, 41–58. [CrossRef]
47. Sun, Y.; Jiang, Q.; Wang, Z.; Fu, P.; Li, J.; Yang, T.; Yin, Y. Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013. *J. Geophys. Res. Atmos.* **2014**, *119*, 4380–4398. [CrossRef]
48. Lv, Z.; Zhu, G.R.; Xiao, H.W.; Luo, L.; Zhang, Z.Y.; Zheng, N.J. Composition characteristics and source apportionment of PM_{2.5} low molecular weight organic acids in autumn in Shijiazhuang. *Environ. Pollut. Prev.* **2018**, *40*, 14–18. (In Chinese)
49. Zhang, Y.Y. Online Monitoring of Water Soluble Ions in PM_{2.5} in Northern Suburb of Nanjing. M.Sc. Thesis, Nanjing University of Information Science and Technology, Nanjing, China, 2018. (In Chinese).
50. Park, S.S.; Kim, Y.J. PM_{2.5} particles and size-segregated ionic species measured during fall season in three urban sites in Korea. *Atmos. Environ.* **2004**, *38*, 1459–1471. [CrossRef]
51. Wang, X.Q.; Zhou, Y.; Cheng, S.Y.; Wang, G. Characterization and regional transmission impact of water-soluble ions in PM_{2.5} during winter in typical cities. *China Environ. Sci.* **2016**, *36*, 2289–2296.

52. Pulles, T.; van der Gon, H.D.; Appelman, W.; Verheul, M. Emission factors for heavy metals from diesel and petrol used in European vehicles. *Atmos. Environ.* **2012**, *61*, 641–651. [[CrossRef](#)]
53. Yatkin, S.; Bayram, A. Elemental composition and sources of particulate matter in the ambient air of a metropolitan city. *Atmos. Res.* **2007**, *85*, 126–139. [[CrossRef](#)]
54. Morawska, L.; Zhang, J. Combustion sources of particles. 1. Health relevance and source signatures. *Chemosphere* **2002**, *49*, 1045–1058. [[CrossRef](#)]
55. Mazzei, F.; D'Alessandro, A.; Lucarelli, F.; Nava, S.; Prati, P.; Valli, G.; Vecchi, R. Characterization of particulate matter sources in an urban environment. *Sci. Total Environ.* **2008**, *401*, 81–89. [[CrossRef](#)] [[PubMed](#)]
56. Callén, M.S.; de la Cruz, M.T.; López, J.M.; Navarro, M.V.; Mastral AM, J.C. Comparison of receptor models for source apportionment of the PM₁₀ in Zaragoza (Spain). *Chemosphere* **2009**, *76*, 1120–1129. [[CrossRef](#)]
57. EPA (Environmental Protection Agency). *Soil Screening Guidance: User's Guide*; EPA/540/R-96/018; Office of Emergency and Remedial Response: Washington, DC, USA, 1996.
58. Fa-Yun, L.; Cheng, H.; Ying, Z.; Man-li, S.; Xiao-nan, Y. Environmental impact and health risk assessment of heavy metals in street dust in Shenyang, Liaoning province. *Liaoning Meteorol.* **2010**, *6*, 59–64.
59. Zheng, N.; Liu, J.; Wang, Q.; Liang, Z. Heavy metals exposure of children from stairway and sidewalk dust in the smelting district, Northeast of China. *Atmos. Environ.* **2010**, *44*, 3239–3245. [[CrossRef](#)]
60. Slezakova, K.; Morais, S.; Pereira, M.D. Trace metals in size-fractionated particulate matter in a Portuguese hospital: Exposure risks assessment and comparisons with other countries. *Environ. Sci. Pollut. Res. Int.* **2014**, *21*, 3604–3620. [[CrossRef](#)]
61. Gray, D.L.; Wallace, L.A.; Brinkman, M.C.; Buehler, S.S.; La Londe, C. Respiratory and cardiovascular effects of metals in ambient particulate matter: A critical review. *Rev. Environ. Contam. Toxicol.* **2015**, *234*, 135–203. [[PubMed](#)]
62. Pope, C.A., III; Bhatnagar, A.; McCracken, J.P.; Abplanalp, W.; Conklin, D.J.; O'Toole, T. Exposure to fine particulate air pollution is associated with endothelial injury and systemic inflammation. *Circ. Res.* **2016**, *119*, 1204–1214. [[CrossRef](#)] [[PubMed](#)]
63. Pope, C.A., III; Muhlestein, J.B.; May, H.T.; Renlund, D.G.; Anderson, J.L.; Horne, B.D. Ischemic heart disease events triggered by short-term exposure to fine particulate air pollution. *Circ. Res.* **2006**, *114*, 2443–2448. [[CrossRef](#)] [[PubMed](#)]
64. Hoffmann, B.; Weinmayr, G.; Hennig, F.; Fuks, K.; Moebus, S.; Weimar, C.; Dragano, N.; Hermann, D.M.; Kälsch, H.; Mahabadi, A.A.; et al. Air quality, stroke, and coronary events: Results of the Heinz Nixdorf Recall study from the Ruhr region. *Dtsch. Arztebl. Int.* **2015**, *112*, 195–201. [[PubMed](#)]
65. Folino, F.; Buja, G.; Zanutto, G.; Marras, E.; Allocca, G.; Vaccari, D.; Gasparini, G.; Bertaglia, E.; Zoppo, F.; Calzolari, V.; et al. Association between air pollution and ventricular arrhythmias in high-risk patients (ARIA study): A multi-centre longitudinal study. *Lancet Planet. Health* **2017**, *1*, e58–e64. [[CrossRef](#)]
66. Brew, K.; Dinakarandian, D.; Nagase, H. Tissue inhibitors of metalloproteinases: Evolution, structure and function. *Biochim. Biophys. Acta* **2000**, *1477*, 267–283. [[CrossRef](#)]
67. Choussat, R.; Montalescot, G.; Collet, J.P.; Jardel, C.; Ankri, A.; Fillet, A.M.; Thomas, D.; Raymond, J.; Bastard, J.P.; Drobinski, G.; et al. Effect of prior exposure to *Chlamydia pneumoniae*, *Helicobacter pylori*, or cytomegalovirus on the degree of inflammation and one-year prognosis of patients with unstable angina pectoris or non-Q-wave acute myocardial infarction. *Am. J. Cardiol.* **2000**, *86*, 379–384. [[CrossRef](#)]
68. Driscoll, K.E.; Carter, J.M.; Hassenbein, D.G.; Howard, B. Cytokines and particle-induced inflammatory cell recruitment. *Environ. Health Perspect.* **1997**, *105* (Suppl. S5), 1159–1164. [[PubMed](#)]
69. Omar, N.Y.; Mon, T.C.; Rahman, N.A.; Abas, M.R. Distributions and health risks of polycyclic aromatic hydrocarbons (PAHs) in atmospheric aerosols of Kuala Lumpur, Malaysia. *Sci. Total Environ.* **2006**, *369*, 76–81. [[CrossRef](#)] [[PubMed](#)]
70. See, S.W.; Balasubramanian, R.; Wang, W. A study of the physical, chemical, and optical properties of ambient aerosol particles in southeast Asia during hazy and non-hazy days. *J. Geophys. Res. Atmos.* **2006**, *111*, D10. [[CrossRef](#)]
71. Giaginis, C.; Gatzidou, E.; Theocharis, S. DNA repair systems as targets of cadmium toxicity. *Toxicol. Appl. Pharmacol.* **2006**, *213*, 282–290. [[CrossRef](#)]
72. Zarros, A. Cadmium (Cd) as a carcinogenetic factor and its participation in the induction of lung cancer. *Pneumon* **2008**, *21*, 172–177.
73. Bandowe, B.A.; Meusel, H.; Huang, R.J.; Ho, K.; Cao, J.; Hoffmann, T.; Wilcke, W. PM_{2.5}-bound oxygenated PAHs, nitro-PAHs and parent-PAHs from the atmosphere of a Chinese megacity: Seasonal variation, sources and cancer risk assessment. *Sci. Total Environ.* **2013**, *473–474*, 77–87. [[CrossRef](#)]
74. Hong, Y.C.; Hwang, S.S.; Kim, J.H.; Lee, K.H.; Lee, H.J.; Lee, K.H.; Yu, S.D.; Kim, D.S. Metals in particulate pollutants affect peak expiratory flow of schoolchildren. *Environ. Health Perspect.* **2007**, *115*, 430–434. [[CrossRef](#)]
75. Mirowsky, J.; Hickey, C.; Horton, L.; Blaustein, M.; Galdanes, K.; Peltier, R.E.; Chillrud, S.; Chen, L.C.; Ross, J.; Nadas, A.; et al. The effect of particle size, location and season on the toxicity of urban and rural particulate matter. *Inh. Toxicol.* **2013**, *25*, 747–757. [[CrossRef](#)] [[PubMed](#)]