



Article

# Understanding How to Improve the Use of Clinical Coordination Mechanisms between Primary and Secondary Care Doctors: Clues from Catalonia

Laura Esteve-Matalí <sup>1,2</sup>, Ingrid Vargas <sup>1,\*</sup>, Franco Amigo <sup>1,3</sup>, Pere Plaja <sup>4</sup>, Francesc Cots <sup>5</sup>, Erick F. Mayer <sup>6</sup>, Joan-Manuel Pérez-Castejón <sup>7</sup> and María-Luisa Vázquez <sup>1</sup>

<sup>1</sup> Health Policy and Health Services Research Group, Health Policy Research Unit, Consortium for Health Care and Social Services of Catalonia, 08022 Barcelona, Spain; lesteve@consorci.org (L.E.-M.); famigo@imim.es (F.A.); mlvazquez@consorci.org (M.-L.V.)

<sup>2</sup> Department for Paediatrics, Obstetrics and Gynaecology, Preventive Medicine, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

<sup>3</sup> Health Services Research Unit, IMIM-Institut Hospital del Mar d'Investigacions Mèdiques, 08003 Barcelona, Spain

<sup>4</sup> Fundació Salut Empordà, 17600 Figueres, Spain; pplaja@salutemporda.cat

<sup>5</sup> Parc de Salut Mar, 08019 Barcelona, Spain; FCots@parcdesalutmar.cat

<sup>6</sup> Serveis de Salut Integrats Baix Empordà, 17230 Palamós, Spain; emayer@ssibe.cat

<sup>7</sup> Badalona Serveis Assistencials, 08911 Badalona, Spain; jpcastejon@bsa.cat

\* Correspondence: ivargas@consorci.org

**Table S1.** Distribution of some characteristics of the doctors of the Catalan national health system (NHS).

Doctors of the Catalan NHS	
<b>Sex</b>	
Male	55.1%
Female	44.9%
<b>Age (years old)</b>	
Mean	44.6
<b>Level of care</b>	
PC	27.4%
SC	72.6%

Source: Solsona et al. [54].