

**Supplementary File S1.** Women excluded vs women included in the final analysis.

Variables	Women Included n= 116	Women excluded n=30	p
<b>Age <math>\bar{x}</math> (SD)</b>	32.08 (4.68)	32.09 (6.23)	0.993 <sup>1</sup>
<b>Country of origin n (%)</b>			
Spain	109 (94.0)	26 (86.7)	0.369 <sup>2</sup>
Other	7 (6.0)	4 (13.3)	
<b>Education n (%)</b>			
No studies/primary	40 (34.5)	10 (33.3)	0.234 <sup>3</sup>
Secondary	53 (45.7)	7 (23.3)	
University/postgraduate	23 (19.8)	13 (43.3)	
<b>Employment situation n (%)</b>			
Student	2 (1.7)	0 (0)	0.573 <sup>2</sup>
Unemployed	41 (35.3)	14 (46.7)	
Work for others	65 (56.0)	16 (53.3)	
Own-account work	8 (6.9)	0 (0)	
<b>Stable partner n (%)</b>	116 (100.0)	30 (100)	1.000 <sup>3</sup>
<b>Maternal Education n (%)</b>			
Yes	46 (39.7)	12 (40.0)	0.963 <sup>3</sup>
No	70 (60.3)	18 (60.0)	
<b>Number of professionals who provided care, n (%)</b>			
1 - 2	37 (31.9)	13 (43.3)	0.329 <sup>2</sup>
3 - 5	67 (57.8)	13 (43.3)	
> 5	26 (10.4)	4 (13.3)	
<b>Number of professionals who performed vaginal examination, n (%)</b>			
0 - 2	63 (54.3)	16 (53.3)	0.852 <sup>2</sup>
3 - 5	52 (44.8)	14 (46.7)	
> 5	1 (0.9)	0 (0)	
<b>Labor Companion, n (%)</b>			
Partner	103 (88.8)	23 (76.7)	0.092 <sup>2</sup>
Other relative	4 (3.4)	0 (0)	
Nobody	9 (7.8)	7 (23.3)	
<b>Puerperium Companion, n (%)</b>			
Partner	111 (95.7)	27 (90.0)	0.327 <sup>2</sup>
Other relative	2 (1.7)	0 (0)	
Nobody	3 (2.6)	3 (10.0)	
<b>The companion was chosen prior to childbirth, n (%)</b>	104 (89.7)	30 (100)	0.460 <sup>3</sup>
<b>Planned epidural analgesia before childbirth, n (%)</b>	85 (73.3)	24 (80.0)	0.618 <sup>3</sup>
<b>Planned mode of new-born feeding (prior to birth)</b>			
Exclusive breastfeeding	91 (78.4)	23 (76.7)	0.653 <sup>2</sup>
Partial breastfeeding	11 (9.5)	2 (6.7)	
Formula feeding	14 (12.1)	5 (16.6)	

<sup>1</sup> t-student test; <sup>2</sup> Fisher test; <sup>3</sup> Chi square test