

Table S1. Tobacco and alcohol use characteristics during the periods of study

	Period of Study		
	Before COVID-19	Lockdown	New Normal
	N(%)	N(%)	N(%)
Tobacco			
<i>Non-smoker</i>			
Total	6780 (72.6)	457 (74.0)	420 (73.8)
Men	1871 (74.1)	102 (72.9)	107 (74.3)
Women	4909 (72.1)	355 (74.3)	313 (73.7)
<i>Ex-smoker</i>			
Total	682 (7.3)	63 (10.2)	49 (8.6)
Men	178 (7.1)	11 (7.9)	18 (12.5)
Women	504 (7.4)	52 (10.9)	31 (7.3)
<i>Current smoker</i>			
Total	1873 (20.1)	98 (15.9)	100 (17.6)
Men	477 (18.9)	27 (19.3)	19 (13.2)
Women	1396 (20.5)	71 (14.9)	81 (19.1)
Alcohol			
<i>Past-30 days</i>			
Total	7334 (78.6)	394 (63.8)	442 (77.7)
Men	2040 (80.8)	89 (63.6)	106 (73.6)
Women	5294 (77.8)	305 (63.8)	336 (79.1)

Figure S1: Percentage of university students starting cannabis use by age and period of study.

