

Table S1. Questionnaire.

Knowledge of COVID-19 infection prevention and control guidelines before, during, and after performing dental procedures among dental students, interns, and dentists in Saudi Arabia.	
Demographic and general characteristics	Gender
	<input type="radio"/> MALE <input type="radio"/> FEMALE
	Age
	<input type="radio"/> 20-29 YRS <input type="radio"/> 30 - 39 YRS <input type="radio"/> 40 -49 YRS <input type="radio"/> 50 -59 YRS <input type="radio"/> ABOVE 60 YRS
	Professional experience
	<input type="radio"/> Students in dental clinics <input type="radio"/> Intern <input type="radio"/> 0-9 yrs <input type="radio"/> 10-19 yrs <input type="radio"/> 20 yrs and above
	Work/Study sector (Check all that apply)
	<input type="radio"/> Academic institute dental clinics <input type="radio"/> Private dental clinics <input type="radio"/> government hospital dental clinics <input type="radio"/> military hospital dental clinics <input type="radio"/> Non practising dentists <input type="radio"/> others
	Region of study/practice in Saudi Arabia
	<input type="radio"/> Riyadh region <input type="radio"/> Makkah region <input type="radio"/> Madinah region <input type="radio"/> Qassim region <input type="radio"/> Eastern region <input type="radio"/> Asir region <input type="radio"/> Tabuk region <input type="radio"/> Hail region <input type="radio"/> The northern border region <input type="radio"/> Jazan region <input type="radio"/> Najran region <input type="radio"/> Al Baha region <input type="radio"/> Al jouf region
	Have you read any guidelines for providing dental services during COVID-19 Pandemic (MOH, WHO, CDC, ADA, NHS, etc)?
	<input type="radio"/> YES <input type="radio"/> NO
	Have you attended any COVID-19 infection prevention workshops?
	<input type="radio"/> YES <input type="radio"/> NO
	The following questions are relevant to COVID-19 infection prevention "BEFORE" performing dental procedures
<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree	
2. Patients with reversible pulpitis and dentine hypersensitivity should be recommended analgesics if needed, avoid stimuli (cold, hot and acidic drinks or food), apply desensitizing toothpaste regularly to the sensitive area with a finger, and advise the patient to call back if symptoms get worse.	
<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree	
3. Prevent crowding in appointment settings by booking appointments	

	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
4. Any dental procedures should be delayed in patients with a history of COVID-19 for at least a month	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
5. High-risk patients like diabetic and immunocompromised patients should be treated early in a dental office opening.	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
6. Telephonic triage / Tele dentistry should not be considered an alternative to in-office care.	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
7. Patients with fracture / loose tooth fragments or broken restorations should be referred to the designated urgent dental clinics during the COVID-19 pandemic.	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
8. The temperature of staff and patients should be monitored daily	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
9. Ask dental health care personnel to stay home if they are sick	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
10. Patients with fever should be referred to a specific medical centre's treating COVID 19	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
11. Accompanying individuals with patients should be allowed in the clinics.	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
12. Hand disinfection with 60 - 75% alcohol should be offered upon entrance to the dental office.	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree ○ Neutral ○ Agree ○ Strongly Agree
13. Emergency dental care can be provided if a patient's temperature is less than 100.4-degrees Fahrenheit and does not have symptoms consistent with COVID-19.	<ul style="list-style-type: none"> ○ Strongly Disagree ○ Disagree

	<input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
14. The waiting area should be large with adequate ventilation.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
15. The 2-meter separation between patients is mandatory in waiting rooms and reception areas.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
16. Remove magazines, toys, and other objects which cannot be easily disinfected	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
17. Posters in the dental office for instructing patients on standard recommendations for respiratory hygiene/cough etiquette and social distancing should be posted in appropriate places.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
18. It is not required by everyone entering the dental office to use facemasks or cloth face coverings.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
19. Dental procedures require professionals to use Personal protective equipment (surgical caps, gloves, N-95 mask, FFP2 mask, goggles, gowns, and face shields).	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
20. It is not required to cover all touchable surfaces with disposable protections.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
21. Patients should not be treated in rooms with negative pressure relative to the surrounding area.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
22. In case hands are visibly soiled, water and soap should be used at least 20 seconds before using an Alcohol-based hand rub.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
23. Preprocedural mouth rinse like 1.5% hydrogen peroxide or 0.2% povidone should not be used before starting any dental procedure in the patient.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral

The following questions are relevant to COVID-19 infection prevention "DURING" the dental procedures

	<input type="radio"/> Agree <input type="radio"/> Strongly Agree
24. Avoid the use of topical spray anaesthesia to prevent gag reflex	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
25. Use of rubber dam and N-95 masks are mandatory for aerosol-generating dental procedures	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
26. High-volume saliva ejectors can increase aerosol or spatter while performing dental procedures.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
27. Panoramic radiographs or cone-beam computed tomographs should not be used intraoral radiography.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
28. Four-handed dentistry should not be practised for aerosol-generating procedures.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
29. Use of 3-in-1 syringes, air-water syringes, and ultrasonic instruments are allowed for all aerosol-generating dental procedures	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
30. Adopt the Atraumatic Restorative Technique and Chemo mechanical caries removal procedure wherever possible	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
31. To reduce the clinical time, preferences should be given to bulk-fill composite resin restorations as it permits increments up to 4mm in thickness.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
32. Treatment should be completed in multiple visits wherever possible.	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree
33. Environmental cleaning and disinfection procedures should be followed after completion of treatment	<input type="radio"/> Strongly Disagree <input type="radio"/> Disagree <input type="radio"/> Neutral <input type="radio"/> Agree <input type="radio"/> Strongly Agree

The following questions are relevant to COVID-19 infection prevention "AFTER" performing dental procedures

34. Clean and disinfect reusable PPE

- Strongly Disagree
 - Disagree
 - Neutral
 - Agree
 - Strongly Agree
-

35. Manage laundry and medical waste following routine procedures

- Strongly Disagree
 - Disagree
 - Neutral
 - Agree
 - Strongly Agree
-