

**Title:** The geriatric G8 score is associated with survival outcomes in older patients with advanced prostate cancer in the ADHERE prospective study of the Meet-URO network.

**Supplementary Table S1.** Patient characteristics in the whole ADHERE study cohort and by ARPI


Characteristic	Abiraterone	Enzalutamide	All patients	P Value
	Cohort ( <i>n</i> = 86)	Cohort ( <i>n</i> = 148)	( <i>n</i> = 234)	
	Value	Value	Value	
	No. (%) [IQR, 95% <i>CI</i> ]	No. (%) [IQR, 95% <i>CI</i> ]	No. (%) [IQR, 95% <i>CI</i> ]	
Age <sup>a</sup> , median, years	78 [73-82]	77.5 [73-82]	78 [73-82]	0.50
Gleason score, median	8 [7-9]	8 [7-9]	8 [7-9]	0.194
Surgery at diagnosis	38 (44)	50 (34)	88 (38)	0.149
Time to CR, mo, median	39.3 [20.5-63.4]	30.1 [11.8-57.1]	31.4 [14.7-58.6]	0.072
Sites of metastases <sup>a</sup>				
Lymph nodes (only)	20 (23)	29 (20)	49 (21)	0.480
Bone (non-visceral)	56 (65)	107 (72)	163 (70)	
Visceral	10 (12)	12 (8)	22 (9)	
PSA at Tx start, median	11.8 [3.1-29.9]	14.3 [5.4-35.9]	14.0 [4.5-33.9]	0.325
Setting of therapy				
Pre-chemotherapy	61 (71)	101 (68)	162 (69)	0.222
Post-chemotherapy	17 (20)	40 (27)	57 (24)	
Post-Abi/Enza	8 (9)	7 (5)	15 (6)	
Steroid use <sup>b</sup> (yes)	86 (100)	14 (9)	100 (43)	<b>&lt;0.001</b>
Charlson score, median	10 [8-11]	10 [10-12]	10 [9-11]	<b>0.028</b>
Geriatric G8, median	14 [12-16]	13 [12-15]	14 [12-15]	0.081
IADL, median	6 [5-8]	6 [5-8]	6 [5-8]	0.639

Concomitant therapies, no.	3 [1-5]	3 [1-5]	3 [1-5]	0.671
Caregiver (yes)	71 (83)	119 (80)	190 (81)	0.816

<sup>a</sup>At the time of initiation of treatment.

<sup>b</sup>During the whole treatment.

Abi, abiraterone; ARPI, androgen receptor pathway inhibitors; CI, confidence intervals; CR, castration resistance; Δ, difference between the PSA basal value and the best response value; Enza, enzalutamide; mo., months; G, grade; IQR, interquartile range; Tx, treatment; yr, year.



**Supplementary Figure S1.** Histogram of distribution of patients based on Geriatric G8 score