

Supplementals

Triggers and Coping Strategies for Fear of Cancer Recurrence in Cancer Survivors: A Qualitative Study

Xu Zhang ¹, Di Sun ², Zhiwen Wang ^{1,*} and Nan Qin ³.

Table S1. Detailed information for each participant.

Number	Gender	Age	Employment Status	Monthly Income	Cancer Treatment	Cancer Type	Time Since First Diagnosis	FCRI-SF Score
P1	Male	67	Retirement	>5000	Chemotherapy	Lung	3	5
P2	Female	62	Retirement	<2000	Chemotherapy+ Surgery	Gynecology	6	19
P3	Female	72	Retirement	2000~5000	Chemotherapy+ Surgery	Breast	10	8
P4	Female	34	Full-time job	2000~5000	Chemotherapy	Gynecology	1	20
P5	Male	42	Full-time job	2000~5000	Surgery	Lung	1	17
P6	Female	50	Retirement	2000~5000	Surgery	Lung	2	22
P7	Female	53	Retirement	<2000	Chemotherapy+ Surgery	Gastrointestinal	4	28
P8	Male	48	Not employed	<2000	Chemotherapy+ Surgery	Gastrointestinal	3	23
P9	Female	30	Full-time job	>5000	Surgery	Breast	1	25
P10	Male	60	Retirement	2000~5000	Surgery	Gastrointestinal	5	10

FCRI-SF score ≥ 13 indicates the presence of fear of cancer recurrence.