

## Supplementary materials

### Flux-based ozone risk assessment for plant injury index in three European cool-temperate deciduous tree species

YASUTOMO HOSHIKA<sup>1\*</sup>, ELISA CARRARI<sup>1</sup>, BARBARA MARIOTTI<sup>1</sup>, SOFIA MARTINI<sup>1</sup>, ALESSANDRA DE MARCO<sup>2</sup>, PIERRE SICARD<sup>3</sup> and ELENA PAOLETTI<sup>1</sup>

1. *Institute of Research on Terrestrial Ecosystems (IRET), National Research Council of Italy (CNR), Via Madonna del Piano, I-50019 Sesto Fiorentino, Italy*
2. *Italian National Agency for New Technologies, Energy and the Environment, C.R. Casaccia, Italy*
3. *ARGANS, Biot, France*

\*Corresponding author: Yasutomo Hoshika, Tel: +39-055-522-5949, Fax: +39-055-522-5920 E-mail: [yasutomo.hoshika@cnr.it](mailto:yasutomo.hoshika@cnr.it)

Supplementary figure and table




Figure S1. Comparison between measured and estimated  $g_s$  in *Alnus glutinosa*, *Sorbus aucuparia* and *Vaccinium myrtillus*. Linear regression analysis: \*\*\*  $p < 0.001$ ; RMSE, Root-Mean-Square Error.

