

## Article

# Fungi Associated with Horse-Chestnut Leaf Miner Moth *Cameraria ohridella* Mortality

Irena Nedveckytė<sup>1,\*</sup>, Dalė Pečiulytė<sup>2</sup> and Vincas Būda<sup>2</sup><sup>1</sup> Institute of Biosciences, Life Sciences Center, Vilnius University, Saulėtekio ave. 7, LT-10257 Vilnius, Lithuania<sup>2</sup> Institute of Ecology, Nature Research Centre, Akademijos 2, LT-08412 Vilnius, Lithuania; dale.peciulyte@gmail.com (D.P.); vincas.buda@gamtc.lt (V.B.)

\* Correspondence: irena.nedveckyte@gf.vu.lt

**Abstract:** The total mortality of the leaf-miner horse-chestnut pest, *Cameraria ohridella*, collected in nature, and the mortality associated with mycoses were assessed under laboratory conditions in stages: for eggs mortality rates of 9.78% and 61.97% were found, respectively; for caterpillars, 45.25% and 5.59%, respectively; and for pupae 21.22% and 100%, respectively. At the egg stage, *Cladosporus cladosporioides* caused mycosis most often (27% of all mycoses); at the caterpillar stage there was no pronounced predominant fungus species; at the pupal stage both *Cordyceps fumosorosea* and *Beauveria bassiana* (32% and 31%, respectively) were most dominant; whereas at the adult stage *Lecanicillium aphanocladii* (43%) were most dominant. *C. ohridella* moths remained the most vulnerable during the pupal and caterpillar stages. Maximum diversity of fungi associated with the leaf-miner moth was reached during the period of development inside the chestnut leaf (Shannon–Wiener index— $H'$  = 2.608 at the caterpillar stage,  $H'$  = 2.619 at the pupal stage), while the minimum was reached in the adult stage ( $H'$  = 1.757). In the caterpillar and pupa stages, saprophytic fungi were most often recorded. Comparative laboratory tests revealed novel properties of the fungus *L. aphanocladii*, its effectiveness as the leaf-miner moth's entomopathogen and its suitability for field application trials while developing environment-friendly methods for horse-chestnut pest control.

**Keywords:** leaf miner; *Cameraria ohridella*; entomopathogenic fungi; *Lecanicillium aphanocladii*

**Citation:** Nedveckytė, I.; Pečiulytė, D.; Būda, V. Fungi Associated with Horse-Chestnut Leaf Miner Moth *Cameraria ohridella* Mortality. *Forests* **2021**, *12*, 58. <https://doi.org/10.3390/f12010058>

Received: 14 December 2020

Accepted: 31 December 2020

Published: 5 January 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Horse-chestnut, *Aesculus hippocastanum* L. (Hippocastanaceae), is widely grown all over Europe in urban greenery, but since the 1980s the decorative value of these trees has fallen sharply due to the spread of the leaf-mining pest *Cameraria ohridella* Deschka and Dimic (Lepidoptera, Gracillariidae). *Cameraria ohridella* causes large-scale browning of leaves and pre-mature defoliation of horse-chestnut trees [1–5]. During the last 30 years, the leafminer spread over Europe, reaching Russia and Turkey [6]. In Lithuania, *C. ohridella* was registered for the first time in 2002 [7] and since then it has dispersed to a great extent and causes multiple infestations annually [8]. The moth's region of origin was obscure for a long time; however based on DNA analysis, the Balkan origin of the *C. ohridella* was revealed [9], and the species has been attributed to invasive ones in many countries. The horse-chestnut leaf miner has been widely studied, including moth biology and distribution [10–14], potential predators [15–18] and competition either with native leaf miners [19] or phytopathogenic fungi [20].

Control of *C. ohridella* populations is complicated due to the lifestyle of the pest: caterpillars develop inside the leaves [13] and host plants occur in urban territories where application of insecticides is unsuitable. The removal of the steadied leaves beneath the trees and subsequently either composting or burning reduces populations of natural enemies (*Chalcidoides* species) [21]. Application of systemic chemicals (e.g., imidacloprid, abamectin) by injection into the trunk of a tree is effective; however, wounds facilitate plant pathogen invasions and damages [22,23]. Removal of the adult moths with sticky

pheromone traps [17,23–25] can decrease pest populations; however, it is expensive and requires a large amount of effort and manual work.

In our opinion, the search for host-specific antagonists should be a priority. However, information on *C. ohridella*-related fungal entomopathogens remains scarce, and most investigations of fungi involved in moth mortality have been performed in a single country only, namely in the Czech Republic [26–30]. Fungi belonging to the genera of known fungal entomopathogens (*Beauveria* Vuill., *Lecanicillium* W. Gams and Zare, *Metarhizium* Sorokin, and *Paecilomyces* Bainier) were isolated from *C. ohridella* habitat and some strains showed entomopathogenic activity against pupae and moth caterpillars [31–35]. Unfortunately, information on fungi virulence throughout moths' developmental stages remains scarce.

The aim of the present study was to identify fungal entomopathogens associated with *C. ohridella* during all developmental stages, from eggs to pupae. The following tasks were set: (a) to examine diseased *C. ohridella* adults, collected from different *A. hippocastanum* trees; (b) to establish the mortality of *C. ohridella* eggs, caterpillars and pupae collected from the field focusing on fungi; (c) to isolate and identify fungi recovered from cadavers; (d) to carry out comparative efficiency tests of dominant fungus species in comparison with *C. ohridella* at different developmental stages.

## 2. Materials and Methods

### 2.1. Insects

Trunk and leaves from a lower crown of trees were examined mid-May to end-May, mid-August to mid-September and at leaf fall periods during 2009–2011, i.e., during the periods of the first and the second generations of *C. ohridella*. Each cadaver of adult moths (mainly found on tree trunks) was placed in a sterile Petri dish individually. Leaves were collected from five heavily infested, randomly selected trees (30 leaves per tree).

Leaves (used for egg treatment) were viewed under a stereomicroscope (Labomed Luxeo 2S Stereo Microscope;  $\times 10$  magnification) to find and determine segments with moth eggs. Leaf segments ( $2\text{--}3\text{ cm}^2$ ) with a known number of eggs were cut out and placed in Petri dishes (5 cm diameter) with sterile moistened cotton lumps and sealed with a lid to avoid desiccation. Eggs were incubated in the lab at  $20 \pm 2\text{ }^\circ\text{C}$ . Up to 150–250 eggs approximately were collected from each tree. The total number of eggs examined was 1451. After incubation, the total number of emerged caterpillars was counted and the egg mortality percentage was calculated. Eggs from which larvae did not hatch were viewed under the stereomicroscope and a light microscope (Motic B1 Series;  $\times 150$  and  $\times 1000$  magnifications) to check for possible fungal infection.

### 2.2. Isolation of Fungi

Leaf segments containing mines (each  $2.5\text{--}3\text{ cm}^2$  in square) were cut out following leaf surface sterilization by application of 0.5% (*v/v*) sodium hypochlorite (NaOCl). Each segment was treated with 75% ethanol for 1 min, followed by immersion for 2 min in sodium hypochlorite and again for 30 sec in 75% ethanol. Then, the segments were rinsed three times in sterile distilled water, and each was placed into a separate Petri dish (5.5 cm in diameter) supplied with a wet cotton lump. Dishes were sealed and incubated at  $20 \pm 2\text{ }^\circ\text{C}$  in the lab. Dishes were examined daily, and adult moth emergence was recorded. The same procedure was performed for the treatment of overwintered pupae, except that only areas of fully developed winter mines were cut from the leaves. Four replicates were prepared, each containing 30 mines. Mines containing caterpillar (or pupal) cadavers were aseptically dissected; each cadaver was taken out of a mine and placed on the moistened paper within a sterile Petri dish. Dishes were kept at  $25\text{ }^\circ\text{C}$  in the dark and examined daily for mycoses. A total of 632 caterpillars and 568 pupae within mines were tested.

Fungi were identified based on their morphology by using a low magnification stereomicroscope ( $\times 40$  magnification) and by preparing slides for light microscopy ( $\times 400$  and  $\times 1000$  magnifications). In cases when a fungi mycelium covered a cadaver and it did not produce conidia (or spores), small segments were dissected and plated on a half-strength

Potato Dextrose Agar (PDA) media supplemented with chloramphenicol ( $0.1 \text{ g L}^{-1} w/v$ ), and sub-cultured on three media: PDA, Sabouroud Dextrose Agar (SDA) and Czapek Dox Agar (CDA) (all from Liofilchem, Italy). Identification of fungi followed standard methods based on macro- and micro-morphological features [36–43]. All fungi species' names were revised according to <http://www.indexfungorum.org/>.

### 2.3. Insect Susceptibility to Fungi

#### 2.3.1. Fungi and Conidia Preparation

Four species of entomopathogenic fungi were used for comparative bioassay trials against *C. ohridella*: *Beauveria bassiana* (Bals.-Criv) Vuill., *Cordyceps fumosorosea* (Wize) Kepler, B. Shrestha and Spatafora (syn., *Isaria fumosorosea*), *Lecanicillium aphanocladii* Zare and W. Gams and *Lecanicillium psalliotae* (Treschew) Zare and W. Gams. All fungal strains are maintained in the Microorganisms Culture Collection at the Biodeterioration Research Laboratory of Nature Research Centre (Vilnius, Lithuania). The tested fungi were grown for 12–14 days on PDA plates; conidia were washed with 1 mL Tween 80 solution ( $0.01\%, v/v$ ). Three plates of each fungus were washed; washings were mixed and conidia suspension was adjusted to the desired concentration using a haemocytometer. Germination of conidia was over 95%. *Cameraria ohridella* eggs, naked pupae, and pupae inside the mines were treated with entomopathogenic fungi conidia suspensions at a concentration of  $3 \times 10^8$  conidia  $\text{mL}^{-1}$ .

#### 2.3.2. Egg Trials

Leaf segments ( $3 \times 3 \text{ cm}$ ) with a counted number of eggs were cut out of a leaf, and their surfaces were gently sterilized, placed into Petri dishes and sprayed with 1.5 mL of the appropriate fungus conidial suspension at  $3 \times 10^8$  conidia  $\text{mL}^{-1}$  concentration. Control Petri dishes were sprayed with 1.5 mL of distilled water. Leaf segments contained a different number of eggs (from 22 to 87); therefore, two or three segments were placed per dish to obtain ~100 eggs for each trial and each control. Each conidial concentration and control was tested three times (~100 eggs per replication, i.e., ~300 eggs in the test and ~300 eggs in control in total). Dishes with eggs were sealed and incubated for 3–5 days under  $20 \pm 2 \text{ }^\circ\text{C}$  in the laboratory. Emerged caterpillars were counted daily.

#### 2.3.3. Pupae Trials

Fallen tree leaves containing overwintered *C. ohridella* mines were washed with a large amount of water and dried-up. Leaf pieces containing mines were cut out (mine area only) and leaf surfaces were sterilized. Half of the sterilized pieces with mines were used directly for bioassay. The other half was aseptically opened; pupae were taken out and used for naked pupae bioassay. In each replicate, 30 pupae were placed on filter paper in a Petri dish (8.5 cm diameter) and sprayed with 1.5 mL of the conidia suspension of the appropriate fungus at a  $3 \times 10^8$  conidia  $\text{mL}^{-1}$  concentration. Control pupae were sprayed with the same volume of distilled water. Five replicates were tested. Pupae were incubated for 4 weeks in a humid chamber at 95% RH (high relative humidity was required to avoid desiccation of pupae) and  $22 \pm 2 \text{ }^\circ\text{C}$ . Pupae were examined weekly and emerged adults were counted.

### 2.4. Data Analyses

To evaluate fungal species diversity at each stage of the leaf-miner moth, the Shannon–Wiener index ( $H'$ ) was used [44]:  $H' = -\sum_{i=1}^S (p_i \ln p_i)$ , where  $S$  is the total number of species recorded and  $p_i$  is the frequency of occurrence of the  $i$ th species. Mortality percentage of the control was calculated, and the mean value of the five replicates (30 individuals each; total  $n = 150$ ) was presented, as well as standard error (SE). The mortality percentages of insects were normalized using  $\arcsin(\sqrt{x/100})$  transformation prior to the analysis of variance. Differences in the mean mortality percentage of the insect between different fungi trials, as well as between certain fungus trials and controls (untreated with fungi) were ascertained by variance analysis (one-way ANOVA) followed

by multiple comparisons of the means conducted using Tukey's test (SPSS 16 statistical package).

### 3. Results

#### 3.1. Fungi Associated with *C. ohridella* Mortality

##### 3.1.1. Adult Moths' Mycoses

Five samplings resulted in a collection of 349 adult moth cadavers, 109 of which had mycosis symptoms (Table 1). During incubation in the laboratory, mycelium development on the cadavers was species-dependent, and started after 2–3 days of incubation, reaching maturity within a week. Fungi were assigned to nine species from seven genera (Table 1). *Lecanicillium aphanocladii* was the most common species with 39.45% frequency of occurrence (FO), followed by *L. psalliotae* (FO = 16.51%) and *Trichothecium roseum* (Pers.) Link (FO = 15.6%). The fungi species diversity index for adult moth cadavers was  $H' = 1.757$ .

**Table 1.** Mycosis symptoms of *Cameraria ohridella* at different stages: from egg to adult.

	Adult Moths	Eggs	Caterpillars	Pupae
Total number (TN) of specimen tested	349	1451	632	568
TN (and%) of dead individuals	349 (100)	142 (9.78)	286 (45.25)	183 (21.22)
TN (and%) with mycosis symptoms	109 (31.23)	88 (61.97)	36 (5.59)	183 (100)
	Number of infected individuals and FO *			
Fungi species	Adult moths	Eggs	Caterpillars	Pupae
<i>Akanthomyces lecanii</i> (Zimm.) Spatafora, Kepler and B. Shrestha				9 (4.91)
<i>Alternaria alternata</i> (Fr.) Keissl.			2 (5.56)	7 (3.82)
<i>Aspergillus parasiticus</i> Speare			2 (5.56)	3 (1.64)
<i>Beauveria bassiana</i> (Bals.-Criv.) Vuill.	6 (5.5)	11 (12.5)	3 (8.33)	31 (16.94)
<i>Cladosporium cladosporioides</i> (Fresen.) G.A. de Vries	11 (10.09)	27 (30.68)	3 (8.33)	11 (6.01)
<i>Clonostachys rosea</i> (Link) Schroers, Samuels, Sefert and W. Gams				8 (4.37)
<i>Colletotrichum gloeosporioides</i> (Penz.) Penz. and Sacc.	2 (1.84)	1 (1.14)	2 (5.56)	
<i>Cordyceps fumosorosea</i> (Wize) Kepler, B. Shrestha and Spatafora	3 (2.75)	2 (2.27)		32 (17.49)
<i>Cordyceps farinosa</i> (Holmsk.) Kepler, B. Shrestha and Spatafora		3 (3.41)		6 (3.28)
<i>Cordyceps tenuipes</i> (Peck) Kepler, B. Shrestha and Spatafora				4 (2.19)
<i>Erysiphe flexuosa</i> (Peck) U. Braun and S. Takam.			3 (8.33)	
<i>Fusarium proliferatum</i> (Matsush.) Nirenberg	2 (1.84)			4 (2.18)
<i>Fusarium oxysporum</i> Schltdl.		6 (6.82)		3 (1.64)
<i>Fusarium fujikuroi</i> Nirenberg	7 (6.42)		1 (2.77)	
<i>Harzia acremonioides</i> (Harz) Costantin			2 (5.56)	
<i>Lecanicillium aphanocladii</i> Zare and W. Gams	43 (39.45)	13 (14.77)	2 (5.56)	2 (1.09)
<i>Lecanicillium psalliotae</i> (Treschew) Zare and W. Gams	18 (16.51)	4 (4.55)		16 (8.74)
<i>Metarhizium anisopliae</i> (Metschn.) Sorokin				4 (2.19)
<i>Microascus brevicaulis</i> S.P. Abbott				2 (1.09)
<i>Oedocephalum glomerulosum</i> (Bull.) Sacc.			3 (8.33)	
<i>Penicillium citrinum</i> Thom		5 (5.68)	1 (2.77)	
<i>Penicillium chrysogenum</i> Thom		2 (2.27)	1 (2.77)	
<i>Penicillium oxalicum</i> Currie and Thom		4 (4.55)		2 (1.09)
<i>Sarocladium strictum</i> (W. Gams) Summerb.			4 (11.12)	16 (8.74)
<i>Stachybotrys chartarum</i> (Ehrenb.) S. Hughes				3 (1.64)
<i>Trichoderma harzianum</i> Rifai		2 (2.27)	2 (5.56)	4 (2.18)
<i>Trichothecium roseum</i> (Pers.) Link	17 (15.6)	8 (9.09)	5 (13.89)	
Other (unidentified four morphotypes)				16 (8.74)
Total number of species	9	13	15	19
Total number of genera	7	9	14	15
Shannon-Wiener diversity index ( $H'$ )	1.757	2.174	2.608	2.619

\* FO—frequency of occurrence (in brackets) was calculated as the percentage of cadavers with the separate fungus of the total number of cadavers with mycosis symptoms.

### 3.1.2. Egg Mortality and Mycoses

A total of 1451 eggs were incubated and checked for hatching daily in the laboratory. The eggs started hatching within two days of incubation and hatched approximately after a week. Only 9.78% of the eggs (142 out of 1451 incubated) did not hatch; growth of fungi was detected on 61.97% of the unhatched eggs and on 6.06% of all eggs (88 out of 142 and out of 1451, respectively) (Table 1). In total, 13 fungi species from nine genera were identified. The species diversity index  $H'$  of fungi found on *C. ohridella* eggs was equal to 2.174. *Cladosporium cladosporioides* was the most frequent isolate (recovered from 30.68% of unhatched eggs), followed by *L. aphanocladii* (14.77%) and *B. bassiana* (12.5%).

### 3.1.3. Caterpillar Mortality and Mycoses

Six hundred thirty-two caterpillars inside the mines were incubated in the laboratory. During the first three days, only 19 (3.0%) dead caterpillars were registered. Mortality of caterpillars reached 4.6% within 6 d, and 5.9% within 9 d of incubation. During two weeks of rearing, only 54.75% of the caterpillars (346 out of 632 incubated) reached the adult moth stage, and thus the total caterpillar mortality was considered as being high. Mortality of 5.59% (36 out of 286) caterpillars was induced by fungi. The fungi species diversity index ( $H' = 2.608$ ) determined for the caterpillars was higher than that for the adult moths ( $H' = 1.757$ ). The frequency of occurrence of any fungus species ranged from 2.77% to 13.89% (Table 1). *Trichothecium roseum* (FO = 13.89%), *Sarocladium strictum* (FO = 11.12%) and *Oedocephalum glomerulosum* (FO = 8.33%) were the most common fungi species isolated. Presence of bacteria and yeasts was noted as well; however those were not analyzed.

### 3.1.4. Pupal Mortality and Mycoses


A total of 568 mines cut from the collected fallen leaves were incubated with overwintered pupae. Adult moths did not emerge from 21.22% of the pupae (183 emerged out of 568 pupae) (Table 1), and fungi were recovered from all diseased pupae (100%). Complexes of fungi species rather than a single species were detected quite often, approximately in 33% of the diseased pupae. The diversity index of the fungi species isolated from the pupae  $H'$  was equal to 2.619, and its value was the highest established in the present analysis. Nineteen fungal species from 15 genera were isolated (Table 1). *Cordyceps fumosorosea* (recovered from 17.49% cadavers), *B. bassiana* (from 16.94%) and *L. psalliotae* (from 8.74%) as well as *S. strictum* (from 8.74% cadavers) were the most common fungi species isolated from the diseased pupae.

## 3.2. Groups of Fungal Entomopathogens

Proportions of recovered fungi species grouped according to their ecology are presented in Figure 1. The highest relative abundance of generalist entomopathogenic fungi (GEF) was revealed in cadavers of *C. ohridella* adult moths (~65%). GEF proportion was smaller in pupae, and even smaller in eggs, comprising 46% and 38%, respectively. As in the case of GEF, the percentage of opportunistic entomopathogenic fungi (OEF) varied depending on the insect development stage. The highest percent of OEF was found in *C. ohridella* eggs and caterpillars (in 48% and 33% of cadavers, respectively), where the lowest abundance of GEF was recorded (38% and 28%, respectively).

As well as the relative abundance of ecological groups, the number of species within the GEF group varied depending on the moth developmental stage (Table 2). *Lecanicillium aphanocladii* prevailed among all isolated entomopathogens (28.30%). Other more common entomopathogenic species isolated were *B. bassiana* (24.06% of isolates), *L. psalliotae* (17.93%) and *C. fumosorosea* (17.45%).





**Figure 1.** Percentage of generalist entomopathogenic fungi (GEF), opportunistic entomopathogenic fungi (OEF) and other saprotrophic fungi (SF) among the fungi recovered from *C. ohridella* moths at different stages of development: adult moths (a), eggs (b), caterpillars (c), and pupae (d).


**Table 2.** Generalist entomopathogenic fungi (GEF) from diseased *Cameraria ohridella* moths at different developmental stages.

GEF Species	Number of Infected Moth Individuals				Total Number (Percentage, %)
	Adult	Egg	Caterpillar	Pupae	
<i>Akanthomyces lecanii</i>				9	9 (4.25)
<i>Beauveria bassiana</i>	6	11	3	31	51 (24.06)
<i>Cordyceps fumosorosea</i>	3	2		32	37 (17.45)
<i>Cordyceps farinosa</i>		3		6	9 (4.25)
<i>Cordyceps tenuipes</i>				4	4 (1.89)
<i>Lecanicillium aphanocladii</i>	43	13	2	2	60 (28.30)
<i>Lecanicillium psalliotae</i>	18	4		16	38 (17.93)
<i>Metarhizium anisopliae</i>				4	4 (1.89)
Total number	70	33	5	104	212 (100)

### 3.3. Susceptibility of *C. ohridella* to Fungal Entomopathogens (Bioassay Tests)

Of the generalist entomopathogenic fungi, which most often damage *C. ohridella* at various stages of development in nature (Table 2), four species were selected for a comparative test: *B. bassiana*, *C. fumosorosea*, *L. aphanocladii* and *L. psalliotae*. The test would have been appropriate for the caterpillar stage of the moth, since this stage is responsible for the damage to the horse-chestnut leaves. However, since caterpillars live inside the leaf, they are very well-protected from any affect, including that of fungi (see Table 2); therefore, the egg and pupal stages of the moth were selected for the comparative analysis.

*Lecanicillium aphanocladii* was among the most virulent fungi species tested on *C. ohridella* (Figure 2). These fungi caused mortality of  $76.83 \pm 4.1\%$  eggs and that was 1.8, 2.0, and 2.8 times more compared to the effects caused either by *B. bassiana*, *C. fumosorosea* or *L. psalliotae*, respectively. *Beauveria bassiana* and *C. fumosorosea* showed similar virulence towards the eggs: induced mortality reached  $57.09 \pm 6.18\%$  and  $53.13 \pm 5.14\%$ , respectively, which was significantly less ( $p < 0.05$ ) from that induced by *L. aphanocladii* ( $76.83 \pm 4.1\%$ ).



**Figure 2.** Mortality of *C. ohridella* eggs, naked pupae and pupae within the mines treated with *Lecanicillium aphanocladii*, *L. psalliotae*, *Beauveria bassiana* and *Cordyceps fumosorosea* at  $3 \times 10^8$  conidia/mL concentrations in the sprayed suspension. Data are presented as the mean of five treatments and standard error (SE) (bars). Different letters at columns indicate statistically significant differences (Tukey's test,  $p < 0.05$ ).

Mortality of the naked pupae significantly ( $p < 0.05$ ) increased comprising,  $57.09 \pm 2.71\%$ ,  $65.28 \pm 4.39\%$ ,  $76.81 \pm 6.52\%$  and  $87.64 \pm 2.76\%$ , respectively, in the *B. bassiana*, *C. fumosorosea*, *L. psalliotae* and *L. aphanocladii* treatments. All four tested fungi showed intensive mycelium growth on the cadavers already after 3–4 days of incubation. The lowest mortality was recorded when pupae inside their mines were sprayed with fungal conidia. Mortality following spraying (concentration  $3 \times 10^8$  conidia  $\text{mL}^{-1}$ ), varied from  $34.29 \pm 13.91\%$  to  $53.11 \pm 4.79\%$  in *B. bassiana* and *L. psalliotae* treatments, respectively. *L. aphanocladii* showed a slightly higher virulence trend ( $56.82 \pm 7.39\%$ ); however, the difference was not statistically significant ( $p > 0.05$ ).

#### 4. Discussion

The analysis of mortality and fungus infestation of the invasive pest leaf miner moth, *C. ohridella*, collected during various stages of development in nature allows to evaluate the role of entomopathogenic fungi in the regulation of its population. In the present

study, eggs, caterpillars and pupae were transferred from nature into development-friendly laboratory conditions, and the results both on total mortality and mortality related to mycoses led to lower estimations than those in nature. There is no doubt that longer exposure should increase fungal infection. On the other hand, predators would remove some infected insects and those would not enter the records of the infestation.

The total mortality of *C. ohridella* eggs was 9.78%, that of larvae was 45.25% and that of overwintering pupae was 21.22%. There are data on egg and pupae mortality obtained under natural conditions, and the values are much higher, reaching 30% for eggs [14] and from 53% up to 97.25% for overwintered pupae [5,14,45]. Unfortunately, due to lack of data both on summer pupae and adults of reproductive age, estimation of general mortality (or survival) throughout the whole life cycle of *C. ohridella* seems not possible. However, there is no doubt that the total survival of preimaginal stages of the moth is very high during the summer period, as indicated by the poor condition of the host-plant leaves. The total percentage of caterpillar mortality was high (45.25%) in comparison with the mortality of eggs and pupae. However, only 5.59% of the dead caterpillars showed mycoses symptoms. Perhaps this could be the result of host-plant leaves, as it is known that the leaves of *A. hippocastanum* contain defensive substances (anthocyanins, phenols and flavonols) which might reduce the body mass and survival of caterpillars within leaf tissues [46,47].

The data obtained on fungi related to *C. ohridella* revealed 27 species from 19 genera, including eight species of entomopathogenic fungi, seven species of secondary (opportunistic) fungal pathogens and 12 species of saprotrophic fungi (Table 1). Common insect pathogens prevailed in the adult and pupal cadavers, comprising 65% and 46% of all isolates, respectively. A comparable proportion of the generalist entomopathogens was revealed in eggs (38%) and caterpillars (28%).

Depending on the moth development stage, the prevailing fungus species detected differed (except for the caterpillar stage, where the detection rate of any of 15 species varied from 2.77% to 13.89% only). At the egg stage, mycoses were mainly caused by *Cladosporium cladosporioides* (30.68% of all mycoses), at the pupa stage *Cordyceps fumosorosea* and *Beauveria bassiana* prevailed (17.49% and 16.94%, respectively), and *Lecanicillium aphanocladii* (39.45%) prevailed at the adult stage. The latter fungus species was recorded for the first time in Lithuania, and thus its identification was checked and confirmed by application of molecular methods and published in a separate paper [48].

Among entomopathogenic species, *L. aphanocladii* (28.30%), *B. bassiana* (24.06%), *L. psalliotae* (17.93%) and *C. fumosorosea* (17.45%) were the most abundant (Table 2). Information on the variety of entomopathogenic fungi involved in *C. ohridella* mortality is very scarce. Investigations of moth-related fungi and possible fungal pathogens have been performed in the Czech Republic [26–30]. Positive insecticidal effects of *L. muscarium* (Petch) Zare and W. Gams against *C. ohridella* caterpillars [33,34] and pupae [32] were reported. *Cordyceps fumosorosea* (syn., *Isaria fumosorosea* or *Paecilomyces fumosoroseus*), *B. bassiana*, and *M. anisopliae* were demonstrated to be pathogenic to the invasive moth as well [31–34].

Some opportunistic fungi were abundant as generalist fungal entomopathogens. They comprised 48% and 33% of the total number of fungi isolated from the *C. ohridella* eggs and caterpillars, respectively. *Cladosporium cladosporioides* and *T. roseum* species were the opportunistic fungal pathogens most often recovered. We found the latter fungi to be common on *A. hippocastanum* foliage (data not presented); thus we predict that direct contact could be the reason of the high infection of *C. ohridella* eggs. *Sarocladium strictum* was common among isolates from caterpillar and pupal cadavers as well. The fungus is known as the most common fungal endophyte of grasses, has been found on leaf surfaces of vascular plants, was frequently observed on other primary fungal colonizers like rusts or powdery mildews, and is widely distributed in the atmosphere [43]. The species is also known as related to the attack on insects [49,50].

Comparative laboratory tests demonstrated (see Figure 2) that the *L. aphanocladii* entomopathogenic efficacy towards *C. ohridella* is equivalent to other well-known ento-



mopathogens: *C. fumosorosea*, *B. bassiana* and *L. psalliotae*. No statistically significant difference was found between the pathogenic potentials of *B. bassiana* and *C. fumosorosea*. Although both fungi increased the mortality of eggs and naked pupae by up to 53.13–65.28%, their induced mortality of pupae inside their mines (34.29–46.57%) was significantly lower than the mortality induced by *L. aphanocladii* and *L. psalliotae* (53.11–56.82%). A surprisingly high difference was revealed in the pathogenic potential of the species within the same *Lecanicillium* genus. Like *L. aphanocladii*, *L. psalliotae* conidia were highly pathogenic to naked pupae, but not to eggs. Based on the results of the present investigation, it cannot be firmly stated that *L. aphanocladii* was the strongest inhibitor of *C. ohridella* development; data on the comparative bioassay of the four fungi tested did not reveal significant interspecific pathogenicity differences, except at the egg stage, when *L. aphanocladii* was the most active. In conclusion, *L. aphanocladii* was commonly recovered from *C. ohridella* cadavers and showed high pathogenic potential in the bioassay. The fungus was found to be associated with *A. hippocastanum* foliage and was particularly common on the leaf surfaces with a higher density of powdery mildew disease inducing fungus, *Erysiphe flexuosa* (Peck) U. Braun and S. T. Takamatsu (syn. *Uncinula flexuosa*) (data not presented). We hypothesize that *L. aphanocladii* is related to *E. flexuosa* development, like three other closely related *Lecanicillium* species (*L. attenuatum*, *L. lecanii* (now renamed *Akanthomyces lecanii*) and *L. longisporum*), which are known to be active both against some insects and powdery mildew [51,52]. One more fungus species of the same genus, known as generalist mycoparasite, *L. muscarium* (formally known as *V. lecanii*) inhibiting powdery mildew [53], was discovered and confirmed as a *C. ohridella* pathogen [34]. This indicates that investigation of *L. aphanocladii*'s relationship with other fungal species could offer perspective and provide more opportunities to develop environment-friendly methods for *C. ohridella* pest control by means of entomopathogens.

## 5. Conclusions

The total mortality of the leaf-miner horse-chestnut pest, *Cameraria ohridella*, collected in nature, and the mortality associated with mycoses, was assessed under laboratory conditions for eggs (9.78% and 61.97%, respectively) for caterpillars (45.25% and 5.59%, respectively) and for pupae (21.22% and 100%, respectively). At the egg stage, *Cladosporus cladosporioides* caused mycosis most often (27% of all mycoses); at the caterpillar stage there was no pronounced predominant fungus species; at the pupal stage both *Cordyceps fumosorosea* and *Beauveria bassiana* were dominant (32% and 31%, respectively); and at the adult stage *Lecanicillium aphanocladii* was dominant (43%). *C. ohridella* moths remained the most vulnerable during the pupal and caterpillar stages. Maximum diversity of fungi associated with the leaf-miner moth was reached during the period of development inside the chestnut leaf (Shannon–Wiener index,  $H' = 2.608$  at the caterpillar stage and  $H' = 2.619$  at the pupal stage), whereas it reached a minimum at the adult stage ( $H' = 1.757$ ). Comparative laboratory tests revealed novel properties of the fungus *L. aphanocladii*, its effectiveness as the leaf-miner moth's entomopathogen and its suitability for field application trials while developing environment-friendly methods needed for horse-chestnut pest control.

**Author Contributions:** Investigation—D.P., I.N.; conceptualization—V.B.; methodology—D.P., writing—original draft, D.P., I.N.; writing—review and editing, I.N., V.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the European Social Fund/European Regional Development Fund under a grant agreement with the Research Council of Lithuania (LMTLT), grant number 09.3.3-LMT-K-712-01-0099, and the Article Processing Charges (APC) was funded by 09.3.3-LMT-K-712-01-0099 and the Institute of Biosciences.

**Institutional Review Board Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author.

**Acknowledgments:** The technical assistance of Vaidilutė Dirginčiūtė-Volodkienė (Institute of Botany, Nature Research Centre, Vilnius, Lithuania) is highly appreciated.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Backhaus, G.F.; Wulf, A.; Kehr, R.; Schröder, T. Die Rosskastanien-Miniermotte (*Cameraria ohridella*): Biologie, Verbreitung und Gegenmassnahmen. *Nachrichtenbl. Deut. Pflanzenschutzbd.* **2002**, *54*, 56–62.
- Gilbert, M.; Grégoire, J.C. Visual, semi-quantitative assessments allow accurate estimates of leafminer population densities: An example comparing image processing and visual evaluation of damage by the horse chestnut leafminer *Cameraria ohridella* (Lep., Gracillariidae). *J. Appl. Entomol.* **2003**, *127*, 354–359. [[CrossRef](#)]
- Heitland, W. Controcam—control of *Cameraria*. *Nachr. Dtsch. Pflanzenschutzbd.* **2006**, *58*, 251–252.
- Heitland, W.; Kopelke, J.P.; Freise, J.; Metzger, J. Ein Kleinschmetterling erobert Europa—die Rosskastanien-Miniermotte *Cameraria ohridella*. *Nat. Mus.* **1999**, *129*, 186–195.
- Syeryebryennikov, B. *Ecology and Control of Horse Chestnut Leaf Miner (Cameraria Ohridella)*; Shmalhausen Institute of Zoology: Kiev, Ukraine, 2008; pp. 1–67.
- Lees, D.C.; Lack, H.W.; Rougerie, R.; Hernandez-Lopez, A.; Raus, T.; Avtzis, N.D.; Augustin, S.; Lopez-Vaamonde, C. Tracking origins of invasive herbivores through herbaria and archival DNA: The case of the horse-chestnut leaf miner. *Front. Ecol. Environ.* **2011**, *9*, 322–328. [[CrossRef](#)]
- Ivinskis, P.; Rimšaitė, J. The horse-chestnut leafminer (*Cameraria ohridella* Deschka & Dimic 1986) (Lepidoptera, Gracillariidae) in Lithuania. *Acta Zool. Litu.* **2006**, *16*, 323–327. [[CrossRef](#)]
- Stankevičienė, A.; Snieškienė, V.; Lugauskas, A. *Erysipher flexuosa*—The new pathogen of *Aesculus hippocastanum* in Lithuania. *Phytopathologia* **2010**, *56*, 67–71. Available online: <https://hdl.handle.net/20.500.12259/50256> (accessed on 10 December 2020).
- Valade, R.; Kenis, M.; Hernandez-Lopez, A.; Augustin, S.; Mari Mena, N.; Magnoux, E.; Rougerie, R.; Lakatos, F.; Roques, A.; Lopez-Vaamonde, C. Mitochondrial and microsatellite DNA markers reveal a Balkan origin for the highly invasive horse-chestnut leaf miner *Cameraria ohridella* (Lepidoptera, Gracillariidae). *Mol. Ecol.* **2009**, *18*, 3458–3470. [[CrossRef](#)]
- Pschorn-Walcher, H. Freiland-Biologie der eingeschleppten Roßkastanien-Miniermotte *Cameraria ohridella* Deschka et Dimic (Lep., Gracillariidae) im Wienerwald. *Linz. Biol. Beiträge* **1994**, *26*, 633–642. Available online: [https://scholar.google.com/scholar?hl=lt&as\\_sdt=0%2C5&q=Freiland-Biologie+der+eingeschleppten+Ro%2C3%9Fkastanien-Miniermotte+Cameraria+ohridella+Deschka+et+Dimic+%28Lep.%2C+Gracillariidae%29+im+Wienerwald&btnG=](https://scholar.google.com/scholar?hl=lt&as_sdt=0%2C5&q=Freiland-Biologie+der+eingeschleppten+Ro%2C3%9Fkastanien-Miniermotte+Cameraria+ohridella+Deschka+et+Dimic+%28Lep.%2C+Gracillariidae%29+im+Wienerwald&btnG=) (accessed on 10 December 2020).
- Skuhravý, V. Zur Kenntnis der Blattminen-Motte *Cameraria ohridella* Desch. & Dim. (Lep., Lithocolletidae) an *Aesculus hippocastanum* L. in der Tschechischen Republik. *Anz. Schädlingsk* **1998**, *71*, 82–84. Available online: [https://scholar.google.com/scholar?hl=lt&as\\_sdt=0%2C5&q=Zur+Kenntnis+der+Blattminen-Motte+Cameraria+ohridella+&btnG=](https://scholar.google.com/scholar?hl=lt&as_sdt=0%2C5&q=Zur+Kenntnis+der+Blattminen-Motte+Cameraria+ohridella+&btnG=) (accessed on 10 December 2020).
- Šefrová, H.; Laštuvka, Z. Dispersal of the horse chestnut leafminer, *Cameraria ohridella* Deschka & Dimic, 1986, in Europe: Its course, ways and causes (Lepidoptera: Gracillariidae). *Zeitschrift Angew.* **2001**, *111*, 194–198.
- Freise, J.; Heitland, W. Bionomics of the horse-chestnut leaf miner *Cameraria ohridella* Deschka & Dimic 1986, a pest on *Aesculus hippocastanum* in Europe (Insecta, Lepidoptera, Gracillariidae). *Senckenb. Biol.* **2004**, *84*, 61–80.
- Girardoz, S.; Quicke, D.L.J.; Kenis, M. Factors favouring the development and maintenance of outbreaks in an invasive leaf miner *Cameraria ohridella* (Lepidoptera: Gracillariidae): A life table study. *Agric. For. Entomol.* **2007**, *9*, 141–158. [[CrossRef](#)]
- Grabenweger, G.; Kehrl, P.; Schlick-Steiner, B.; Steiner, F.; Stolz, M.; Bacher, S. Predator complex of the horse chestnut leafminer *Cameraria ohridella*: Identification and impact assessment. *J. Appl. Entomol.* **2005**, *129*, 353–362. [[CrossRef](#)]
- Grabenweger, G.; Hopp, H.; Schmolling, S.; Koch, T.; Balder, H.; Jäckel, B. Laboratory rearing and biological parameters of the eulophid *Pnigalio agraulis*, a parasitoid of *Cameraria ohridella*. *J. Appl. Entomol.* **2009**, *133*, 1–9. [[CrossRef](#)]
- Volter, L.; Kenis, M. Parasitoid complex and parasitism rates of the horse chestnut leafminer, *Cameraria ohridella* (Lepidoptera: Gracillariidae) in the Czech Republic, Slovakia and Slovenia. *Eur. J. Entomol.* **2006**, *103*, 365–370. [[CrossRef](#)]
- Ferracini, C.; Alma, A. Evaluation of the community of native eulophid parasitoids on *Cameraria ohridella* Deschka and Dimic in Urban areas. *Environ. Entomol.* **2007**, *36*, 1147–1153. [[CrossRef](#)]
- Péré, C.; Augustin, S.; Tomov, R.; Peng, L.H.; Turlings, T.C.J.; Kenis, M. Species richness and abundance of native leaf miners are affected by the presence of the invasive horse-chestnut leaf miner. *Biol. Invasions* **2010**, *12*, 1011–1021. [[CrossRef](#)]
- Jagiello, R.; Łakomy, P.; Łukowski, A.; Giertych, M.J. Spreading-the-risk hypothesis may explain *Cameraria ohridella* oviposition in relation to leaf blotch disease. *Arthropod Plant Interact.* **2019**, *13*, 787–795. [[CrossRef](#)]
- Kehrl, P.; Lehmann, M.; Bacher, S. Mass-emergence devices: A biocontrol technique for conservation and augmentation of parasitoids. *Biol. Control* **2005**, *32*, 191–199. [[CrossRef](#)]
- Baraniak, E.; Walczak, U.; Zduniak, P. Appearance and migration of the horse-chestnut leafminer *Cameraria ohridella* in relation to city size and leaf-raking, using the example of two cities in Western Poland. *J. Pest. Sci.* **2005**, *78*, 145–149. [[CrossRef](#)]
- Ferracini, C.; Alma, A. How to preserve horse chestnut trees from *Cameraria ohridella* in the urban environment. *Crop Prot.* **2008**, *27*, 1251–1255. [[CrossRef](#)]
- Kindl, J.; Kalinová, B.; Freise, J.; Heitland, W.; Augustin, S.; Guichard, S.; Avtzis, N.; Svatoš, A. Monitoring the population dynamics of the horse chestnut leafminer *Cameraria ohridella* with a synthetic pheromone in Europe. *Plant Prot. Sci.* **2002**, *38*, 131–137. [[CrossRef](#)]

25. Svatoš, A.; Kaliniová, B.; Hrdý, I. *Cameraria ohridella*: 10 years of sex pheromone and kairomone research. *J. Appl. Entomol.* **2009**, *133*, 319–327. [[CrossRef](#)]
26. Zemek, R.; Prenerova, E.; Weyda, F. The first record of entomopathogenic fungus *Paecilomyces fumosoroseus* (Deuteromycota: Hyphomycetes) on the hibernating pupae of *Cameraria ohridella* (Lepidoptera: Gracillariidae). *Entomol. Res.* **2007**, *37*, 135–136.
27. Zemek, R.; Prenerova, E.; Volter, L.; Weyda, F.; Skuhravy, V. Perspectives for the biological control of *Cameraria ohridella*. *Commun. Agric. Appl. Biol. Sci.* **2007**, *72*, 521–526.
28. Prenerova, E.; Zemek, R.; Weyda, F.; Volter, L. Sampling and occurrence of entomopathogenic fungi in soil from *Cameraria ohridella* (Lepidoptera: Gracillariidae) habitats. *IOBS/WPRS Bull.* **2008**, *31*, 204.
29. Prenerova, E.; Zemek, R.; Weyda, F.; Volter, L. Entomopathogenic fungi isolated from soil in the vicinity of *Cameraria ohridella* infested chestnut trees. *IOBS/WPRS Bull.* **2009**, *45*, 321–324.
30. Awad, M.; Prenerova, E.; Volter, L.; Zemek, R. Evaluation of *Isaria fumosorosea* CCM 8367 for control of *Cameraria ohridella*, and effects on beneficial parasitoids. In Proceedings of the 43rd Annual Meeting of the Society for Invertebrate Pathology, Trabzon, Turkey, 11–15 July 2010. 138p.
31. Richter, D.; Sermann, H.; Jäckel, B.; Büttner, C.B. Pathogenicity of entomopathogenic fungi on hibernating pupae of *Cameraria ohridella* Ceschka & Dimic 1986 (Lepidoptera, Gracillariidae). Part 1: Pathogenicity against the naked pupa. *Commun. Agric. Appl. Biol. Sci.* **2007**, *72*, 399–410. Available online: <https://europepmc.org/article/med/18399468> (accessed on 10 December 2020).
32. Richter, D.; Sermann, H.; Jäckel, B.; Büttner, C.B. Pathogenicity of entomopathogenic fungi on hibernating pupae of *Cameraria ohridella* Descha & Dimic 1986 (Lepidoptera, Gracillariidae). Part 2: Efficacy of entomopathogenic fungi against pupa in her pupas cell. *Commun. Agric. Appl. Biol. Sci.* **2007**, *72*, 411–421. Available online: <https://europepmc.org/article/med/18399469> (accessed on 10 December 2020).
33. Kalmus, M.; Sermann, H.; Lerche, S.; Büttner, C. Efficiency of the entomopathogenic fungus *Lecanicillium muscarium* on hibernating pupae of *Cameraria ohridella* Descha & Dimic, 1986 (Lepidoptera, Gracillariidae). *IOBS/WPRS Bull.* **2008**, *31*, 223–227.
34. Lerche, S.; Sermann, H.; Büttner, C. Persistence of the entomopathogenic fungus *Lecanicillium muscarium* Zare & Gams under ambient conditions in the field. *Commun. Agric. Appl. Biol. Sci.* **2009**, *74*, 353–356. [[PubMed](#)]
35. Schemmer, R.; Chládeková, P.; Medo, J.; Barta, M. Natural prevalence of entomopathogenic fungi in hibernating pupae of *Cameraria ohridella* (Lepidoptera: Gracillariidae) and virulence of selected isolates. *Plant Protect. Sci.* **2016**, *52*, 199–208. [[CrossRef](#)]
36. Gams, W. *Cephalosporium-Artige Schimmelpilze (Hyphomycetes)*; Gustav Fisher Verlag: Stuttgart, Germany, 1971; 262p.
37. Samson, R.A. *Paecilomyces and Some Allied Hyphomycetes*; Studies in Mycology; Centraalbureau voor Schimmelcultures: Baarn, The Netherlands, 1974; Volume 6, 119p.
38. Matsushima, T. *Icones Microfungorum a Matsushima Lectorum*; Takashi Matsushima: Kobe, Japan, 1975; 209p.
39. Pitt, J.I. *A Laboratory Guide to Common Penicillium Species*; Commonwealth Scientific and Industrial Research organization, Division of Food Research: Canberra, Australia, 1985; 122p.
40. Kiffer, E.; Morelet, M. *The Deuteromycetes, Mitosporic Fungi: Classification and Generic Keys*; Science Publishers Inc.: Enfield, CT, USA, 2000; 274p.
41. Zare, R.; Gams, W. A revision of *Verticillium* section Prostrata IV. The genus *Lecanicillium* and the new genus *Simplicillium*. *Nova Hedwig.* **2001**, *73*, 1–50. Available online: <https://www.cabdirect.org/cabdirect/abstract/20023039635> (accessed on 10 December 2020).
42. Sung, G.H.; Hywel-Jones, N.L.; Sung, J.M.; Luangsa-ard, J.J.; Shrestha, B.; Spatafora, J.W. Phylogenetic classification of *Cordyceps* and the clavicipitaceae fungi. *Stud. Mycol.* **2007**, *57*, 1–64. [[CrossRef](#)] [[PubMed](#)]
43. Domsch, K.H.; Gams, W.; Anderson, T.H. *Compendium of Soil Fungi*, 2nd ed.; IHW-Verlag: Eching, Germany, 2007; Volume 2, 672p. [[CrossRef](#)]
44. Krebs, C.J. *Ecological Methodology*; Harper and Row Publisher: New York, NY, USA, 1989; 645p.
45. Girardoz, S.; Tomov, R.; Eschen, R.; Quicke, D.L.J.; Kenis, M. Two methods to assess the mortality factors affecting larvae and pupae of *Cameraria ohridella* in leaves of *Aesculus hippocastanum* in Switzerland and Bulgaria. *Bull. Entomol. Res.* **2007**, *97*, 445–453. [[CrossRef](#)] [[PubMed](#)]
46. Paterska, M.; Bandurska, H.; Wysłouch, J.; Molińska-Glura, M.; Moliński, K. Chemical composition of horse-chestnut (*Aesculus*) leaves and their susceptibility to chestnut leaf miner *Cameraria ohridella* Deschka & Dimić. *Acta Physiol. Plant.* **2017**, *39*, 1–16. [[CrossRef](#)]
47. Jagiełło, R.; Baraniak, E.; Guzicka, M.; Karolewski, P.; Łukowski, A.; Giertych, M.J. One step closer to understanding the ecology of *Cameraria ohridella* (Lepidoptera: Gracillariidae): The effects of light conditions. *Eur. J. Entomol.* **2019**, *116*, 42–51. [[CrossRef](#)]
48. Pečiulytė, D.; Kačergius, A. *Lecanicillium Aphanocladii*—A New Species to the Mycoflora of Lithuania and a New Pathogen of Tree Leaves Mining Insects. *Bot. Lith.* **2012**, *18*, 133–146. [[CrossRef](#)]
49. Vidal, S. Changes in suitability of tomato for whiteflies mediated by a non-pathogenic endophytic fungus. *Entomol. Exp. Appl.* **1996**, *80*, 272–274. [[CrossRef](#)]
50. Jallow, M.F.A.; Gobena, D.D.; Vidal, S. Influence of an endophytic fungus on host plant selection by a polyphagous moth via volatile spectrum changes. *Arthropod Plant Interact.* **2008**, *2*, 53–62. [[CrossRef](#)]
51. Askary, H.; Carriere, Y.; Belanger, R.R.; Brodeur, J. Pathogenicity of the fungus *Verticillium lecanii* to aphids and powdery mildew. *Biocontrol. Sci. Technol.* **1998**, *8*, 23–32. [[CrossRef](#)]

- 
52. Kim, J.J.; Goettel, M.; Gillespie, D.R. Potential of *Lecanicillium* species for dual microbial control of aphids and the cucumber mildew fungus *Sphaerotheca fuliginea*. *Biol. Control* **2007**, *40*, 327–332. [[CrossRef](#)]
  53. Heintz, C.; Blaich, R. Ultrastructural and histochemical studies on interactions between *Vitis vinifera* L. and *Uncinula necator* (Schw.) Burr. *New Phytol.* **1990**, *115*, 107–117. [[CrossRef](#)]