

The Evaluation of Radiation Use Efficiency and Leaf Area Index Development for Estimation of Biomass Accumulation in Short Rotation Poplar and Annual Field Crops

Abhishek Mani Tripathi^{1*}, Eva Pohanková^{1,2}, Milan Fischer^{1,2}, Matěj Orság^{1,2}, Miroslav Trnka^{1,2}, Karel Klem¹, Michal V. Marek¹

¹ Global Change Research Institute CAS, Bělidla 986/4a, 603 00 Brno, Czech Republic; eva.pohankova@seznam.cz (E.P.); fischer.milan@gmail.com (M.F.); orsag.matej@gmail.com (M.O.); mirek_trnka@yahoo.com (M.T.); klem.k@czechglobe.cz (K.K.); marek.mv@czechglobe.cz (M.V.M.)

² Department of Agrosystems and Bioclimatology, Faculty of Agronomy, Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czech Republic

*Correspondence: manicfre@gmail.com or tripath.a@czechglobe.cz; Tel.: +420511192215

Figure S1: Developed an allometric relationship between aboveground dry mass (AGDM, kg stump⁻¹) and in diameter at breast height (DBH, mm) of poplar clone J-105 in coppice rotation after two years of plant growth.

