

Table S6. Protein motifs found in fungal effectoromes

Fungus	Reported effectorome*	RXLR	CRN	LysM	YFWxC	[LI]xAR	EAR	ToxA	Reference
<i>Venturia inaequalis</i>	1369	37	6	74	285	56	214	10	[79]
<i>Pestalotiopsis fici</i>	381	13	0	20	29	20	31	1	[80]
<i>Xylona heveae</i>	84	3	0	5	14	0	12	0	[80]
<i>Colletotrichum orbiculare</i>	615	28	2	46	82	50	76	6	[80]
<i>Mycosphaerella graminicola</i>	331	17	1	32	43	17	53	4	[80]
<i>Puccinia graminis f. sp. tritici</i>	1154	64	12	84	217	52	148	7	[80]
<i>Ustilago maydis</i>	212	5	0	13	18	9	48	1	[80]
<i>Fusarium verticillioides</i>	415	20	2	26	39	30	41	3	[80]
<i>Sclerotinia sclerotiorum</i>	329	9	1	16	25	16	52	0	[80]
<i>Leptosphaeria maculans</i>	1070	33	8	60	160	52	188	6	[81]
<i>Cochliobolus sativus</i>	196	7	0	18	20	14	20	0	[82]
<i>Parastagonospora nodorum</i>	246	15	0	17	37	20	27	2	[83]
<i>Mycosphaerella graminicola</i>	94	6	1	10	20	9	7	2	[55]

*the effectoromes were downloaded from the supplementary material of each report