

Article Risks in Major Cryptocurrency Markets: Modeling the Dual Long Memory Property and Structural Breaks

Zhuhua Jiang¹, Walid Mensi^{2,3} and Seong-Min Yoon^{4,*}

- ¹ Division of Chinese Foreign Affairs and Commerce, Hankuk University of Foreign Studies, Seoul 02450, Republic of Korea
- ² Department of Economics and Finance, College of Economics and Political Science, Sultan Qaboos University, Muscat 123, Oman
- ³ Department of Finance and Accounting, University of Tunis El Manar and IFGT, Tunis 248, Tunisia
- ⁴ Department of Economics, Pusan National University, Busan 46241, Republic of Korea
- * Correspondence: smyoon@pusan.ac.kr

Abstract: This study estimates the effects of the dual long memory property and structural breaks on the persistence level of six major cryptocurrency markets. We apply the Bai and Perron structural break test, Inclán and Tiao's iterated cumulative sum of squares (ICSS) algorithm, and the fractionally integrated generalized autoregressive conditional heteroscedasticity (FIGARCH) model, with different distributions. The results show that long memory and structural breaks characterize the conditional volatility of cryptocurrency markets, confirming our hypothesis that ignoring structural breaks leads to an underestimation of the persistence of volatility modeling. The ARFIMA-FIGARCH model, with structural breaks and a skewed Student–*t* distribution, fits the cryptocurrency market's price dynamics well.

Keywords: cryptocurrency; dual long memory (LM); structural breaks (SBs); efficient market hypothesis; ARFIMA-FIGARCH model



Citation: Jiang, Z.; Mensi, W.; Yoon, S.-M. Risks in Major Cryptocurrency Markets: Modeling the Dual Long Memory Property and Structural Breaks. *Sustainability* **2023**, *15*, 2193. https://doi.org/10.3390/su15032193

Academic Editor: Sajid Anwar

Received: 11 December 2022 Revised: 31 December 2022 Accepted: 19 January 2023 Published: 24 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

1. Introduction

If a certain financial market is not sufficiently efficient, prices do not accurately reflect information in that financial market, and the market function of resource allocation cannot be exercised. If a market does not develop in a more efficient way, it may not be sustainable in the long run.

The cryptocurrency market is relatively new and thus, has attracted much attention since its creation in 2008 [1]. The growing literature has addressed the stylized facts and efficiency hypothesis of this digital currency market. Urguhart [2] showed that the Bitcoin market has tended to be increasingly efficient since August 2013. Tiwari et al. [3] confirmed Urquhart's [2] findings, and Nadarajah and Chu [4] and Bariviera [5] revealed evidence of the efficiency of the Bitcoin market. Katsiampa [6] discovered that the AR (autoregressive)-CGARCH (component generalized autoregressive conditional heteroscedasticity) model is a suitable model to examine the volatility of the Bitcoin market. Takaishi [7] examined the multi-fractal properties of Bitcoin using intra-day data and the MF-DFA approach and found evidence of multi-fractality. Sensoy [8] found that the Bitcoin market is an efficient market in its weak-form version, adding that liquidity positively affects the information efficiency level, while volatility decreases this level. Vidal-Tomás and Ibañez [9] explored the semi-strong efficiency of Bitcoin. Using detrended fluctuation analysis and the Hurst exponent, Alvarez-Ramirez et al. [10] found evidence of asymmetric correlations in the Bitcoin market. Dyhrberg [11] used GARCH models to analyze the volatility of Bitcoin, gold, and the U.S. dollar and found that the models are crucial tools for portfolio risk management. Alvarez-Ramirez and Rodriguez [12] suggested that the cryptocurrency markets are informationally efficient during most of the sample period. Duan et al. [13]

showed that Bitcoin markets are close to an efficient market, although the efficiency varies over time. López-Martín et al. [14] revealed that the efficiency degree in six cryptocurrency markets increased over time. Kakinaka and Umeno [15] revealed that the cryptocurrency markets became more inefficient in the short-term, but not in the long-term during the COVID-19 pandemic. Phiri [16] tested random walk behavior in Bitcoin return dynamics in time-frequency space and determined that Bitcoin returns are the most predictable or the least weak-form market efficient. In the meantime, Aggarwal et al. [17], Lamoureux and Lastrapes [18], and Lastrapes [19] showed that not monitoring the structural breaks (SBs) when investigating a long memory (LM) process may result in a spurious estimation of the volatility persistence. In this regard, some studies on stock and commodity markets incorporated LM and SB together in their volatility modeling [20–23].

The above literature has ignored the modeling of dual LM property that may exist in both the conditional mean and conditional variance time series movements, along with SBs, which are major stylized facts in the volatility model of cryptocurrencies. LM and SBs have significant implications for asset allocations and portfolio management because they are related to market efficiency. In fact, the determination of the optimal investment strategies requires accurate modeling of the LM and SBs. The presence of LM rejects the market efficiency hypothesis. The evidence of the LM feature in volatility reveals volatility persistence, indicating that uncertainty is a key factor for a cryptocurrency market, and it should be managed. Thus, understanding the dual LM in cryptocurrency markets is essential for crypto investors and forecasters. In addition, the previous literature has focused only on Bitcoin, ignoring the rest of the major cryptocurrencies, such as Ethereum, which is the second-most important cryptocurrency in terms of market size. Modeling the dual LM and SBs in returns and volatility is hence an appealing topic of study on cryptocurrency markets.

This paper differs from, and adds to, the existing literature on cryptocurrency markets in at least two ways. First, this paper contributes to the related empirical studies by examining the presence of dual LM (in the mean and the variance) in six major digital currencies (Bitcoin, Dash, Ethereum, Litecoin, Monero, and Ripple). Second, we analyze the impacts of the SBs and LM on the variance persistence level. To the best of our knowledge, these two objectives have not yet been examined in cryptocurrency markets. Empirically, we consider ARFIMA-FIGARCH models with different residual distributions. These models have a particularity that allows them to assess the LM coefficients of the conditional mean and variance simultaneously. These models are widely employed in the empirical literature [24–33]. Third, not accounting for these two stylized facts can bring about sizeable upward biases in the degree of volatility persistence [34] because GARCH models do not consider SB variables. These variables have important implications regarding the expected changes and arbitrage activities, as well as on forecasting portfolio risk. The marginalization of the SBs and LM leads to the underestimation or overestimation of volatility for long stretches, leading to spurious portfolio risk assessments. To avoid the misestimation of the persistence and the distortion of information inflows, we apply Inclán and Tiao's [35] ICSS algorithm and identify multiple SBs for the markets to include receiving input of SBs in their GARCH volatility model.

The results provide evidence that the cryptocurrency markets are characterized by the presence of SBs and LM in their price dynamics, particularly during volatility. Moreover, the SB variables affect the conditional mean dynamics in the Ripple market and the conditional variances for all markets. In addition, the LM property characterizes the cryptocurrency markets. Fourth, the persistence in the variance coefficient values increases for all cryptocurrencies and differs in magnitude across the markets. Lastly, the ARFIMA-FIGARCH model, with SBs and a skewed Student–*t* distribution, is the best model, since it improves the accuracy of estimating the volatility of the cryptocurrency markets. The presence of LM and SBs in the variance model are crucial aspects for allocating assets and forecasting prices. The remainder of this paper is organized as follows. Section 2 discusses the methodology. Section 3 describes the data and conducts a preliminary analysis. Section 4 discusses the empirical results. Section 5 provides the concluding remarks.

2. Dual Long Memory Models with Structural Breaks

We assume that the returns for the cryptocurrency at time t, r_t can be described using the ARMA (m, n) model with SBs, which is written as follows:

$$r_t = \mu + \sum_{k=1}^m \psi_k r_{t-k} + \sum_{h=1}^n \theta_h \varepsilon_{t-h} + \sum_l \phi_l dum_{ml} + \varepsilon_t, \tag{1}$$

$$\varepsilon_t = z_t \sigma_t,$$
 (2)

where dum_m is the dummy variable to consider the possible sudden changes in the conditional mean process, which are captured by the Bai and Perron [36] test; ε_t is the independently distributed error, with zero mean and variance σ_t^2 , and z_t is random innovation.

This model can be extended to the ARFIMA-FIGARCH model, which allows for a dual LM feature in conditional mean and conditional variance.

The ARFIMA (m, ξ, n) model can be written as follows:

$$\psi(L)(1-L)^{\xi}\left(r_t - \sum_l \phi_l dum_{m_l}\right) = \theta(L)\varepsilon_t,\tag{3}$$

where ξ is the LM parameter in the above conditional mean equation; *L* denotes the lag or backshift operator; and $\psi(L) = 1 - \psi_1 L - \psi_2 L^2 \dots - \psi_m L^m$ and $\theta(L) = 1 + \theta_1 L + \theta_2 L^2 \dots + \theta_n L^n$ are the autoregressive (AR) and the moving-average (MA) polynomials, respectively.

The ARFIMA process is non-stationary when $\xi \leq -0.5$ or $\xi \geq 0.5$. The process exhibits anti-persistency for $-0.5 < \xi < 0$ and LM for $0 < \xi < 0.5$. The process shows short memory for $\xi = 0$.

The FIGARCH (p, d, q) model introduced by Baillie et al. [37] can be represented as follows:

$$\alpha(L)(1-L)^{d}\varepsilon_{t}^{2} = \omega + [1-\beta(L)]v_{t} + \sum_{i}\delta_{i}dum_{v_{i}},$$
(4)

where $v_t = \varepsilon_t^2 - \sigma_t^2$, $\alpha(L) = \alpha_1 L + ... + \alpha_q L^q$ and $\beta(L) = \beta_1 L + ... + \beta_p L^p$, and $(1 - L)^d$ denotes the fractional differencing operator. dum_v is the dummy variable that considers possible SBs in the conditional variance process, which can be captured by the ICSS algorithm of Inclán and Tiao [35]. We have a stationary LM process when 0 < d < 1. The FIGARCH (p, d, q) process exhibits anti-persistency in the volatility for 0 < d < 0.5 and LM in the volatility for 0.5 < d < 1. The process shows short memory in the volatility for d = 0.5.

Chung [38] suggested a slightly different process to include LM features in the conditional variance as:

$$\sigma_t^2 = \sigma^2 + \left\{ 1 - [1 - \beta(L)]^{-1} \varnothing(L) (1 - L)^d \right\} \left(\varepsilon_t^2 - \sigma^2 \right), \tag{5}$$

where σ^2 is the unconditional variance of ε_t .

The idea of fractional integration has been applied to other types of GARCH models, including the fractionally integrated exponential GARCH (FIEGARCH) of Bollerslev and Mikkelsen [39] and the fractionally integrated asymmetric power ARCH (FIAPARCH) of Tse [40].

Similar to the GARCH (p,q) process, the FIEGARCH (p,d,q) is specified as follows:

$$\log\left(\sigma_t^2\right) = \omega + \varnothing(L)^{-1}(1-L)^{-d}[1+\alpha(L)]g(z_{t-1}),\tag{6}$$

where function $g(z_t) = \theta_1 z_t + \theta_2(|z_t| - E|z_t|)$ [41].

In the model, the parameters θ_1 and θ_2 depict the sign effect and the magnitude effect, respectively. More specifically, good news has $(\theta_1 + \theta_2)$ impact on volatility, while bad

news has a $(\theta_1 - \theta_2)$ impact on volatility. For $\theta_1 > 0$ and $\theta_2 > 0$, positive return shocks will have a greater influence on volatility than negative return shocks; for $\theta_1 < 0$ and $\theta_2 > 0$, negative return shocks cause greater volatility changes than positive return shocks. As for the FIGARCH specification, the FIEGARCH model nests the conventional EGARCH model for d = 0, and the IEGARCH model for d = 1.

The FIAPARCH (p, d, q) model is specified as follows:

$$\sigma_t^{\delta} = \omega + \left\{ 1 - \left[1 - \beta(L) \right]^{-1} \varnothing(L) (1 - L)^d \right\} (|\varepsilon_t| - \gamma \varepsilon_t)^{\delta}, \tag{7}$$

where $\delta > 0$ and $-1 < \gamma < 1$. When $\gamma > 0$, negative shocks have more impact on volatility than positive shocks, and the inverse is also true. The conditional variance has the LM property if 0 < d < 1. The FIAPARCH model also nests the FIGARCH model when $\delta = 2$ and $\gamma = 0$. Thus, the FIAPARCH model is superior to the FIGARCH model, since the former model allows for asymmetric LM features in the conditional variance.

Davidson [42] suggested the hyperbolic GARCH (HYGARCH) model, which is an extended model of the FIGARCH process, with hyperbolic convergence rates. The HY-GARCH (p, d, q) model is represented as follows:

$$\sigma_t^2 = \omega [1 - \beta(L)]^{-1} + \left\{ 1 - [1 - \beta(L)]^{-1} \varnothing(L) (1 + k) \left[(1 - L)^d - 1 \right] \right\} \varepsilon_t^2, \tag{8}$$

where $k \ge 0$ and $d \ge 0$. HYGARCH nests the FIGARCH model with k = 1 and the GARCH model with k = 0. The process is stationary for 0 < k < 1, but nonstationary for k > 1.

 z_t is an innovation process. A common choice for the density of this process is an independent and identical normal distribution, $z_t \sim N(0, 1)$. The log-likelihood function for the normal innovation is expressed as follows:

$$L_{norm} = -\frac{1}{2} \sum_{t=1}^{T} \left[ln(2\pi) + ln(h_t) + z_t^2 \right].$$
(9)

However, we assume that z_t follows the skewed Student-*t* distribution to catch the skewness and kurtosis features in a return series process. The log-likelihood function for the skewed Student-*t* innovation is expressed as follows:

$$L_{norm} = ln \left[\Gamma\left(\frac{\nu+1}{2}\right) \right] - ln \left[\Gamma\left(\frac{\nu}{2}\right) \right] - \frac{1}{2} ln [\pi(\nu-2)] + ln \left(\frac{2}{\lambda+1/\lambda}\right) + ln(s) - \frac{1}{2} \sum_{t=1}^{T} \left[ln(h_t) + (1+\nu) ln \left(1 + \frac{sz_t + m}{\nu - 2} \lambda^{-l_t}\right) \right],$$
(10)

where $I_t = 1$ if $z_t \ge -\frac{m}{s}$ and $I_t = -1$ if $z_t < -\frac{m}{s}$.

The constants $m = m(\lambda, \nu)$ and $s = \sqrt{s^2(\lambda, \nu)}$ are the mean and standard deviations of the skewed Student-*t* distribution, respectively, and are as follows:

$$m(\lambda,\nu) = \frac{\Gamma\left(\frac{\nu-1}{2}\right)\sqrt{\nu-2}}{\sqrt{\pi}\Gamma\left(\frac{\nu}{2}\right)} \left(\lambda - \frac{1}{\lambda}\right), \ s^2(\lambda,\nu) = \left(\lambda^2 + \frac{1}{\lambda^2} - 1\right) - m^2, \tag{11}$$

where λ and ν identify the skewness and kurtosis of this innovation distribution, respectively. The symmetry of the distribution increases as $\lambda \to 0$, and the thickness of the tails decreases as $\nu \to \infty$.

3. Data and Preliminary Statistics

For the empirical analysis, we collected daily closing price data for six popular cryptocurrencies: Bitcoin (BTC), Dash (DASH), Ethereum (ETH), Litecoin (LTC), Monero (XMR), and Ripple (XRP). The sample period spans from 29 April 2013–29 October 2020 for BTC, 14 February 2014–29 October 2020 for Dash (DASH), 9 August 2015–29 October 2020 for ETH, 29 April 2013–29 October 2020 for LTC, 21 May 2014–29 October 2020 for XMR, and 4 August 2013–29 October 2020 for XRP. The data were sourced from Cryptocurrency Market Capitalizations (https://coinmarketcap.com, accessed on 30 October 2020). Figure 1 displays the logarithmic price dynamics of each cryptocurrency, which show significant upwards movements, particularly in 2016 and 2017. The return dynamics of each cryptocurrency market are plotted in Figure 2, showing evidence of volatility clustering and fat tails. Figure 2 plots the presence of at least three SB events in the return series for all markets.



Figure 1. Dynamics of the logarithmic prices of cryptocurrencies.



Figure 2. Dynamics of the returns of cryptocurrencies. The dotted lines show the ± 3 standard deviation bands. The jumps in these lines imply the SB points measured by the ICSS algorithm.

Table 1 shows that the average returns for all cryptocurrencies are positive. Dash is the most volatile market, while Bitcoin is the least volatile. All return series exhibit asymmetry and fat tails. The Jarque–Bera test results reject the null hypothesis of normal distributions for all return series, which are stationary according to the conventional unit root tests results. Serial correlations of the squared returns and ARCH effects have been found in all cryptocurrencies, implying that the GARCH-class models could fit our return series well.

	BTC	DASH	ETH	LTC	XMR	XRP		
	A. Descriptive statistics							
Mean	0.1635	0.2117	0.2584	0.0921	0.1848	0.1406		
Maximum	35.7451	127.0565	41.2337	82.8968	58.4637	102.7356		
Minimum	-46.4729	-46.7565	-130.2105	-51.3925	-49.4208	-61.6273		
Std. Dev	4.2497	7.4348	6.8929	6.3026	6.8015	6.9866		
Kurtosis	12.2749	45.1602	71.8075	25.9537	7.8084	31.6827		
Skewness	-0.5889	2.8712	-3.4718	1.5024	0.4495	2.0074		
Jarque–Bera	17,366 ***	211,559 ***	414,411 ***	77,961.5 ***	6059.6 ***	112,360 ***		
B. Results of the unit root tests for the logarithmic prices								
ADF	-1.12	-2.21	-2.05	-1.29	-0.70	-1.45		
PP	-0.81	-2.22	-1.45	-1.11	-0.70	-1.13		
KPSS	0.47 ***	0.92 ***	1.09 ***	0.55 ***	0.87 ***	0.57 ***		
	(C. Results of the	e unit root tests	for the returns				
ADF	-10.63 ***	-49.67 ***	-7.69 ***	-17.24 ***	-12.79 ***	-14.19 ***		
PP	-53.14 ***	-49.71 ***	-42.30 ***	-51.88 ***	-49.07 ***	-49.45 ***		
KPSS	0.08	0.34	0.26	0.09	0.16	0.08		
D. Statistical properties								
Q(12)	34.59 ***	31.9 ***	28.3 *	36.4 ***	30.5 **	62.9 ***		
$Q_{S}(12)$	304.7 ***	293.7 ***	126.0 ***	488.4 ***	350.5 ***	391.3 ***		
ARCH(12)	16.3 ***	31.8 ***	9.5 ***	32.3 ***	16.1 ***	24.1 ***		

Table 1. Descriptive statistics, results of the unit root tests, and statistical properties.

Notes: ADF, PP, and KPSS are the augmented Dickey–Fuller unit root test statistic, the Phillips–Perron unit root test statistic, and the Kwiatkowski–Phillips–Schmidt–Shin stationarity test statistic, respectively. Q(20) and $Q_S(20)$ denote the Box–Pierce Q tests, with 20 lags for no serial correlation of returns and squared returns, respectively. ARCH (10) denotes the ARCH LM test, with 10 lags for homoscedasticity. ***, **, and * denote the rejection of the null hypothesis at the 1%, 5%, and 10% significance levels, respectively.

4. Empirical Results

4.1. Estimation Results, with and without Structural Breaks

We have estimated various GARCH-type models using the OxMetrics ver. 8 and determined that the ARFIMA (m, ξ, n) -FIGARCH (p, d, q) model is the best model, according to the information criteria. The estimation results of the ARFIMA (m, ξ, n) -FIGARCH (p, d, q)model reported in Table 2 show significant evidence of the anti-persistency in the conditional mean in some cryptocurrency markets (DASH, LTC, XMR, and XRP). Regarding the variance equation, we found significant evidence of LM in the conditional variance in all cryptocurrency markets, and that LTC is the market with the highest persistence, followed by BTC and XMR. For the remaining markets, the persistence level ranged from 0.5227 to 0.5345. Overall, the results reveal that the cryptocurrency markets exhibit LM behavior in their variances, and a lack of LM features in their return dynamics. These results indicate a non-linear dependence on the conditional variance and also disconfirm the weak-form efficiency hypothesis, which has implications in terms of asset allocation and efficiency levels. The short-term ARCH coefficient is statistically significant for the series, except for DASH and ETH. The long-term GARCH coefficient is statistically significant at the 10% level in all cryptocurrency markets. The diagnostic test results (see Panel C) demonstrate that there is no evidence of misspecification.

To avoid the underestimation of the persistence, we consider the SB variables in both the mean and the variance models. We employ the Bai and Perron [36] test to capture the SBs for the conditional mean dynamics and the ICSS algorithm of Inclán and Tiao [35] to identify the SBs for the conditional variance dynamics. This ICSS model is used to identify multiple sudden jumps in the unconditional volatility in a series of independent observations. For detailed information on these tests, see Bai and Peron [36] and Inclán and Tiao [35].

	BTC	DASH	ETH	LTC	XMR	XRP	
		A. Con	ditional mean e	quation			
	0.1039 **	0.0238	0.1868 *	-0.0334	0.1432 **	-0.1408 ***	
μ	(0.0447)	(0.0705)	(0.0977)	(0.0357)	(0.0713)	(0.0419)	
z	-0.0048	-0.0275 *	-0.0116	-0.0607 ***	-0.0492 ***	-0.0505 ***	
ζ	(0.0150)	(0.0144)	(0.0154)	(0.0125)	(0.0145)	(0.0134)	
		B. Condi	tional variation	equation			
	102.83 *	330.11 ***	232.60 ***	117.80 *	683.23 ***	48.932 ***	
ω	(53.063)	(13.694)	(14.771)	(60.299)	(24.150)	(14.858)	
J	0.6978 ***	0.5345 ***	0.5052 ***	0.7212 ***	0.6344 ***	0.5227 ***	
и	(0.0650)	(0.0396)	(0.0465)	(0.0679)	(0.0314)	(0.0560)	
A 1	0.2517 ***	0.1813	0.0893	0.2950 ***	0.1679 **	0.1738 ***	
<i>u</i> ₁	(0.0524)	(0.2120)	(0.1297)	(0.0567)	(0.0775)	(0.0667)	
R.	0.7364 ***	0.4101 *	0.3654 ***	0.7919 ***	0.4635 ***	0.6029 ***	
ρ_1	(0.0607)	(0.2478)	(0.1324)	(0.0518)	(0.0843)	(0.0666)	
Acummetru	-0.0380*	380 * 0.0781 *** 0	0.0646 **	0.0274	0.0581 **	0.0562 **	
215ymmetry	(0.0205)	(0.0237)	(0.0258)	(0.0208)	(0.0292)	(0.0269)	
Tail	3.2116 ***	3.2621 ***	3.0608 ***	2.9527 ***	3.6222 ***	3.5099 ***	
1411	(0.1170)	(0.1582)	(0.1646)	(0.0819)	(0.2208)	(0.2094)	
C. Diagnostic checking							
logL	-7018.98	-7463.62	-5783.06	-7785.53	-5868.94	-7374.36	
BĪC	5.1446	6.1182	6.0872	5.7060	6.1739	6.2944	
Q(20)	58.64 ***	51.87 ***	4.49 **	54.95 ***	56.44 ***	53.34 ***	
$Q_{S}(20)$	5.97	5.81	0.36	2.62	17.38	16.89	
ARCH(10)	0.34	0.34	1.19	0.09	1.20	1.17	

Table 2. Estimation results of the ARFIMA-FIGARCH model.

Notes: Q(20) and $Q_5(20)$ denote the Box–Pierce Q tests, with 20 lags for no serial correlation of returns and squared returns, respectively. ARCH (10) denotes the ARCH test, with 10 lags for homoscedasticity. See also the note for Table 1. *, ** and *** denote the rejection of the null hypothesis at the 1%, 5%, and 10% significance levels, respectively.

The results of the SB identification are reported in Tables 3 and 4. As shown in Table 3, we identify evidence of SBs in the conditional mean equation only for the DASH cryptocurrency market dynamics, while we determine strong evidence of SBs in the conditional variances for all cryptocurrencies, as shown Table 4. These results confirm that economic and financial shocks affect the volatility of cryptocurrencies. The dates for each SB of each market are reported in Table 4. These SB variables are explanatory variables and are included in both the mean and variance equations of the GARCH models for most accuracy estimations.

Table 3. Structural breaks in the conditional mean detected by the Bai and Perron test.

Cryptocurrency	Conditional Mean	Time Period
BTC	0.1654	30 April 2013~29 October 2020
DASH	0.5925 - 0.3010	15 February 2014~20 December 2017 21 December 2017~29 October 2020
ETH	0.2584	10 August 2015~29 October 2020
LTC	0.0922	30 April 2013~29 October 2020
XMR	0.1849	22 May 2014~29 October 2020
XRP	0.1406	5 August 2013~29 October 2020

Note: The SBs for the conditional mean are identified using the Bai and Perron [36] test.

Table 5 shows that the SBs insignificantly affect the conditional mean of the DASH market. In addition, we find evidence of the anti-persistency in the mean for four out of the six markets (DASH, LTC, XMR and XRP), which is similar to the results shown in

Table 2. Interestingly, we find that the LM coefficient (d) is significant for all cases, and it increases significantly for four out of the six markets, with the exception of BTC and LTC, indicating higher predictability. For example, by comparing Tables 2 and 5, we find that DASH's LM coefficient increases from 0.5345 to 0.6207 and that of ETH increases from 0.5052 to 0.5970. These results confirm our hypothesis that ignoring SB variables causes an underestimation of the persistence of the variance modeling. These results indicate that the cryptocurrency markets respond progressively to information flows. Past prices can be used to predict future prices in these cryptocurrency markets, disconfirming the random walk hypothesis and signaling an inefficient market. A plausible explanation for these results can be attributed to the relatively small size of the total cryptocurrency market compared to classical financial markets. Cryptocurrency markets have fewer regulations and deserve financial reforms. In addition, cryptocurrency prices are not backed by commodities, as is the case with stock prices [43].

Cryptocurrency	Standard Deviation	Time Period
BTC	6.4391	30 April 2013~14 April 2014
	3.1868	16 April 2014~5 May 2017
	5.5929	6 May 2017~24 April 2018
	3.6916	25 April 2018~29 October 2020
DASH	17.6279	15 February 2014~19 August 2014
	5.3276	20 August 2014~31 December 2016
	7.8378	1 January 2017~12 February 2018
	5.2387	13 February 2018~29 October 2020
ETH	8.5226	10 August 2015~7 February 2018
	4.9148	8 February 2018~29 October 2020
LTC	5.3273	30 April 2013~15 November 2013
	14.0939	16 November 2013~13 April 2014
	5.5548	14 April 2014~29 October 2020
XMR	15.3774	22 May 2014~23 July 2014
	7.2966	24 July 2014~12 March 2018
	4.8902	13 March 2018~29 October 2020
XRP	10.9748	5 August 2013~28 May 2014
	4.4208	29 May 2014~20 March 2017
	12.5448	21 March 2017~12 February 2018
	4.6843	13 February 2018~29 October 2020

Table 4. Structural breaks in volatility detected by the ICSS algorithm.

Note: The structural break points for the conditional variance are identified using the ICSS algorithm of Inclán and Tiao [35].

Note that the misspecification of the volatility model affects the asset allocation and portfolio risk design. The ARCH effect decreased significantly after accounting for the SB variables, while the coefficient of the GARCH effect increased with the presence of the SB variables. Again, this result shows the importance of these variables when modeling the persistence of cryptocurrencies.

More importantly, we find a significant tail for all markets, indicating that the ARFIMA $(0, \xi, 0)$ -FIGARCH (1, d, 1) model, with SBs and a skewed Student–*t* distribution, is the better model because it outperforms the ARFIMA $(0, \xi, 0)$ -FIGARCH (1, d, 1) model without SB variance and with a normal distribution. This result is confirmed by the Bayesian information criterion (BIC) and log likelihood values, suggesting that considering the LM properties and SBs can enhance the estimation of the conditional mean and conditional variance. The estimation results of the ARFIMA-HYGARCH and ARFIMA-FIAPARCH models, with structural breaks and a skewed Student–*t* distribution, are summarized in Tables A1 and A2.

	BTC	DASH	ETH	LTC	XMR	XRP
		A. Cond	ditional mean e	equation		
11	0.0937 **	0.0938	0.2136 **	-0.0319	0.1454 **	-0.1652 ***
μ	(0.0440)	(0.0929)	(0.0990)	(0.0349)	(0.0713)	(0.0573)
dum m	-	-0.1526	_	_	-	-
uum_m		(0.1194)				
۶	-0.0077	-0.0324 **	-0.0104	-0.0607 ***	-0.0494 ***	-0.0256 *
5	(0.0143)	(0.0149)	(0.0157)	(0.0125)	(0.0145)	(0.0132)
		B. Condit	tional variation	equation		
	132.34 ***	933.77 ***	441.10 ***	62.953	830.21 ***	735.43 *
ω	(14.012)	(140.81)	(17.116)	(38.906)	(134.47)	(423.29)
dum o	-42.464 ***	-285.41 ***	123.17 ***	-11.456	-41.883	-78.947
uum_0_1	(12.923)	(47.521)	(21.982)	(20.308)	(128.53)	(84.269)
dum 72-	135.64 ***	-29.852	_	-82.862 ***	2.5130	572.12
uum_0 <u>2</u>	(11.104)	(134.33)	-	(25.994)	(28.504)	(413.70)
dum vo	126.63 ***	87.502	_	_	_	664.26
<i>uum_0</i> 3	(12.202)	(142.55)				(465.98)
d	0.6591 ***	0.6207 ***	0.5970 ***	0.4587 ***	0.6536 ***	0.6589 ***
и	(0.0486)	(0.0409)	(0.0366)	(0.0648)	(0.0344)	(0.0619)
Na	0.2626 ***	0.1805	0.0914	0.3868 ***	0.1689 **	0.2684 ***
ul	(0.0638)	(0.1843)	(0.1093)	(0.0876)	(0.0691)	(0.0909)
Ba	0.6919 ***	0.4708 **	0.4469 ***	0.6459 ***	0.6138 ***	0.4536 ***
ρ_1	(0.0710)	(0.2029)	(0.1023)	(0.0797)	(0.0654)	(0.1001)
Asummetru	-0.0434 **	0.0771 ***	0.0759 ***	0.0305	0.0577 **	0.0202
215ymmetry	(0.0211)	(0.0247)	(0.0264)	(0.0210)	(0.0270)	(0.0244)
Tail	3.0250 ***	3.1176 ***	3.0441 ***	3.0322 ***	3.4819 ***	2.9105 ***
1411	(0.1074)	(0.1536)	(0.1616)	(0.1025)	(0.2061)	(0.1072)
		С. Г	Diagnostic chec	king		
logL	-7001.55	-7451.21	-5766.36	-7777.29	-7373.98	-7588.03
BĪC	5.1424	6.1233	6.0737	5.7057	6.3007	5.7748
Q(20)	58.91 ***	43.92 ***	50.22 ***	56.02 ***	53.31 ***	38.78 ***
$Q_{S}(20)$	6.28	7.58	15.99	2.89	16.60	2.66
ARCH(10)	0.39	0.47	1.19	0.06	0.08	0.15

Table 5. Estimation results of the ARFIMA-FIGARCH model, with structural breaks and a skewed Student–t distribution.

Notes: See the notes of Table 2. *dum_m* in the conditional mean equation is the dummy variable that considers the SB for Period_1 in Table 3. Similarly, *dum_v*₁, *dum_v*₂, *and dum_v*₃ in the conditional variance equation are the dummy variables that consider the SBs for Period_1, Period_2, and Period_3 in Table 4, respectively. The dummy variable for Period_0 is not included in the conditional mean and variance equations to avoid the multi-collinearity problem. *, ** and *** denote the rejection of the null hypothesis at the 1%, 5%, and 10% significance levels, respectively.

4.2. Evaluation of Forecasting Accuracy and Diebold and Mariano (DM) Test

We checked whether the inclusion of SBs in the mean and conditional variance model can enhance the forecasting accuracy of the model. To quantify forecasting accuracy, we measured the mean of absolute errors (MAE) and the mean squared errors (MSE), as follows:

$$MAE = \frac{1}{T} \sum_{i=1}^{T} \left| \sigma_{f,t}^2 - \sigma_{a,t}^2 \right|,$$
 (12)

$$MSE = \frac{1}{T} \sum_{i=1}^{T} \left(\sigma_{f,t}^2 - \sigma_{a,t}^2 \right)^2,$$
(13)

where *T* denotes the number of forecasting data points and $\sigma_{f,t}^2$ is the volatility forecast for day *t*, whereas $\sigma_{a,t}^2$ denotes actual volatility on day *t*. In accordance with the relevant literature [44–46], daily ex post actual volatility was calculated by the squared returns as follows:

$$\sigma_t^2 = r_t^2 \tag{14}$$

We also performed the DM test (Diebold and Mariano, 2002) to compare the 1-dayahead predictive accuracy between each of considered models and the benchmark model (the standard GARCH model, in our analysis). Specifically, the DM test identifies whether the two forecasts are equally good. A rejection of the null hypothesis determines that there is a significant difference between the forecasts. Each panel of Table 6 displays the results of the DM test based on the MSE and MAE loss functions. A positive value for the DM test statistics means that the forecasting results from the compared model are more accurate

	BTC	DASH	ETH	LTC	XMR	XRP	
(A) Standard GARCH model without structural break dummies							
MAE	15.2040	28.5379	29.4993	49.2557	21.5686	27.7451	
MSE	497.11	2702.96	1484.65	4433.68	1018.60	1464.10	
(B) ARFIMA-FIGARCH model without structural break dummies							
MAE	8.7251	22.4812	22.8393	21.8084	19.2678	13.0244	
DM test	8.36 ***	7.24 ***	6.15 ***	8.43 ***	5.64 ***	6.98 ***	
MSE	317.93	2391.58	1135.21	2035.86	950.62	440.12	
DM test	4.51 ***	3.03 ***	2.26 ***	3.73 ***	2.39 ***	3.36 ***	
	(C) ARFIMA-FIGARCH model with structural break dummies						
MAE	9.6617	23.9853	23.3647	21.2344	19.4171	14.7833	
DM test	8.06 ***	6.29 ***	6.38 ***	8.36 ***	5.56 ***	7.36 ***	
MSE	323.72	2449.47	1158.05	2003.24	954.80	494.91	
DM test	4.94 ***	2.89 ***	2.40 ***	3.70 ***	2.40 ***	3.62 ***	

Table 6. Forecast evaluation and DM test using the recent 100 one-step-ahead forecasts.

that those from the benchmark model.

Notes: This table is created from the out-of-sample results of the one-step-ahead forecasting for robustness checks. See also the note of Table 1. *** denote the rejection of the null hypothesis at the 10% significance levels.

We evaluated the forecasting accuracy of the recent 100 one-step-ahead forecasts generated from the ARFIMA-FIGARCH model, with and without SBs. The results are reported in Table 6. Smaller forecasting error statistics imply the superior forecasting ability of a given model. An overall evaluation demonstrates that all the dual LM models with SB dummies provide relatively good forecasts of the volatility of six cryptocurrency markets, whereas those models which do not consider LM features and do not include SB dummies seem to be a poor alternative. Thus, the results of one-step-ahead forecasting analysis indicate that the volatility models with SBs and dual LM parameters produce excellent out-of-sample predictability.

5. Conclusions

This paper is the first to estimate the effects of the dual LM features in the mean and variance and SBs on the persistence levels of the Bitcoin (BTC), Dash (DASH), Ethereum (ETH), Litecoin (LTC), Monero (XMR), and Ripple (XRP) cryptocurrency markets. These are interesting findings that can improve investors' decision making. First, the cryptocurrency markets exhibit SBs and LM features in their price dynamics (particularly in volatility). Litecoin is the market with the highest persistence, followed by Bitcoin and Monero. Second, SBs affect the conditional mean dynamics in the Dash market. Third, evidence of LM in the mean dynamics is found in the Dash, Litecoin, Monero, and Ripple cryptocurrency markets after accounting for the SBs. Fourth, there are persistent increases in the variance coefficient values for all cryptocurrencies and different magnitudes across the markets. Fifth, the cryptocurrency markets exhibit LM behavior in their variances and a lack of LM in their returns. The presence of LM reflects the inefficiency of the cryptocurrency markets. Lastly, we show that the ARFIMA $(0, \xi, 0)$ -FIGARCH (1, d, 1) model, with SBs and a skewed Student -t distribution, is a very good model since it improves the accuracy of the estimations of the volatility of the cryptocurrency markets. The presence of LM in the variance shows that consideration of the LM feature is important in measuring risk

(uncertainty), and that it is a key factor in the dynamics of cryptocurrency prices. The obtained results bolster the existing literature [17–19]. These results can assist investors and portfolio managers in terms of their decision-making processes. In practice, ignoring the SBs and LM features in the volatility model leads to spurious results for the persistence level and incorrect inferences regarding the efficiency hypothesis, thus affecting asset allocation, price prediction, and portfolio risk. The cryptocurrency market is widely affected by information flow. Investors should be cautious in the presence of SBs that affect the price dynamics and market volatility. As an example, favorable news increases cryptocurrency prices, which then induce speculation in these markets. The price dynamics of the cryptocurrency market are highly volatile, and market conditions change frequently. Therefore, it is necessary to continuously search for the most suitable model that can explain cryptocurrency market fluctuations.

Author Contributions: All the authors contributed to the entire process of writing this paper. Conceptualization, Z.J., W.M. and S.-M.Y.; data curation, W.M.; methodology, Z.J., W.M. and S.-M.Y.; formal analysis, Z.J. and W.M.; funding acquisition, Z.J. and S.-M.Y.; investigation, Z.J. and W.M.; project administration, Z.J. and S.-M.Y.; software, Z.J.; supervision, S.-M.Y.; validation, Z.J. and W.M.; visualization, Z.J. and S.-M.Y.; writing—original draft, Z.J., W.M. and S.-M.Y.; writing—review and editing, Z.J. and S.-M.Y. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2020S1A5B8103268).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The sample data are sourced from Cryptocurrency Market Capitalizations (https://coinmarketcap.com) (accessed on 2 February 2022).

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Estimation results of the ARFIMA (m, ξ , n)-HYGARCH (p, d, q) model, with structural breaks and a skewed Student-t distribution.

	BTC	DASH	ETH	LTC	XMR	XRP		
A. Conditional mean equation								
	0.0730	0.1010	0.1961 **	-0.0295	0.1273 *	-0.1678 ***		
μ	(0.0455)	(0.0958)	(0.0991)	(0.0353)	(0.0722)	(0.0546)		
dum m		-0.1695						
uum_m	-	(0.1225)	-	-	-	-		
7	-0.0102	-0.0318 **	-0.0113	-0.0644 ***	-0.0466 ***	-0.0277 **		
5	(0.0138)	(0.0149)	(0.0155)	(0.0118)	(0.0146)	(0.0133)		
		B. Condit	tional variation	ı equation				
	2.6424 *	18.155	3.5248	3.4735	60.753 ***	14.127		
ω	(1.3817)	(14.606)	(2.5354)	(2.2756)	(22.070)	(15.236)		
	-2.4547 *	-15.358	-0.3786	-1.0387	-56.585 ***	-14.353		
uum_{01}	(1.3651)	(13.832)	(1.9222)	(5.5905)	(21.776)	(15.269)		
dum n-	5.2440 *	-8.6414		-3.5754	-58.460 ***	-3.1874		
uum_{02}	(2.8119)	(11.425)	-	(2.3052)	(21.881)	(10.132)		
dum no	-2.1341 *	-14.767	_	_	_	-11.901		
uum_03	(1.2693)	(13.432)				(13.249)		
d	0.6940 ***	0.6257 ***	0.5933 ***	0.6212 ***	0.6189 ***	0.4328 ***		
и	(0.1562)	(0.1988)	(0.1489)	(0.0944)	(0.1247)	(0.0592)		
N 1	0.3453 ***	0.2790 *	0.1017	0.5388 ***	0.1935 **	0.3413		
μŢ	(0.1054)	(0.1649)	(0.1316)	(0.1285)	(0.0816)	(0.6512)		
B1	0.7659 ***	0.5465 **	0.4982 ***	0.8345 ***	0.5934 ***	0.3000		
P1	(0.0819)	(0.2563)	(0.1380)	(0.0504)	(0.0810)	(0.6510)		

	BTC	DASH	ETH	LTC	XMR	XRP
B. Conditional variation equation						
A	-0.0501 **	0.0760 ***	0.0693 ***	0.0271	0.0469 *	0.0156
Asymmetry	(0.0220)	(0.0245)	(0.0269)	(0.0205)	(0.0269)	(0.0239)
Tail	2.4971 ***	3.1631 ***	2.7898 ***	2.2380 ***	3.6152 ***	2.4985 ***
1011	(0.1345)	(0.2170)	(0.2124)	(0.0947)	(0.2791)	(0.1588)
$Log \; \alpha \; (HY)$	0.2053 *	-0.0189	0.1744	0.5508 **	-0.0308	0.6386 **
	(0.1222)	(0.0772)	(0.1322)	(0.2627)	(0.0564)	(0.2563)
C. Diagnostic checking						
logL	-6983.22	-7452.73	-5767.67	-7762.52	-7374.35	-7579.83
BĬC	5.1319	6.1278	6.0790	5.6979	6.3043	5.7716
Q(20)	63.43 ***	42.32 ***	52.16 ***	51.13 ***	52.32 ***	35.07 ***
$Q_{S}(20)$	6.35	5.80	11.58	4.37	16.60	1.14
ARCH(10)	0.43	0.35	1.01	0.07	1.21	0.06

Table A1. Cont.

Notes: See the notes for Table 2. *dum_m* in the conditional mean equation is the dummy variable that considers the structural break for Period_1 in Table 3. Similarly, *dum_v*₁, *dum_v*₂, and *dum_v*₃ in the conditional variance equation are the dummy variables that consider the SBs for Period_1, Period_2, and Period_3 in Table 4, respectively. The dummy variable for Period_0 is not included in the conditional mean and variance equations to avoid the multi-collinearity problem. *, ** and *** denote the rejection of the null hypothesis at the 1%, 5%, and 10% significance levels, respectively.

Table A2. Estimation results of the ARFIMA (m, ξ , n)-FIAPARCH (p, d, q) model, with structural breaks and a skewed Student–t distribution.

	BTC	DASH	ETH	LTC	XMR	XRP
		A. Cond	litional mean e	equation		
μ	0.0987 **	0.0853	0.2253 **	-0.0351	0.1878 **	-0.1255 **
	(0.0427)	(0.0892)	(0.0990)	(0.0354)	(0.0766)	(0.0530)
darran ana		-0.1502				
uum_m	-	(0.1146)	-	-	-	-
7	-0.0127	-0.0373 ***	-0.0116	-0.0636 ***	-0.0478 ***	-0.0269 *
ç	(0.0153)	(0.0139)	(0.0158)	(0.0122)	(0.0142)	(0.0155)
		B. Condit	ional variation	equation		
	127.58 ***	95.258 ***	248.27	167.92 ***	304.58 ***	86.419 ***
W	(13.697)	(35.191)	(673.67)	(16.933)	(103.76)	(26.356)
dum 71.	-42.149 **	-14.238	67.692	-24.909	-19.594	-2.1122
<i>uum_0</i> 1	(20.290)	(10.453)	(247.51)	(18.485)	(135.00)	(6.8424)
dum no	128.42 ***	16.185	_	-267.27 ***	8.0732	49.194 **
uum_02	(8.5360)	(28.737)		(27.069)	(218.50)	(22.530)
dum v2	119.02 ***	23.391	-	_	-	54.502 **
<i>uum_</i> 03	(11.214)	(33.446)				(22.185)
d	0.6631 ***	0.5065 ***	0.5621 **	0.4116 ***	0.6007 ***	0.5227 ***
	(0.0492)	(0.0548)	(0.2224)	(0.0684)	(0.1003)	(0.0335)
<i>α</i> ₁	0.2668 ***	0.1265	0.1049	0.4331 ***	0.1930 ***	0.2881 *
n1	(0.0697)	(0.1491)	(0.1082)	(0.1408)	(0.0749)	(0.1610)
B1	0.7037 ***	0.3715 **	0.4453 ***	0.6113 ***	0.6103 ***	0.3973 **
P1	(0.0737)	(0.1734)	(0.1643)	(0.1412)	(0.0604)	(0.1883)
$APARCH(\gamma)$	-0.0741	0.0253	-0.0580	-0.1761 ***	-0.2238 ***	-0.1120 *
(1)	(0.0570)	(0.0685)	(0.0681)	(0.0529)	(0.0604)	(0.0615)
$APARCH(\delta)$	1.9995 ***	1.2106 ***	1.8034 **	2.3373 ***	1.680 1***	1.0804 ***
	(0.1631)	(0.1589)	(0.7282)	(0.0577)	(0.1793)	(0.0971)
Asymmetry	-0.0438 **	0.0771 ***	0.0767 ***	0.0233	0.0613 **	0.0299
1 10 y 11 11 10 11 y	(0.0212)	(0.0242)	(0.0265)	(0.0214)	(0.0290)	(0.0228)
Tail	3.0358 ***	3.3493 ***	3.0959 ***	2.7428 ***	3.6221 ***	2.8861 ***
1	(0.1063)	(0.2109)	(0.4091)	(0.0706)	(0.2920)	(0.1458)

	BTC	DASH	ETH	LTC	XMR	XRP
		C. 1	Diagnostic checl	king		
logL	-7000.46	-7446.49	-5765.72	-7765.63	-7366.23	-7581.62
BĨC	5.1474	6.1259	6.0809	5.7030	6.3007	5.7759
Q(20)	59.43 ***	47.08 ***	50.19 ***	51.82 ***	49.49 ***	39.01 ***
$Q_{S}(20)$	5.92	9.85	15.02	8.51	13.32	2.28
ARCH(10)	0.38	0.67	1.14	0.06	1.05	0.10

Table A2. Cont.

Notes: See the notes for Table A1. *, ** and *** denote the rejection of the null hypothesis at the 1%, 5%, and 10% significance levels, respectively.

References

- 1. Urquhart, A. What causes the attention of Bitcoin? *Econ. Lett.* **2018**, *166*, 40–44. [CrossRef]
- 2. Urquhart, A. The inefficiency of Bitcoin. Econ. Lett. 2016, 148, 80-82. [CrossRef]
- 3. Tiwari, A.K.; Jana, R.K.; Das, D.; Roubaud, D. Informational efficiency of Bitcoin—An extension. *Econ. Lett.* **2018**, *163*, 106–109. [CrossRef]
- 4. Nadarajah, S.; Chu, J. On the inefficiency of Bitcoin. Econ. Lett. 2017, 150, 6–9. [CrossRef]
- 5. Bariviera, A.F. The inefficiency of Bitcoin revisited: A dynamic approach. *Econ. Lett.* **2017**, *161*, 1–4. [CrossRef]
- 6. Katsiampa, P. Volatility estimation for Bitcoin: A comparison of GARCH models. Econ. Lett. 2017, 158, 3–6. [CrossRef]
- 7. Takaishi, T. Statistical properties and multifractality of Bitcoin. *Phys. A* **2018**, *506*, 507–519. [CrossRef]
- 8. Sensoy, A. The inefficiency of Bitcoin revisited: A high-frequency analysis with alternative currencies. *Financ. Res. Lett.* **2019**, *28*, 68–73. [CrossRef]
- 9. Vidal-Tomás, D.; Ibañez, A. Semi-strong efficiency of Bitcoin. Financ. Res. Lett. 2018, 27, 259–265. [CrossRef]
- 10. Alvarez-Ramirez, J.; Rodriguez, E.; Ibarra-Valdez, C. Long-range correlations and asymmetry in the Bitcoin market. *Phys. A* **2018**, 492, 948–955. [CrossRef]
- 11. Dyhrberg, A.H. Bitcoin, gold and the dollar—A GARCH volatility analysis. Financ. Res. Lett. 2016, 16, 85–92. [CrossRef]
- 12. Alvarez-Ramirez, J.; Rodriguez, E. A singular value decomposition approach for testing the efficiency of Bitcoin and Ethereum markets. *Econ. Lett.* **2021**, *206*, 109997. [CrossRef]
- 13. Duan, K.; Li, Z.; Urquhart, A.; Ye, J. Dynamic efficiency and arbitrage potential in Bitcoin: A long-memory approach. *Int. Rev. Financ. Anal.* **2021**, *75*, 101725. [CrossRef]
- 14. López-Martín, C.; Benito Muela, S.; Arguedas, R. Efficiency in cryptocurrency markets: New evidence. *Eurasian Econ. Rev.* 2021, 11, 403–431. [CrossRef]
- 15. Kakinaka, S.; Umeno, K. Cryptocurrency market efficiency in short-and long-term horizons during COVID-19: An asymmetric multifractal analysis approach. *Financ. Res. Lett.* **2022**, *46*, 102319. [CrossRef]
- 16. Phiri, A. Can wavelets produce a clearer picture of weak-form market efficiency in Bitcoin? *Eurasian Econ. Rev.* **2022**, *12*, 373–386. [CrossRef]
- 17. Aggarwal, R.; Inclan, C.; Leal, R. Volatility in emerging stock markets. J. Financ. Quant. Anal. 1999, 34, 33–55. [CrossRef]
- 18. Lamoureux, C.G.; Lastrapes, W.D. Persistence in variance, structural change, and the GARCH model. *J. Bus. Econ. Stat.* **1990**, *8*, 225–234.
- 19. Lastrapes, W.D. Exchange rate volatility and U.S. monetary policy: An ARCH application. *J. Money Credit. Bank.* **1989**, *21*, 66–77. [CrossRef]
- Yıldırım, D.Ç.; Cevik, E.I.; Esen, Ö. Time-varying volatility spillovers between oil prices and precious metal prices. *Resour. Policy* 2020, 68, 101783. [CrossRef]
- 21. Caporale, G.M.; Karanasos, M.; Yfanti, S.; Kartsaklas, A. Investors' trading behaviour and stock market volatility during crisis periods: A dual long-memory model for the Korean Stock Exchange. *Int. J. Financ. Econ.* **2021**, *26*, 4441–4461. [CrossRef]
- 22. Luo, J.; Demirer, R.; Gupta, R.; Ji, Q. Forecasting oil and gold volatilities with sentiment indicators under structural breaks. *Energy Econ.* **2022**, *105*, 105751. [CrossRef]
- 23. Pan, Q.; Mei, X.; Gao, T. Modeling dynamic conditional correlations with leverage effects and volatility spillover effects: Evidence from the Chinese and US stock markets affected by the recent trade friction. *N. Am. J. Econ. Financ.* **2022**, *59*, 101591. [CrossRef]
- 24. Vougas, D.V. Analysing long memory and volatility of returns in the Athens stock exchange. *Appl. Financ. Econ.* **2004**, *14*, 457–460. [CrossRef]
- 25. Kang, S.H.; Yoon, S.-M. Long memory properties in return and volatility: Evidence from the Korean stock market. *Phys. A* 2007, 385, 591–600. [CrossRef]
- 26. Kasman, A.; Torun, E. Long memory in the Turkish stock market return and volatility. Cent. Bank Rev. 2007, 7, 13–27.
- Kasman, A.; Kasman, S.; Torun, E. Dual long memory property in returns and volatility: Evidence from the CEE countries' stock markets'. *Emerg. Mark. Rev.* 2009, 10, 122–139. [CrossRef]

- Mensi, W.; Hammoudeh, S.; Yoon, S.-M. Structural breaks and long memory in modeling and forecasting volatility of foreign exchange markets of oil exporters: The importance of scheduled and unscheduled news announcements. *Int. Rev. Econ. Financ.* 2014, 30, 101–119. [CrossRef]
- Marchese, M.; Kyriakou, I.; Tamvakis, M.; Di Iorio, F. Forecasting crude oil and refined products volatilities and correlations: New evidence from fractionally integrated multivariate GARCH models. *Energy Econ.* 2020, 88, 104757. [CrossRef]
- Rehman, M.U. Do bitcoin and precious metals do any good together? An extreme dependence and risk spillover analysis. *Resour. Policy* 2020, 68, 101737. [CrossRef]
- 31. Boubaker, H.; Saidane, B.; Ben Saad Zorgati, M. Modelling the dynamics of stock market in the gulf cooperation council countries: Evidence on persistence to shocks. *Financ. Innov.* **2022**, *8*, 1–22. [CrossRef]
- 32. González-Pla, F.; Lovreta, L. Modeling and forecasting firm-specific volatility: The role of asymmetry and long-memory. *Financ. Res. Lett.* **2022**, *48*, 102931. [CrossRef]
- 33. Lovcha, Y.; Perez-Laborda, A. Long-memory and volatility spillovers across petroleum futures. *Energy* **2022**, 243, 122950. [CrossRef]
- Mensi, W.; Hammoudeh, S.; Yoon, S.-M. Structural breaks, dynamic correlations, asymmetric volatility transmission, and hedging strategies for petroleum prices and USD exchange rate. *Energy Econ.* 2015, 48, 46–60. [CrossRef]
- 35. Inclán, C.; Tiao, G.C. Use of cumulative sums of squares for retrospective detection of changes of variance. *J. Am. Stat. Assoc.* **1994**, *89*, 913–923.
- 36. Bai, J.; Perron, P. Computation and analysis of multiple structural change models. J. Appl. Econom. 2003, 18, 1–22. [CrossRef]
- 37. Baillie, R.T.; Bollerslev, T.; Mikkelsen, H.O. Fractionally integrated generalized autoregressive conditional heteroskedasticity. *J. Econom.* **1996**, *74*, 3–30. [CrossRef]
- 38. Chung, C.-F. *Estimating the Fractionally Integrated GARCH Model;* Discussion Paper; National Taiwan University: New Taipei, Taiwan, 1999.
- 39. Bollerslev, T.; Mikkelsen, H.O. Modeling and pricing long memory in stock market volatility. *J. Econom.* **1996**, *73*, 151–184. [CrossRef]
- 40. Tse, Y.K. The conditional heteroscedasticity of the yen–dollar exchange rate. J. Appl. Econom. 1998, 13, 49–55. [CrossRef]
- 41. Nelson, D.B. Conditional heteroskedasticity in asset returns: A new approach. Econometrica 1991, 59, 347–370. [CrossRef]
- 42. Davidson, J. Moment and memory properties of linear conditional heteroscedasticity models, and a new model. *J. Bus. Econ. Stat.* **2004**, 22, 16–29. [CrossRef]
- 43. Bartos, J. Does Bitcoin follow the hypothesis of efficient market? Int. J. Econ. Sci. 2015, 4, 10–23. [CrossRef]
- 44. Brailsford, T.J.; Faff, R.W. An evaluation of volatility forecasting techniques. J. Bank. Financ. 1996, 20, 419–438. [CrossRef]
- 45. Brook, C.; Persand, G. Volatility forecasting for risk management. J. Forecast. 2003, 22, 1–22. [CrossRef]
- Degiannakis, S. Volatility forecasting: A fractional integrated asymmetric power ARCH skewed-t model. *Appl. Financ. Econ.* 2004, 14, 1333–1342. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.