

**Table S1:** Micronutrient intakes in adults with and without dyslipidemia

<b>Variables</b>	<b>Total population (n=610)</b>	<b>Dyslipidemia (n=75)</b>	<b>Non-Dyslipidemia (n=535)</b>	<b>P-value<sup>§</sup></b>
Vitamin C	155.2 (100.4, 231.1)	158.0 (109.2,237.0)	155.16 (99.0,230.5)	0.879
Vitamin D	4.5 (2.7, 7.1)	3.8 (2.5,7.1)	4.6 (2.8,7.2)	0.160
Vitamin E	8.5 (6.0, 11.9)	9.2 (6.1, 13.2)	8.4 (5.9,11.9)	0.135
Thiamin	2.4 (1.5, 3.7)	2.3 (1.3,3.7)	2.4 (1.5,3.7)	0.310
Riboflavin	2.4 (1.5, 4.7)	2.2 (1.4, 4.0)	2.4 (1.5, 4.2)	0.243
Pyridoxine	2.5 (1.8, 3.4)	2.4 (1.8,3.5)	2.5 (1.8,3.4)	0.976
Folate	327.3 (237.8, 467.4)	339.2 (248.8,488.8)	325.4 (236.1, 464.8)	0.484
vitamin B <sub>12</sub>	6.0 (3.4, 9.9)	7.02 (3.3,11.0)	5.8 (3.4, 9.8)	0.523
Calcium (mg)	787.6 (541.4, 1061.6)	711.8 (464.2,1076.8)	800.9 (544.4,1059.9)	0.214
Iron (mg)	14.0 (10.1, 19.0)	13.2 (9.7,19.3)	14.2 (10.1,19.0)	0.495

<sup>§</sup>Significance at  $\alpha=5\%$ ; Comparisons between variables were performed using Kruskal Wallis test. All values presented as median (25% -75%)