

Table S1. Number of portions (median (25th-75th percentile)) of food groups per tray consumed at the worksite canteens across the study and by time periods.

	Total n=30,384	T0 (June-Sept) n=7371	T1 (Sept-Dec) n=11,014	T2 (Jan-Apr) n=11,999	p-value¹ main effect
Water (portion/tray)	1.0 (1.0-1.0)	1.0 (1.0-1.0)	1.0 (1.0-1.0)	1.0 (1.0-1.0)	0.093
Beverages (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.495
Juice (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.579
Fruit (portion/tray)	1.0 (0.0-1.0)	1.0 (0.0-1.0) ^b	1.0 (0.5-1.0) ^a	1.0 (0.0-1.0) ^a	< 0.001
Vegetables (portion/tray)	1.0 (0.0-1.0)	1.0 (0.0-1.0)	1.0 (0.0-1.0)	1.0 (0.0-1.0)	0.180
Tubers (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^a	0.0 (0.0-0.0) ^b	< 0.001
Cereals (portion/tray)	1.0 (0.0-1.0)	1.0 (0.0-1.0) ^b	1.0 (0.0-1.0) ^b	1.0 (0.0-1.0) ^a	< 0.001
Whole-grain cereals (portion/tray)	0.0 (0.0-1.0)	0.0 (0.0-0.0) ^b	0.0 (0.0-1.0) ^a	0.0 (0.0-1.0) ^a	< 0.001
Pulses (portion/tray)	0.0 (0.0-0.5)	0.0 (0.0-0.0) ^b	0.0 (0.0-0.5) ^a	0.0 (0.0-0.5) ^a	< 0.001
Dairy products (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^a	< 0.001
Egg (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^a	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^a	< 0.001
Fish (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^b	0.0 (0.0-0.5) ^a	0.0 (0.0-0.0) ^{ab}	0.004
White meat (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^a	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^c	< 0.001
Red meat (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^a	0.0 (0.0-0.0) ^a	< 0.001
Cured meat (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^a	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^b	< 0.001
Desert (portion/tray)	0.0 (0.0-0.0)	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^b	0.0 (0.0-0.0) ^a	< 0.001

¹ Kruskal-Wallis test with pairwise comparisons. Different letters in the same row indicate significantly different values ($p < 0.05$).