



Figure S1. Effect of FDP dietary intervention on hepatic steatosis in normal or MASLD-induced mice. Representative photographs of liver tissues

Table S1. The primary antibodies used in immunoblotting analyses.

	Antibody	Dilution Factor	Corporation	Clone
Primary antibody	CYP2E1	1:5,000	Abcam	Rabbit
	iNOS	1:5,000	Abcam	Rabbit
	3NT	1:5,000	Abcam	Mouse
	FAS	1:1,000	Santa Cruz	Mouse
	PPARγ	1:1,000	Santa Cruz	Mouse
	SREBP-1	1:1,000	Santa Cruz	Mouse
	α-SMA	1:5,000	Sigma-Aldrich	Mouse
	MMP2	1:1,000	Santa Cruz	Rabbit
	MMP9	1:1,000	Santa Cruz	Mouse
	TGF-β	1:1,000	Santa Cruz	Mouse
	Collagen-1	1:1,000	Santa Cruz	Mouse
	Pro collagen-1	1:1,000	Santa Cruz	Mouse
	ZO-1	1:5,000	Abcam	Mouse
	Claudin-4	1:1,000	Santa Cruz	Mouse
	β-catenin	1:1,000	Santa Cruz	Mouse
	α-tubulin	1:1,000	Santa Cruz	Mouse
	E-cadherin	1:1,000	Santa Cruz	Mouse
	Bax	1:1,000	Santa Cruz	Mouse
	Occludin	1:1,000	Santa Cruz	Mouse
	Cleaved caspase 3	1:1,000	Cell Signaling	Rabbit
	P-JNK	1:1,000	Santa Cruz	Mouse
	JNK	1:1,000	Santa Cruz	Mouse
	GAPDH	1:1,000	Santa Cruz	Mouse