



Article

Fusarium Species and Mycotoxins Associated with Sorghum Grains in Uruguay

Ana Belén Corallo ¹, Agustina del Palacio ¹, María Oliver ¹, Susana Tiscornia ¹, Macarena Simoens ², Jaqueline Cea ², Inés de Aurrecochea ³, Inés Martínez ⁴ , Alicia Sanchez ⁴, Silvina Stewart ⁵  and Dinorah Pan ^{1,*}

- ¹ Sección Micología, Facultad de Ciencias Facultad de Ingeniería, Udelar, Julio Herrera y Reissig 565, Montevideo 11300, Uruguay; belencfabiano@gmail.com (A.B.C.); agustinadelpalacio@gmail.com (A.d.P.); moliver24@gmail.com (M.O.); tiscorniasusana@gmail.com (S.T.)
- ² Laboratorio Tecnológico del Uruguay, Departamento de Análisis de Productos Agropecuarios, Avenida Italia 6201, Montevideo 11500, Uruguay; msimoens@latu.org.uy (M.S.); jacqueline.cea6@gmail.com (J.C.)
- ³ Departamento de Granos, Dirección General de Servicios Agrícolas, Ministerio de Ganadería Agricultura y Pesca, Avenida Millán 4703, Montevideo 12900, Uruguay; ideaurrecochea@mgap.gub.uy
- ⁴ Latitud, Fundación del Laboratorio Tecnológico del Uruguay, Avenida Italia 6201, Montevideo 11500, Uruguay; imartin@latitud.org.uy (I.M.); asanchez@latitud.org.uy (A.S.)
- ⁵ Programa Cultivos de Secano, Instituto Nacional de Investigación Agropecuaria, Estación Experimental La Estanzuela, Ruta 50, Km 11, Colonia 70000, Uruguay; sstewart@inia.org.uy
- * Correspondence: dpan@fing.edu.uy

Abstract: Grain mold and stalk rot are among the fungal diseases that cause significant losses in sorghum worldwide and are caused by different *Fusarium* spp. The presence of *Fusarium* species in sorghum grains causes yield losses and mycotoxin contamination, which represents a risk to consumers. In this study, *Fusarium graminearum* species complex (FGSC) had a high incidence, followed by *Fusarium fujikuroi* species complex (FFSC) and *F. incarnatum-equiseti* species complex. Within FFSC, *F. proliferatum*, *F. andiyazi*, *F. fujikuroi*, *F. thapsinum*, *F. verticillioides* and *F. subglutinans* were identified, and this was the first report of *F. fujikuroi* in sorghum. The most frequent toxins found in sorghum samples were deoxynivalenol (DON) and zearalenone (ZEN). The presence of fumonisins and nivalenol (NIV) was detected at low levels. This study adds new knowledge about the occurrence of *Fusarium* species and mycotoxins in sorghum grains. Furthermore, this is the first report in Uruguay on fungicide sensitivity for *Fusarium* isolates from sorghum, which constitutes an important starting point for defining management practices to minimize fungal infection and mycotoxin contamination.

Keywords: *Fusarium graminearum*; *Fusarium fujikuroi* species complex; trichothecene; zearalenone; fumonisin; fungicide sensitivity

Key Contribution: *Fusarium graminearum* and *Fusarium fujikuroi* species complexes were isolated from sorghum in Uruguay. *Fusarium fujikuroi* was isolated for the first time from sorghum grains. The most frequent toxins found in sorghum samples were DON and ZEN. Metconazole was the fungicide that showed the best performance against *F. proliferatum* and *F. graminearum*.



Citation: Corallo, A.B.; del Palacio, A.; Oliver, M.; Tiscornia, S.; Simoens, M.; Cea, J.; de Aurrecochea, I.; Martínez, I.; Sanchez, A.; Stewart, S.; et al. *Fusarium* Species and Mycotoxins Associated with Sorghum Grains in Uruguay. *Toxins* **2023**, *15*, 484. <https://doi.org/10.3390/toxins15080484>

Received: 28 June 2023
Revised: 23 July 2023
Accepted: 25 July 2023
Published: 31 July 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Sorghum (*Sorghum bicolor* L. Moench), with a world production of 62×10^6 t in 2021, is one of the most important cereals [1,2].

With the discovery of sorghum's associated health benefits, food-grade sorghum has been produced and is becoming more popular in different regions of the world as a human food [3]. These nutritional qualities, in addition to increasing the profitability of sorghum, have increased the total cultivated area worldwide [4].

In Uruguay, sorghum is one of the three major grain fed summer crops, and it is used for feeding cattle, dairy cattle, poultry, and swine. It's is considered one of the main

protagonists, both for its stability and for its low production cost, which contributes to increasing the profitability of the livestock and dairy production system. Stems and leaves are used for silage, and grains are destined for animal feeding or bioethanol production [5]. For this reason, the sorghum cultivated area has increased by 237% from 2000 to 2021, with 233×10^3 t harvested in the 2020–2021 season [6].

Although sorghum is considered to have habitual resistance to microbial infection, fungal contamination is a major limitation for sorghum production [3]. Among the fungal diseases that causes significant losses in the world are grain mold and stalk rot, which are caused by different *Fusarium* spp. [7]. The presence of *Fusarium* species in sorghum grains causes yield losses and represents a risk to the health of consumers due to the capacity to produce mycotoxins, secondary toxic metabolites.

The *Fusarium fujikuroi* species complex (FFSC) is usually associated with sorghum in different countries [3,8–12]. In addition, some species of this complex can produce fumonisins (FBs), a group of mycotoxins responsible for the contamination of sorghum grains [13]. The presence of FBs in grains and feed has been associated with several diseases such as leukoencephalomalacia equine and porcine pulmonary edema. Further, human consumption has been epidemiologically associated with esophageal cancer and neural tube defects [14]. Other major *Fusarium* spp. associated with sorghum are the members of the *Fusarium graminearum* species complex (FGSC), which produce type B trichothecene mycotoxins such as deoxynivalenol (DON), nivalenol (NIV) and zearalenone (ZEN). DON is associated with feed refusal, vomiting, and suppressed immune functions, and ZEN can cause changes in reproductive organs and fertility loss in animals and humans due to the affinity for estrogen receptors [15,16].

Considering the above and the lack of information that exists in Uruguay, the aims of the present study were (1) to identify the *Fusarium* species associated with sorghum grains in the major sorghum-growing area in Uruguay, (2) to determine the presence of mycotoxins in sorghum samples, (3) to assess the ability of *Fusarium* spp. to produce mycotoxins, and (4) to evaluate the in vitro sensitivity of the main *Fusarium* species towards four fungicides corresponding to triazoles and benzimidazoles, currently used on several crops.

2. Results

2.1. Species Identification

A total of 1567 *Fusarium* spp. from the 2016 harvest and 1098 from the 2017 harvest were isolated in 95 and 72 sorghum grain samples, respectively.

In 2016, mean *Fusarium* spp. incidence in grain samples was 17%, with a maximum of 84%. During 2017, the incidence was lower (15%), although the maximum incidence was 100%. Among the *Fusarium* species isolated and based on morphological characteristics, FGSC had the highest incidence (10%, $n = 1744$), followed by species belonging to FFSC (4%, $n = 609$) and the *F. incarnatum-equiseti* species complex (FIESC) (2%, $n = 282$) (Figure 1).

Within FGSC, *F. graminearum* s.s was the predominant species, with only two isolates identified as *F. boothii*. Meanwhile, within the FFSC, 28% were identified as *F. proliferatum*, 23% as *F. andiyazi*, 21% as *F. fujikuroi*, 18% as *F. thapsinum*, 5% as *F. verticillioides* and 5% as *F. subglutinans* (Figure S1 Supplemental Material). Other identified species were *F. heterosporum* (2%), *F. tricinctum* (1%), *F. armeniacum* (1%), *F. pseudograminearum* (0.5%), *F. chlamydosporum* (0.5%) and *F. oxysporum* (0.2%).

2.2. Trichothecene Genotype Determination

The multiplex PCR assays indicated that most *F. graminearum* s.s isolates were of the 15AcDON genotype (98%), with only two isolates indicating NIV genotypes. No 3-AcDON genotypes were determined in the growing seasons studied.




Figure 1. Incidence (%) of the different *Fusarium* species infecting the sorghum samples during 2016 and 2017 as assessed by visual observations. Incidence was calculated as the percentage of grain infected out of the total analyzed grains per sorghum sample. *Fusarium graminearum* species complex (FGSC); *Fusarium fujikuroi* species complex (FFSC); *Fusarium incarnatum-equiseti* species complex (FIESC).

2.3. *Fusarium* Mycotoxin Production

The mycotoxin production ability of *F. graminearum* s.s and *F. proliferatum* isolates was analyzed, as they were the most frequent species. Of the *F. graminearum* s.s studied, ZEN was produced by 92% ($n = 86$) of the isolates, DON by 88% ($n = 83$) and NIV by 6% ($n = 6$). Production of FB1 and FB2 was detected in 75% ($n = 41$) and 64% ($n = 35$) of the *F. proliferatum* isolates, respectively. Thirteen *F. proliferatum* isolates (24%) did not produce fumonisins.

2.4. Determination of Mycotoxins in Sorghum

One hundred twelve sorghum samples harvested during 2016 and 2017 were analyzed for the presence of mycotoxins (Table 1). The percentage of DON-positive samples was high during the two years (88% and 93%), while mean DON content was low, varying from 1315 $\mu\text{g}/\text{kg}$ to 565 $\mu\text{g}/\text{kg}$, respectively. Regarding ZEN, positive samples (97 and 90%) and the mean content (1453 $\mu\text{g}/\text{kg}$ and 752 $\mu\text{g}/\text{kg}$) were high in both years of study. Also, the annual maximum levels ranged from 6560 $\mu\text{g}/\text{kg}$ to 9280 $\mu\text{g}/\text{kg}$. Low levels of FB1 and FB2 were also detected in the sorghum samples. Similarly, low levels of NIV were only detected in three samples from 2017.

Table 1. Contamination by toxins in sorghum samples harvested during 2016 and 2017 in Uruguay.

	Total Mean ($\mu\text{g}/\text{kg}$) \pm SD		Positive Samples (%)		Contamination in Positive Samples ($\mu\text{g}/\text{kg}$)			
	2016	2017	2016	2017	Mean \pm SD	Range	Mean \pm SD	Range
					2016		2017	
DON	1315 \pm 1597	565 \pm 567	88	93	1507 \pm 1623	59–6071	619 \pm 564	59–2133
NIV	nd *	4 \pm 19	0	5	--	--	79 \pm 36	50–120
ZEN	1453 \pm 2194	752 \pm 1297	97	90	1567 \pm 2239	30–9280	841 \pm 1344	30–6560
FB1	67 \pm 244	58 \pm 148	16	23	411 \pm 493	80–1693	255 \pm 217	80–753
FB2	5 \pm 36	13 \pm 98	2	2	--	80–270	--	80–742

nd *: not detected.

2.5. Fungicide Sensitivity Assay

EC₅₀ values of metconazole, tebuconazole, epoxiconazole and carbendazim were calculated for 20 isolates of *F. proliferatum* and *F. graminearum* s.s. Both *Fusarium* species showed different behaviors to the tested fungicides (Figures 2 and 3). The two *Fusarium* species investigated showed the highest sensitivity levels for metconazole, with EC₅₀ values below 0.50 mg/L and significantly different from those of the other fungicides tested ($p < 0.05$). Only one isolate of *F. proliferatum* was shown to be less sensitive to metconazole, with EC₅₀ values of 2.18 mg/L. Besides metconazole, EC₅₀ values lower than 0.75 mg/L were recorded for carbendazim and tebuconazole for all isolates of *F. proliferatum*, with mean EC₅₀ values of 0.53 mg/L and 0.33 mg/L, respectively. In the case of epoxiconazole, isolates S400 and S677 were observed to be the least sensitive, with EC₅₀ values of 0.99 and 1.04 mg/L, respectively.




Figure 2. Distribution of EC₅₀ (mg/L) values of *Fusarium proliferatum* isolates for epoxiconazole, tebuconazole, carbendazim and metconazole.

Generally, isolates of *F. graminearum* were shown to be less sensitive to the fungicides evaluated. The EC₅₀ values for triazole fungicide class ranged from 0.31 mg/L to 15.29 mg/L with a mean of 1.97 mg/L for epoxiconazole, 0.05 mg/L to 0.97 mg/L with a mean of 0.17 mg/L for metconazole and 0.14 mg/L to 3.54 mg/L with a mean of 1.06 mg/L for tebuconazole (Figure 3). Additionally, carbendazim EC₅₀ values ranged between 0.34 mg/L and 1.37 mg/L with a mean of 0.62 mg/L (Figure 3). In the case of epoxiconazole, isolate S501 was observed to have high EC₅₀ values of 15.29 mg/L. Also, isolates S58, S68 and S501 showed low levels of sensitivity for tebuconazole, with values of 2.76, 2.88 and 3.54 mg/L, respectively.




Figure 3. Distribution of EC50 (mg/L) values of *Fusarium graminearum s.s.* isolates for epoxiconazole, tebuconazole, carbendazim and metconazole.

3. Discussion

In Uruguay, little is known about the presence of fungi and mycotoxins in sorghum. Most of the studies that have been carried out have focused on describing the occurrence and incidence of *Fusarium* species in wheat and barley [17–19]. Only one study was found for this crop, and it was limited to silages [20]. Considering this, this work represents the first extended study about the occurrence of *Fusarium* species in sorghum grain in Uruguay. *Fusarium graminearum s.s.* appeared as the predominant species, followed by *F. proliferatum*, *F. andiyazi*, *F. fujikuroi* and *F. thapsinum*. Although all of these species have previously been present in sorghum in other parts of the world, our data indicate that *F. graminearum s.s.* is the main contaminant of sorghum, which differs from that reported in other studies where FFSC resulted in the most frequent contaminants [8–11,21,22]. This can be partially explained by the crop rotation system in Uruguay, where sorghum is generally planted after wheat or barley, so the prevalence of *F. graminearum s.s.* could be due to inoculum carryover from the previous crop. The presence of *F. graminearum* in sorghum grains has a great toxicological risk since this species can produce DON, 15-ADON, 3-ADON, NIV and ZEN. Also, infection of sorghum by species within FGSC can cause reduced germination and nutritional quality of the grains [23].

Within FFSC, *F. proliferatum*, *F. andiyazi* and *F. thapsinum* were the most frequent species, with all these species being reported for the first time in Uruguay. These species have been isolated from sorghum in Argentina, Australia, Brazil, Ethiopia, Nigeria, South Africa, Tunisia, and the USA and are well documented as pathogens of sorghum that cause grain mold and stalk rot [10,11,21,22,24–26]. However, frequencies of FFSC in this study were in contrast with species recovered from sorghum in other areas, including Argentina and Brazil, where *F. thapsinum* and *F. verticillioides* are the most frequent [8,11,12].

Fusarium fujikuroi was also recovered with high frequency within FFSC and, to our knowledge, this is the first report in sorghum. It has been isolated from rice, where it causes Bakanae disease, and from other crops such as soybeans, corn, barley, wheat, and peanuts.

These differences in the recovery and composition of FFSC may be associated with differences in agroclimatic and crop rotation systems associated with our country [26]. Moreover, little is known concerning the growth conditions, host range and pathogenicity of the *Fusarium* species associated with sorghum in Uruguay. This knowledge is essential for developing appropriate management strategies.

Fusarium graminearum s.s. with the DON/15ADON genotype was the predominant species and represents the first data of *F. graminearum* s.s. strains isolated from sorghum in Uruguay. Our results agree with previous reports from different crops in Uruguay and other regions of South America, where the DON-15ADON genotype prevails for *F. graminearum* s.s., showing the stability of genotype composition across crops and years [17,27,28].

Mycotoxins in sorghum samples showed contamination with FB1 and FB2, but in low levels, and none of the samples exceeded the limit established by the FDA for animal feed and the current regulations for other grains [29]. Incidence and DON levels found in the present study were high when compared to other reports from sorghum [30,31]. However, only 27% ($n = 30$) of samples had DON levels that exceeded the recommended levels in Uruguay for swine and equines (1000 $\mu\text{g}/\text{kg}$); 10% ($n = 11$), the levels for cattle consumption (2000 $\mu\text{g}/\text{kg}$); and 3% ($n = 3$), the levels for ruminants, sheep and poultry (5000 $\mu\text{g}/\text{kg}$) [32]. No sample exceeded the Uruguayan levels for raw materials destined for the production of animal feed or the FDA levels for grains destined for ruminants and chickens (10,000 $\mu\text{g}/\text{kg}$) [29,32]. Zearalenone was the most dominant mycotoxin and occurred at high levels and incidence rates, with 64% ($n = 72$) of the samples exceeding the recommended levels in Uruguay of 250 $\mu\text{g}/\text{kg}$ [32]. Although there are very few published works on ZEN contamination in sorghum worldwide, our results are similar to previous reports which positioned ZEN as the most frequent mycotoxin in sorghum [33–35]. High concentrations of DON and ZEN in our country could be explained by the fact that *F. graminearum* s.s. was the main contaminant species for sorghum. In this sense, we found a positive correlation between the incidence of *F. graminearum* and DON ($r = 0.521$, $p < 0.001$) and ZEN ($r = 0.406$, $p < 0.001$). In addition, climatic conditions, agricultural practices and the sorghum varieties used in Uruguay could also explain the levels of mycotoxins found.

In Uruguay, chemical protection for the control of *Fusarium* grain mold in sorghum is still not widely used, and little is known about the activity of fungicides used on crops, especially against FFSC. Therefore, this study provided information on fungicide sensitivity to two of the main toxigenic *Fusarium* species associated with sorghum crops. Metconazole was the fungicide that showed the best performance against *F. proliferatum* and *F. graminearum*, while epoxiconazole showed a lower effect compared to the other fungicides. In this sense, few studies have investigated the sensitivity of *Fusarium* spp. to epoxiconazole and particularly have been carried out with *F. graminearum* isolated from wheat. The sensitivity of *F. proliferatum* was similar to that reported by other authors in other geographical regions [36–38]. A single isolate of *F. proliferatum* exhibited EC50 values of 2.18 mg/L for metconazole, suggesting reduced sensitivity compared with the other isolates collected. However, except for this isolate, the sensitivity to metconazole was higher than those detected for the other fungicides. Regarding *F. graminearum* s.s. sensitivity, metconazole was the most effective fungicide in inhibiting the growth of *F. graminearum* s.s. isolates with EC50 values lower than 0.97 mg/L. However, the mean EC50 values found for epoxiconazole (1.97 mg/L) and tebuconazole (1.06 mg/L) were higher than those reported by other authors. For epoxiconazole, reported sensitivities in wheat from China ranged from 0.019 to 0.93 mg/L for isolates of *F. graminearum* [39]. In our case, 55% of the isolates had EC50 values higher than 0.93 mg/L, which was the highest value reported for Chinese isolates; moreover, we found one isolate that had an EC50 value of 15.29 mg/L. In the case of sensitivity to tebuconazole, our study showed a mean EC50 value (1.06 mg/L) higher than 0.891 mg/L and 0.29 mg/L, which were the mean values reported in other studies

for isolates of *F. graminearum* s.s. from barley and wheat in Uruguay, respectively [17,19]. Sensitivity surely increases as years go by in relation to numerous exposures, especially to control *Fusarium* head blight. Moreover, our results agree with those obtained from wheat isolates in Brazil, China, Croatia and the United States [40–43]. Additionally, our findings were similar to previous reports that the EC50 values of isolates for metconazole were lower than the EC50 values of the same isolates for tebuconazole and the other fungicides [44]. To our knowledge, this is the first fungicide sensitivity analysis of different classes of fungicides for *F. proliferatum* and *F. graminearum* isolates from sorghum and constitutes an important starting point for defining management practices to control *Fusarium* grain mold in this crop. Further studies are needed to investigate the effectiveness of fungicide applications in vivo and the genetic basis for the reduced sensitivity of *Fusarium* species to epoxiconazole and tebuconazole.

4. Conclusions

Sorghum in Uruguay is exposed to colonization by a wide variety of toxigenic *Fusarium* species that could cause grain contamination with different mycotoxins, mainly from ZEN. These findings indicate the need to establish monitoring programs for mycotoxin levels in sorghum to ensure animal health and implement strategies to minimize fungal infection in the field. In addition to this, metconazole was found to be the most effective fungicide for control of the main *Fusarium* species analyzed.

5. Materials and Methods

5.1. Sorghum Samples

A total of 167 sorghum samples were collected during two harvest seasons from fields in the main sorghum cultivation area of Uruguay (Figure 4). Ninety-five sorghum samples from the 2016 harvest season and 72 samples from the 2017 harvest season were analyzed. Samples were collected according to the ISO 24333: cereals and cereal products - sampling. Each 2 kg grain sample was divided into two subsamples (1 kg each), one for mycological analysis and the other for mycotoxin analysis.




Figure 4. Map depicting the location of departments in Uruguay where sorghum samples were harvested in two sorghum growing seasons, 2016 and 2017.

5.2. *Fusarium* Isolation

From each sample, 100 grains were taken and surface disinfected by placing them in 0.4% hypochlorite for 2 min, rinsing with sterile distilled water and drying on sterile filter

paper. Then, they were placed in Petri dishes, 10 grains per plate, with potato dextrose agar (PDA) culture medium (Oxoid) and incubated at 25 ± 2 °C for 5 days. Monosporic cultures were obtained from colonies identified morphologically as *Fusarium* spp. according to Leslie and Summerell [45]. A subset of representative isolates from different morphologically identified *Fusarium* species was selected for molecular identification, 223 isolates from the 2016 harvest and 221 isolates from the 2017 harvest. The isolates were stored at -80 °C in the fungal collection of the Sección Micología, Facultad de Ciencias, Universidad de la República.

5.3. Molecular Identification

Genomic DNA was extracted using the cetyltrimethylammonium bromide (2% CTAB) method [45].

Transcription elongation factor gene (*TEF 1- α*) RFLP analysis was performed to identify *Fusarium graminearum sensu stricto* (*F. graminearum s.s.*) from the remaining species within FGSC with a protocol previously described [46]. A fragment of the *TEF 1- α* gene was amplified by PCR with primers EF-1 and EF-2 [47]. The PCR cycling parameters used for the amplification were as follows: an initial denaturation cycle at 95 °C for 3 min; 35 cycles of denaturation (1 min at 95 °C), hybridization (1 min at 55 °C) and extension (1:30 min at 72 °C); a final extension cycle of 10 min at 72 °C and 5 min at 4 °C. *TEF 1- α* amplicons were digested for 45 min at 37 °C with restriction enzyme *Bsa*HI. RFLP patterns were visualized in a 1.5% agarose gel containing GreenSafe Premium (NZYTech, Lisboa, Portugal). Fragment sizes were estimated by comparison with a DNA standard marker GeneRuler 1 Kb DNA ladder (Thermo Scientific, Waltham, MA, USA).

For the other *Fusarium* species, the *TEF 1- α* gene was amplified in the same way as previously described, and PCR products were purified and sequenced by MacroGen Inc., Republic of Korea. Consensus sequences were obtained using the SeqMan software (Lasergene, Madison, WI, USA) and compared to GenBank database sequences using the Basic Local Alignment Search Tool (BLAST) and to the *Fusarium* MLST website (<https://fusarium.mycobank.org>) of CBS-KNAW Fungal Biodiversity Centre. The generated sequences were deposited in the GenBank database (Supplementary Table S1).

The sequences of FFSC were aligned with reference strain sequences from FFSC using the MUSCLE algorithm [48] with Mega version 11 [49]. A phylogenetic tree was inferred by using the maximum likelihood method, using Kimura's 2-parameter model (G + I) and the Tree-Bisection-Regrafting (TBR) algorithm [50]. *Fusarium oxysporum* strain CBS 132475 and *Fusarium inflexum* NRRL 20433 were used as an outgroup. The analyses were conducted with 1000 bootstrap replicates.

5.4. Trichothecene Genotype Determination

Three different PCR assays were used to determine the trichothecene genotypes of 93 isolates of *F. graminearum s.s.* First, two multiplex assays targeted a portion of trichothecene biosynthesis genes *Tri3* and *Tri12* to determine NIV, 15-ADON and 3-ADON genotypes, while the third assay targeted portions of the *Tri13* gene to determine DON genotypes [51,52]. PCR for *Tri3* and *Tri12* consisted of an initial denaturation of 2 min at 94 °C, followed by 25 cycles of 30 s at 94 °C, 30 s at 52 °C and 1 min at 72 °C. PCR amplification of *Tri13* consisted of an initial denaturation of 2 min at 94 °C, followed by 35 cycles of 30 s at 94 °C, 30 s at 65 °C and 30 s at 72 °C. The specific bands for each amplification were resolved on 1.5% agarose gels containing GreenSafe Premium (NZYTech) and scored by size in comparison to a 100 bp DNA size ladder (Invitrogen Life Technologies, Carlsbad, CA, USA).

5.5. *Fusarium* Mycotoxin Production

The ability of 93 isolates of *F. graminearum s.s.* and 55 isolates of *F. proliferatum* to produce mycotoxin was analyzed, as they were the most frequent species. For mycotoxin production assays, 25 g of rice grains was used for DON, NIV and ZEN or 20 g of rice

was used for FB1 and FB2; rice grains were placed in bags with 10 mL of distilled water, and then autoclaved twice at 121 °C for 30 min. Autoclaved rice bags were inoculated with 3 mycelium plugs taken from pure cultures of each selected isolate and incubated for 28 days at 25 °C in the dark. After this time, the contents of the bags were dried at 50 °C, finely ground with a laboratory blender and stored at −20 °C until toxin analysis. DON and NIV were detected using a method described by Reynoso et al., and ZEN and FBs were detected based on AOAC 985.18 and AOAC: 995.15 methods, respectively [53,54]. All toxins were analyzed using a high-performance liquid chromatography (HPLC) instrument connected to a reversed-phase C18 column (150 mm × 4.6 mm i.d., 5 µm particle size; Nucleodur, Macherey-Nagel, Düren, Germany) connected to a Security Guard pre-column (8 mm × 4 mm i.d., 5 µm particle size; Nucleodur, Macherey-Nagel, Düren, Germany). The HPLC system consisted of a Shimadzu LC-10ADvp pump. For DON and NIV, the HPLC system was coupled with a Shimadzu SPD photodiode array detector, while for ZEN and FBs, an RF-10AxI fluorescence detector was used. Toxin production was determined by comparison with external standards supplied by Trilogy Analytical Laboratory Inc., Washington, MO, USA.

5.6. Determination of Mycotoxins in Sorghum

Fifty-five representative samples of sorghum grain from the 2016 harvest and 57 samples from the 2017 harvest were selected for mycotoxin determination according to AOAC methods [54]. All toxins were determined using an HPLC instrument connected to a reversed-phase Gemini C18 (4.60 mm × 150 mm, 5 µm) analytical column (Phenomenex, Torrance, CA, USA). The HPLC system consisted of a Waters 1525 pump, Waters 717 Injector and Waters TCM oven. For FBs and ZEN, the HPLC system was coupled with a Waters 2475 Fluorescence detector, while for DON and NIV, a Waters 2996 photodiode array detector was used. For FB analysis, the excitation and emission wavelengths were set at 335 and 440 nm, respectively. The mobile phase used was methanol–NaH₂PO₄ 0.1 M (77:23 v/v) pH = 3.3, at a flow rate of 1.0 mL/min and at 25 °C. For ZEN analysis, the excitation and emission wavelengths were set at 270 and 465 nm, respectively. The mobile phase was methanol–orthophosphoric acid 0.01 M (58:42 v/v), the flow rate was 1.0 mL/min at 25 °C. For DON and NIV analysis, the mobile phase was water–methanol (86:14 v/v), at a flow of 1.0 mL/min and at 30 °C. Standards of DON, NIV, FB1, FB2 and ZEN from Trilogy (Washington, MO, USA) were used. The accuracy of the method was evaluated through studies of accuracy and precision by using sorghum grains fortified with each toxin (DON, NIV, ZEN, FB1 or FB2 as appropriate). For the accuracy of the method, replicates of certified samples were performed, and the relative standard deviation was calculated between them. The percentage of recovery of the method was 93% for DON, 87% for NIV, 76% for FB1, 77% for FB2 and 77% for ZEN.

5.7. Fungicide Sensitivity Assay

The sensitivity of *F. proliferatum* and *F. graminearum* s.s. isolates were evaluated against two different chemical classes of fungicides, triazoles and benzimidazoles. Fungicides belonging to triazoles were metconazole, tebuconazole and epoxiconazole, while carben-dazim belongs to the benzimidazole class. The assay was performed on a total of 20 isolates per species, 10 isolates from each crop season (2016 and 2017). Fungicides were diluted in dimethylsulfoxide (DMSO) and incorporated into PDA to achieve concentrations of 0.05, 0.25, 0.50, 0.75, 1.5, 3, 6 and 10 mg/L. Then, each PDA plate was inoculated with a mycelial plug in the center. *Fusarium proliferatum* isolate plates were incubated for 7 days at 25 °C in darkness, and *F. graminearum* s.s. strains were incubated for 5 days under the same conditions. Three replicates of each fungicide concentration per isolate were performed.

Evaluation was performed based on the radial growth on PDA containing different concentrations of each fungicide, compared to control plates (PDA + DMSO) [55,56]. The effective concentration of fungicides leading to a 50% inhibition (EC₅₀) of mycelial growth of each strain was determined. The EC₅₀ was calculated based on Probit analysis [57].

Analysis of variance (ANOVA) of the EC50 values was conducted to determine differences in sensitivity for each fungicide. Means were compared using Fisher's least significant difference (LSD) ($\alpha = 0.05$). Statistical analysis was performed using SigmaStat Version 3.5.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/toxins15080484/s1>, Figure S1: Maximum likelihood tree inferred from the transcription elongation factor gene (*TEF 1- α*) sequences of species belonging to the *Fusarium fujikuroi* species complex (FFSC). Table S1: GenBank accession numbers of the FFSC used in phylogenetic analysis.

Author Contributions: Conceptualization, A.B.C., S.S. and D.P.; Methodology, A.B.C., J.C. and D.P.; Formal Analysis, A.B.C. and D.P.; Investigation, A.B.C., A.d.P., M.O., S.T., M.S., I.d.A., I.M., A.S., S.S. and D.P.; Resources, S.S. and D.P.; Data Curation, A.B.C. and D.P.; Writing – Original Draft Preparation, A.B.C.; Writing – Review & Editing, M.S., I.d.A., A.S., S.S. and D.P.; Visualization, S.S. and D.P.; Supervision, S.S. and D.P.; Project Administration, S.S. and D.P.; Funding Acquisition, S.S. and D.P. Conceptualization, A.B.C., S.S. and D.P.; Methodology, A.B.C., J.C. and D.P.; Formal Analysis, A.B.C. and D.P.; Investigation, A.B.C., A.d.P., M.O., S.T., M.S., I.d.A., I.M., A.S., S.S. and D.P.; Resources, S.S. and D.P.; Data Curation, A.B.C. and D.P.; Writing – Original Draft Preparation, A.B.C.; Writing – Review & Editing, M.S., I.d.A., A.S., S.S. and D.P.; Visualization, S.S. and D.P.; Supervision, S.S. and D.P.; Project Administration, S.S. and D.P.; Funding Acquisition, S.S. and D.P. All authors have read and agreed to the published version of the manuscript.

Funding: This work was financed by Agencia Nacional de Investigación e Innovación (ANII) (grant number: FSA-I-2017-1-139531).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data will be made available on request.

Conflicts of Interest: The authors declare no conflict of interests.

References

1. FAOSTAT. FAOSTAT Statistical Database. 2015. Available online: <http://faostat3.fao.org/home/E> (accessed on 6 May 2022).
2. United States Department of Agriculture (USDA). World Agricultural Production. 2022. Available online: <https://apps.fas.usda.gov/psdonline/circulars/production.pdf> (accessed on 6 May 2022).
3. Astoreca, A.; Emateguy, G.; Alconada, T. Fungal contamination and mycotoxins associated with sorghum crop: Its relevance today. *Eur. J. Plant Pathol.* **2019**, *155*, 381–392. [[CrossRef](#)]
4. United States Department of Agriculture (USDA). Office of the Chief Economist. In *Agricultural Projections to 2024*; World Agricultural Outlook Board: Washington, DC, USA, 2015.
5. DIEA. Anuario estadístico agropecuario, Ministerio de Ganadería Agricultura y Pesca. *Montevideo. DIEA*. 2013. Available online: https://descargas.mgap.gub.uy/DIEA/Documentos%20compartmentados/Anuario2013/Diea_Anuario_2013.pdf (accessed on 10 August 2022).
6. Mehtol, M. Maiz y sorgo: Situación y perspectivas, 2021. In *Anuario de OPYPA de 2021*; Ministerio de Ganadería, Agricultura y Pesca: Montevideo, Uruguay, 2021; p. 193.
7. Ackerman, A.; Wennndt, A.; Boyles, R. The sorghum grain mold disease complex: Pathogens, host responses, and the bioactive metabolites at play. *Front. Plant Sci.* **2021**, *12*, 660171. [[CrossRef](#)] [[PubMed](#)]
8. Alves dos Reis, T.; Zorzete, P.; Rodrigues Pozzi, C.; Nascimento da Silva, V.; Ortega, E.; Correa, B. Mycoflora and fumonisin contamination in Brazilian sorghum from sowing to harvest. *J. Sci. Food Agric.* **2010**, *90*, 1445–1451. [[CrossRef](#)] [[PubMed](#)]
9. Divakara, S.; Santosh, P.; Aiyaz, M.; Ramana, M.; Hariprasad, P.; Nayaka, S.; Niranjana, S. Molecular identification and characterization of *Fusarium* spp. associated with sorghum seeds. *J. Sci. Food Agric.* **2014**, *94*, 1132–1139. [[CrossRef](#)]
10. Lahouar, A.; Crespo-Sempere, A.; Marín, S.; Saïd, S.; Sanchis, V. Toxicogenic molds in Tunisian and Egyptian sorghum for human consumption. *J. Stored Prod. Res.* **2015**, *63*, 57–62. [[CrossRef](#)]
11. Mohamed Nor, N.M.I.; Salleh, B.; Leslie, J.F. *Fusarium* Species from Sorghum in Thailand. *Plant Pathol. J.* **2019**, *35*, 301–312. [[CrossRef](#)]
12. Pena, G.A.; Cavaglieri, L.R.; Chulze, S.N. *Fusarium* species and moniliformin occurrence in sorghum grains used as an ingredient for animal feed in Argentina. *J. Sci. Food Agr.* **2019**, *99*, 47–54. [[CrossRef](#)]
13. Leslie, J.F.; Zeller, K.A.; Lamprecht, S.C.; Rheeder, J.P.; Marasas, W.F. Toxicity, pathogenicity, and genetic differentiation of five species of *Fusarium* from sorghum and millet. *Phytopathology* **2005**, *95*, 275–283. [[CrossRef](#)]

14. Marasas, W.F.O.; Rheeder, J.P.; Lamprecht, S.C.; Zeller, K.A.; Leslie, J.F. *Fusarium andiyazi* sp. nov., a new species from sorghum. *Mycologia* **2001**, *93*, 1203–1210. [CrossRef]
15. McCormick, S.P.; Stanley, A.M.; Stover, N.A.; Alexander, N.J. Trichothecenes: From simple to complex mycotoxins. *Toxins* **2011**, *3*, 802–814. [CrossRef]
16. Zinedine, A.; Soriano, J.M.; Molto, J.C.; Manes, J. Review on the toxicity, occurrence, metabolism, detoxification, regulations and intake of zearalenone: An oestrogenic mycotoxin. *Food Chem. Toxicol.* **2007**, *45*, 1–18. [CrossRef]
17. Garmendia, G.; Pattarino, L.; Negrin, C.; Martinez-Silveira, A.; Pereyra, S.; Ward, T.; Vero, S. Species composition, toxigenic potential and aggressiveness of *Fusarium* isolates causing Head blight of barley in Uruguay. *Food Microbiol.* **2018**, *76*, 426–433. [CrossRef]
18. Pan, D.; Mionetto, A.; Calero, N.; Reynoso, M.M.; Torres, A.; Bettucci, L. Population genetic analysis and trichothecene profiling of *Fusarium graminearum* from wheat in Uruguay. *Genet. Mol. Res.* **2016**, *15*, 1–11. [CrossRef]
19. Umpiérrez-Failache, M.; Garmendia, G.; Pereyra, S.; Rodríguez-Haralambides, A.; Ward, T.J.; Vero, S. Regional differences in species composition and toxigenic potential among *Fusarium* head blight isolates from Uruguay indicate a risk of nivalenol contamination in new wheat production areas. *Int. J. Food Microbiol.* **2013**, *166*, 135–140. [CrossRef]
20. del Palacio, A.; Mionetto, A.; Bettucci, L.; Pan, D. Evolution of fungal populations and mycotoxins in sorghum silage. *Food Addit. Contam. A* **2016**, *33*, 1864–1872. [CrossRef]
21. Kelly, L.; Tan, Y.; Ryley, M.; Aitken, E. *Fusarium* species associated with stalk rot and head blight of grain sorghum in Queensland and New South Wales. *Plant Pathol.* **2017**, *66*, 1413–1423. [CrossRef]
22. Sharma, R.; Thakur, R.P.; Senthilvel, S.; Nayak, S.; Reddy, S.V.; Rao, V.P.; Varshney, R.K. Identification and characterization of toxigenic *Fusaria* associated with sorghum grain mold complex in India. *Mycopathologia* **2011**, *171*, 223–230. [CrossRef]
23. Balota, M. Sorghum diseases, head mold. Virginia Polytechnic Institute and State University, Extension Program, Virginia State, Petersburg, VA, USA. 2012.
24. Funnell-Harris, L.D.; Pedersen, J.F. Presence of *Fusarium* spp. in air and soil associated with sorghum fields. *Plant Dis.* **2011**, *95*, 648–656. [CrossRef]
25. Leslie, J.F.; Marasas, W.F.O.; Diourté, M. Naturally occurring mycotoxins in sorghum and millet from Mali (abstract). In Proceedings of the INTSORMIL Principal Investigators Conference, Addis Ababa, Ethiopia, 18–20 November 2002.
26. Petrovic, T.; Walsh, J.L.; Burgess, L.W.; Summerell, B.A. *Fusarium* species associated with stalk rot of grain sorghum in the northern grain belt of eastern Australia. *Aust. Plant Pathol.* **2009**, *38*, 373–379. [CrossRef]
27. Pan, D.; Calero, N.; Mionetto, A.; Bettucci, L. Trichothecene genotypes of *Fusarium graminearum* from wheat in Uruguay. *Int. J. Food Microbiol.* **2013**, *162*, 120–123. [CrossRef]
28. Del Ponte, E.M.; Moreira, G.M.; Ward, T.J.; O'Donnell, K.; Nicolli, C.P.; Machado, F.J.; Duffeck, M.R.; Alves, K.S.; Tessmann, D.J.; Waalwijk, C.; et al. *Fusarium graminearum* Species Complex: A Bibliographic Analysis and Web-Accessible Database for Global Mapping of Species and Trichothecene Toxin Chemotypes. *Phytopathology* **2022**, *112*, 741–751. [CrossRef] [PubMed]
29. FDA. Mycotoxins. 2010. Available online: <https://www.fda.gov/food/natural-toxins-food/mycotoxins> (accessed on 4 May 2022).
30. Chilaka, C.; De Boevre, M.; Atanda, O.; De Saeger, S. The status of *Fusarium* mycotoxins in Sub-Saharan Africa: A review of emerging trends and post-harvest mitigation strategies towards food control. *Toxins* **2017**, *9*, 19. [CrossRef] [PubMed]
31. Hanvi, D.; Lawson-Evi, P.; De Boever, M.; Goto, C.E.; De Saeger, S.; Eklu-Gadegbeku, K. Natural occurrence of mycotoxins in maize and sorghum in Togo. *Mycotoxin Res.* **2019**, *35*, 321–327. [CrossRef] [PubMed]
32. Resolución S/N/001. 2001. Límites Máximos de DON en Alimentos Para Animales. 2001. Available online: <https://www.gub.uy/ministerio-ganaderia-agricultura-pesca/institucional/normativa/resolucion-sn001-limites-maximos-don-alimentos-para-animales> (accessed on 6 May 2022).
33. Aoyama, K.; Ishikuro, E.; Nishiwaki, M.; Ichinoe, M. Zearalenone contamination and the causative fungi in sorghum. *Shokuhin Eiseigaku Zasshi* **2009**, *50*, 47–51. [CrossRef]
34. Ayalew, A.; Fehrmann, H.; Lepschy, J.; Beck, R.; Abate, D. Natural occurrence of mycotoxins in staple cereals from Ethiopia. *Mycopathologia* **2006**, *162*, 57–63. [CrossRef]
35. Chala, A.; Taye, W.; Ayalew, A.; Krska, R.; Sulyok, M.; Logrieco, A. Multi mycotoxin analysis of sorghum (*Sorghum bicolor* L. Moench) and finger millet (*Eleusine coracana* L. Garten) from Ethiopia. *Food Control* **2014**, *45*, 29–35. [CrossRef]
36. Cendoya, M.; Nichea, J.; Monge, M.; Zachett, V.; Stella, M.; Ramirez, M. Effect of fungicides commonly used for *Fusarium* head blight management on growth and fumonisin production by *Fusarium proliferatum*. *Rev. Argent. Microbiol.* **2021**, *53*, 64–74. [CrossRef]
37. Marín, P.; De Ory, A.; Cruz, A.; Magan, N.; González-Jaén, M.T. Potential effects of environmental conditions on the efficiency of the antifungal tebuconazole controlling *Fusarium verticillioides* and *Fusarium proliferatum* growth rate and fumonisin biosynthesis. *Int. J. Food Microbiol.* **2013**, *165*, 251–258. [CrossRef]
38. Xu, S.; Wang, J.; Wang, H.; Bao, Y.; Li, Y.; Govindaraju, M.; Yao, W.; Chen, B.; Zhang, M. Molecular characterization of carbendazim resistance of *Fusarium* species complex that causes sugarcane pokkah boeng disease. *BMC Genomics* **2019**, *20*, 115. [CrossRef]
39. Liu, S.; Fu, L.; Chen, J.; Chen, J.; Wang, S.; Liu, J.; Jiang, J.; Che, Z.; Tian, Y.; Chen, G. Baseline sensitivity and control efficacy of epoxiconazole against *Fusarium graminearum* in Henan Province, China. *Eur J Plant Pathol.* **2020**, *157*, 825–833. [CrossRef]
40. Ivić, D.; Sever, Z.; Kumanovska, B. In vitro sensitivity of *Fusarium graminearum*, *F. avenaceum* and *F. verticillioides* to carbendazim, tebuconazole, flutriafol, metconazole and prochloraz. *Pestic i Phytomed (Belgrade)* **2011**, *26*, 35–42. [CrossRef]

41. Machado, F.J.; Santana, F.M.; Lau, D.; Del Ponte, E.M. Quantitative Review of the effects of triazole and benzimidazole fungicides on *Fusarium* head blight and wheat yield in Brazil. *Plant Dis.* **2017**, *101*, 1633–1641. [[CrossRef](#)]
42. Spolti, P.; Del Ponte, E.M.; Dong, Y.; Cummings, J.A.; Bergstrom, G.C. Triazole sensitivity in a contemporary population of *Fusarium graminearum* from New York wheat and competitiveness of a tebuconazole-resistant isolate. *Plant Dis.* **2014**, *98*, 607–613. [[CrossRef](#)]
43. Yin, Y.; Liu, X.; Li, B.; Ma, Z. Characterization of sterol demethylation inhibitor-resistant isolates of *Fusarium asiaticum* and *F. graminearum* collected from wheat in China. *Phytopathology* **2009**, *99*, 487–497. [[CrossRef](#)]
44. Anderson, N.; Freije, A.; Bergstrom, G.; Bradley, C.; Cowger, C.; Faske, T.; Hollier, C.; Kleczewski, N.; Padgett, G.; Paul, P.; et al. Sensitivity of *Fusarium graminearum* to Metconazole and Tebuconazole Fungicides Before and After Widespread Use in Wheat in the United States. *Plant Health Prog.* **2020**, *21*, 85–90. [[CrossRef](#)]
45. Leslie, J.F.; Summerell, B.A. *The Fusarium Laboratory Manual*; Blackwell Professional: Ames, IA, USA, 2006.
46. Garmendia, G.; Umpierrez-Failache, M.; Ward, T.; Vero, S. Development of a PCR-RFLP method based on the transcription elongation factor 1- α gene to differentiate *Fusarium graminearum* from other species within the *Fusarium graminearum* species complex. *Food Microbiol.* **2018**, *70*, 28–32. [[CrossRef](#)]
47. O'Donnell, K.; Cigelnik, E.; Nirenberg, H.I. Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* **1998**, *90*, 465–493. [[CrossRef](#)]
48. Edgar, R.C. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **2004**, *32*, 1792–1797. [[CrossRef](#)]
49. Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis version 11. *Molecular Biology and Evolution* **2021**, *38*, 3022–3027. [[CrossRef](#)]
50. Nei, M.; Kumar, S. *Molecular Evolution and Phylogenetics*; Oxford University Press: New York, NY, USA, 2000.
51. Ward, T.J.; Bielawski, J.P.; Kistler, H.C.; Sullivan, E.; O'Donnell, K. Ancestral polymorphism and adaptive evolution in the trichothecene mycotoxin gene cluster of phytopathogenic *Fusarium*. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 9278–9283. [[CrossRef](#)]
52. Chandler, E.; Simpson, D.; Thomsett, M.; Nicholson, P. Development of PCR Assays to Tri7 and Tri13 Trichothecene Biosynthetic Gene Regions in *Fusarium graminearum* and Characterization of Chemotypes of *Fusarium*. *Physiol. Mol. Plant Pathol.* **2003**, *62*, 355–367. [[CrossRef](#)]
53. Reynoso, M.M.; Ramirez, M.L.; Torres, A.M.; Chulze, S.N. Trichothecene genotypes and chemotypes in *Fusarium graminearum* strains isolated from wheat in Argentina. *Int J Food Microbiol.* **2011**, *145*, 444–448. [[CrossRef](#)] [[PubMed](#)]
54. AOAC. *International Official Methods of Analysis of AOAC International*, 20th ed.; AOAC: Gaithersburg, MD, USA, 2016.
55. Becher, R.; Hettwer, U.; Karlovsky, P.; Deising, H.B.; Wirsal, S. Adaptation of *Fusarium graminearum* to tebuconazole yielded descendants diverging for levels of fitness, fungicide resistance, virulence, and mycotoxin production. *Phytopathology* **2010**, *100*, 444–453. [[CrossRef](#)]
56. Tateishi, H.; Miyake, T.; Mori, M.; Kimura, R.; Sakuma, Y.; Saishoji, T. Sensitivity of Japanese *Fusarium graminearum* species complex isolates to metconazole. *J. Pestic. Sci.* **2010**, *35*, 419–430. [[CrossRef](#)]
57. Finney, D.J. Probit Analysis (2nd Ed). *J. Inst. Actuar.* **1952**, *78*, 388–390.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.