

Supplementary Materials: Evaluation of Carbon Ion Radiation-Induced Trismus in Head and Neck Tumors Using Dose-Volume Histograms

Atsushi Musha ^{1,2,*}, Hirofumi Shimada ¹, Nobuteru Kubo ¹, Hidemasa Kawamura ¹, Naoko Okano ¹, Yuhei Miyasaka ¹, Hiro Sato ¹, Katsuyuki Shirai ³, Jun-ichi Saitoh ⁴, Satoshi Yokoo ², Kazuaki Chikamatsu ⁵ and Tatsuya Ohno ¹

Table S1. Univariate analysis of risk factors for carbon ion radiotherapy-induced trismus.

Parameter	<i>n</i>	<i>p</i> -Value	Odds Ratio (95% CI)
Age(years)			
59 ≤	14	0.517	1.846(0.285-11.978)
59 >	17		
Gender			
man	14	0.517	1.846(0.285-11.978)
woman	17		
Primary site			
Maxillary sinus and Nasal cavity	16	0.93	1.083(0.182-6.439)
others	15		
Histological type			
Adenoid cystic carcinoma	17	0.118	0.185(0.19-1.816)
others	14		
T stage			
T2-3	12	0.527	0.563(0.93-3.391)
T4	19		
GTV(Gross tumor volumes, cm ³)			
28.68 ≤	16	0.411	2.167(0.334-14.057)
28.68 >	15		