Globular Adiponectin Inhibits Breast Cancer Cell Growth through Modulation of Inflammasomes Activation: Critical Role of Sestrin2 and AMPK Signaling

Duc-Vinh Pham, Pawan Kumar Raut, Mahesh Pandit, Jae-Hoon Chang, Nikita Katila, Dong-Young Choi, Jee-Heon Jeong and Pil-Hoon Park

Supplementary Materials




Figure S1. Effect of globular adiponectin on SESN2 mRNA expression level in breast cancer cells. MCF-7 cells were treated with gAcrp (1 μ g/mL) for indicated time periods. Sestrin2 mRNA levels were determined by qT-PCR analysis.








Figure S2. Figure S2. Effect of inflammasomes inhibitors on the viability of breast cancer cells. T47D cells (A) or MDA-MB-231 cells (B) were treated with Ac-YVAD-cmk, MCC950, or IL-1Ra for indicated time durations. Cell viability was determined by MTS assay.















Whole Western blots for Figure 2














Whole Western blots for Figure 3

































Whole Western blots for Figure 5












Whole Western blots for Figure 6










Figure S3. Blots from Western blot analysis.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).