

Table S1. Primers used for qPCR.

Gene	Forward (5'→3')	Reverse (5'→3')
BGN	CCCTCTCCAGGTCCATCCGC	GAGCTGGGTAGGTTGGGCGGG
CK19	ATGGCCGAGCAGAACCGGAA	CCATGAGCCGCTGGTACTCC
CLDN7	AGCTGCAAAATGTACGACTCG	GGAGACCACCATTAGGGCTC
ECAD	TCTTCCCCGCCCTGCCAATC	GCCTCTCTCGAGTCCCCTAG
EPCAM	AATCGTCAATGCCAGTGTACTT	TTCATCGCAGTCAGGATCATAA
GAPDH	TTGCCATCAATGACCCCTTCA	CGCCCCACTTGATTTTGGA
GRHL2	GGGGAAGAGCAACGAGTGG	GCGCTGGTCGTCTTTGAGA
INS	GCAGCCTTTGTGAACCAACAC	CCCCGCACACTAGGTAGAGA
NEUROG 3	GGCGCTGACTCAAACGCTGC	AAGCCAGACTGCCTGGGCTC
OVOL2	ACAGGCATTCGTCCCTACAAA	CGCTGCTTATAGGCATACTGC
RAC1	ACCATGCAGGCCATCAAGTGTGTGG	TTACAACAGCAGGCATTTTCTCT TC
RAC1b	GGAGAAACGTACGGTAAGGATATAA CC	GGCAATCGGCTTGTCTTTGCCC
SERPINE 1	CTTCTTCAGGCTGTTCCGGAGC	GGGTCAGGGTTCCATCACTTGG
SNAI1	CTGCTCCACAAGCACCAAGAGTC	CCAGCTGCCCTCCCTCCAC
SNAI2	ATATTCGGACCCACACATTACCT	GCAAATGCTCTGTTGCAGTGA
TBP	GCTGGCCCATAGTGATCTTT	CTTCACACGCCAAGAAACAG
VIM	TGGCAGGTCTTGACCTTGAA	GGTCATCGTGATGCTGAGAA