

SUPPLEMENTARY MATERIAL: The relationship between the four most common symptoms  
A) *neck lump*, B) *dysphagia*, C) *oral exulceration* and the two study periods.

*Supplementary Table S1A*

	<b>Neck lump</b>	<b>Other</b>	<b>p</b>
<b>Before Covid-19</b> <b>n=402 (%)</b>	110 (27.4)	292 (72.6)	0.514
<b>During Covid-19</b> <b>n=123 (%)</b>	30 (24.4)	93 (75.6)	
<b>Total n=525 (%)</b>	140 (26.7)	385 (73.3)	

*Supplementary Table S1B*

	<b>Pain</b>	<b>Other</b>	<b>p</b>
<b>Before Covid-19</b> <b>n=402 (%)</b>	103 (25.6)	299 (74.4)	0.926
<b>During Covid-19</b> <b>n=123 (%)</b>	31 (25.2)	92 (74.8)	
<b>Total n=525 (%)</b>	134 (25.5)	391 (74.5)	

*Supplementary Table S1C*

	<b>Dysphagia</b>	<b>Other</b>	<b>p</b>
<b>Before Covid-19</b> <b>n=402 (%)</b>	99 (24.6)	303 (75.4)	0.241
<b>During Covid-19</b> <b>n=123 (%)</b>	24 (19.5)	99 (80.5)	
<b>Total n=525 (%)</b>	123 (23.4)	402 (76.6)	

*Supplementary Table S1D*

	<b>Exulceration in the oral cavity</b>	<b>Other</b>	<b>p</b>
<b>Before Covid-19</b> <b>n=402 (%)</b>	85 (21.1)	317 (78.9)	0.256
<b>During Covid-19</b> <b>n=123 (%)</b>	32 (26%)	91 (74)	
<b>Total n=525 (%)</b>	117 (22.3)	408 (77.7)	