

Effect of the deposition of vanadium-oxide on the photocatalytic activity of TiO₂ nanotubes and its photodiode performance interfaced with CH₃NH₃PbI₃ single crystal

Jelena Vujančević^{1†}, Pavao Andričević^{2,3}, Veljko Djokić^{2,4}, Vladimir Blagojević⁵, Vera P. Pavlović⁶, Jovana Ćirković⁷, Endre Horváth^{2,8}, László Forró^{2,9}, Abdennaceur Karoui¹⁰, Vladimir B. Pavlović¹¹ and Djordje Janačković⁴*

¹ Institute of Technical Sciences of SASA, Knez Mihailova 35/IV, 11000 Belgrade, Serbia;

² Ecole Polytechnique Fédérale de Lausanne, Laboratory of Physics of Complex Matter, CH-1015 Lausanne, Switzerland;

³ Department of Physics, Technical University of Denmark, DTU Risø Campus, 4000 Roskilde, Denmark

⁴ University of Belgrade, Faculty of Technology and Metallurgy, Karnegijeva 4, 11120 Belgrade, Serbia

⁵ TBP software, Solunska 11, 11000 Belgrade, Serbia

⁶ University of Belgrade, Faculty of Mechanical Engineering, Kraljice Marije 16, 11120 Belgrade, Serbia

⁷ University of Belgrade, Institute for Multidisciplinary Research, Kneza Višeslava 1, 11000 Belgrade, Serbia

⁸ Haute école du paysage, d'ingénierie et d'architecture de Genève – HEPIA, 1202 Genève, Switzerland

⁹ Department of Physics and Astronomy, University of Notre Dame, 433 Nieuwland Science Hall
Notre Dame, IN 46556, USA

¹⁰ North Carolina Central University, Department of Physics, 1801 Fayetteville St Durham, North Carolina 27707, USA

¹¹ University of Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Belgrade-Zemun, Serbia

*Corresponding author. jelena.vujancevic@itn.sanu.ac.rs




Figure S1. Survey XPS spectra of TiO₂ nanotubes after vanadium oxide deposition.

Table S1. Band gaps of the samples with and without vanadium oxide deposition.

Samples	T-10	T-15	T-20	TV-10	TV-15	TV-20
Band gap, eV	3.44	3.40	3.20	3.27	3.38	3.08