

TiO₂/p-BC Composite Photocatalyst for Efficient Removal of Tetracycline from Aqueous Solutions under Simulated Sunlight

Jianhui Liu ¹, Liwen Zheng ^{1,*}, Yongchao Gao ^{1,*}, Lei Ji ¹, Zhongfeng Yang ¹, Hailun Wang ¹, Ming Shang ¹, Jianhua Du ² and Xiaodong Yang ³

¹ Ecology Institute, Qilu University of Technology(Shandong Academy of Sciences), Jinan 250103, China; 15665891128@163.com (J.L.); jilei1010@qlu.edu.cn (L.J.); yangzf@qlu.edu.cn (Z.Y.); wanghl0215@163.com (H.W.); sm17667939671@163.com (M.S.)

² WSP Australia Pty Limited, Level 3, Mia Yellagonga Tower 2, 5 Spring Street, Perth 6000, Australia; jason.du@wsp.com

³ Department of Geography & Spatial Information Technology, Ningbo University, Ningbo 315211, China; yangxiaodong@nbu.edu.cn

* Correspondence: zhenglw@qlu.edu.cn (L.Z.); gaoyc@sdas.org (Y.G.)




Figure S1 *m/z* of degrading TC over TiO₂/p-BC composite photocatalyst in different pathways (a-d) and *m/z* of the degraded small molecular fragments (e,f)