catalysts

Supplementary Materials: Mixed Zinc/Manganese on Highly Reduced Graphene Oxide: A Highly Active Nanocomposite Catalyst for Aerial Oxidation of Benzylic Alcohols

Mohamed E. Assal [1], Mohammed Rafi Shaik [1], Mufsir Kuniyil [1,2], Mujeeb Khan [1], Abdulrahman Yahya Alzahrani [3], Abdulrahman Al-Warthan [1], Mohammed Rafiq H Siddiqui [1,*] and Syed Farooq Adil [1,*]

[1] Department of Chemistry, College of Science, King Saud University, P.O. 2455, Riyadh 11451, Saudi Arabia; mhd.elshahat@gmail.com (M.E.A.); rafiskm@gmail.com (M.R.S.); mufsir@gmail.com (Muf.K.); kmujeeb@ksu.edu.sa (Muj.K.); awarthan@ksu.edu.sa (A.A.-W.)
[2] Department of Chemistry, K L University, Guntur 522502, Andhra Pradesh, India
[3] Petrochemical Research Centre, King Abdul-Aziz City for Science and Technology, Riyadh 11451, Saudi Arabia; aalzahrani@kacst.edu.sa
* Correspondence: rafiqs@ksu.edu.sa (M.R.H.S.); sfadil@ksu.edu.sa (S.F.A.); Tel.: +966-11-4676-082 (M.R.H.S.); +966-11-4670-439 (S.F.A)



Figure S1. XRD patterns of ZnO_x(1%)–$MnCO_3$/(1%)HRG catalyst (a) fresh and (b) recycled.



Figure S2. TGA isothermal pattern of ZnO_x(1%)–$MnCO_3$/(1%)HRG catalyst.