

Supporting Information for Publication

Effects of Charge Density on Photophysics and Aggregation Behaviour of Anionic Fluorene-arylene Conjugated Polyelectrolytes

Liliana M. Martelo^{1,*,#}, Sofia M. Fonseca¹, Ana T. Marques², Hugh D. Burrows^{1,*}, Artur J. M. Valente¹, Licinia L. G. Justino¹, Ullrich Scherf², Swapna Pradhan², Qiu Song^{2, §}.

¹ Departamento de Química, Universidade de Coimbra, 3004-535 Coimbra (Portugal); liliana.martelo@tecnico.ulisboa.pt (L.M.M.); sfonseca@qui.uc.pt (S.M.F.); burrows@ci.uc.pt (H.D.B.); avalente@ci.uc.pt (A.J.M.V.); liciniaj@ci.uc.pt (L.L.G.J.)

² Makromolekulare Chemie, Bergische Universität Wuppertal, DE-42097 Wuppertal (Germany); anatmarques@yahoo.com (A.T.M.); scherf@uni-wuppertal.com (U.S.); swapna_shivasan@yahoo.com (S.P.); squi2010@sinano.ac.com (Q.S.)

Present address:

[#] Centro de Química-Física Molecular, Instituto Superior Técnico, 1049-001 Lisbon, Portugal

[§] Printable Electronics Research Center, Suzhou Institute of Nano-Tech and Nano-Bionics, CAS, 215123, Suzhou, P. R. China

* Correspondence: liliana.martelo@tecnico.ulisboa.pt (LMM); Tel.: +351-218-419-259 and burrows@ci.uc.pt (HDB); Tel.: +351-239-852-083




Figure S1 Normalised fluorescence spectra of aqueous solutions (ca. 50 μM in terms of repeat units) of PBS-PPF (λ_{excit} 370 nm), PBS-PPF2 (λ_{excit} 365 nm) and PBS-PPF3 (λ_{excit} 370 nm).