

Electronic Supplementary Information

A sacrificial PLA block mediated route to injectable and degradable PNIPAAm based hydrogels

Vernon Tebong Mbah^{1,2}, Vincent Pertici¹, Pierluigi Stipa², Bernard Verrier³, Céline Lacroix³, Didier Gigmes¹, Thomas Trimaille^{1*}

¹ Aix Marseille Univ, CNRS, ICR, Marseille, France

² Dipartimento di Scienze dei Materiali e della Terra, Università Politecnica delle Marche, via Brecce Bianche, 60131 Ancona, Italy

³ Université Claude Bernard Lyon 1, Laboratoire de Biologie Tissulaire et d'Ingénierie Thérapeutique, UMR 5305, Université Lyon 1, CNRS, IBCP, 69367 Lyon, France




Figure S1. ¹H NMR spectrum of PLA-HEA (CDCl₃, 400 MHz), *: methanol




Figure S2. ¹H NMR spectrum of PLA-SG1 (CDCl₃, 400 MHz), *: methanol




Figure S3. ¹H NMR spectrum of PEGMA (CDCl₃, 400 MHz)




Figure S4. ¹H NMR spectrum of PLA-b-P(NIPAAm-co-PEGMA) (CDCl₃, 400 MHz)




Figure S5. ¹H NMR spectrum of P(NIPAAm-co-PEGMA) (CDCl₃, 400 MHz)