

Supporting Information

Association of Hydrophobic Carboxyl-Terminal Dendrimers with Lymph Node-Resident Lymphocytes

Yutaka Nishimoto ¹, Misaki Nishio ¹, Shu Nagashima ¹, Kohei Nakajima ², Takayuki Ohira ², Shinya Nakai ³, Ikuhiko Nakase ³, Kei Higashikawa ⁴, Yuji Kuge ⁴, Akikazu Matsumoto ¹, Mikako Ogawa ² and Chie Kojima ^{1*}

¹ Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, 1-1 Gakuen-cho, Naka-ku, Sakai, Osaka 599-8531, Japan

² Laboratory of Bioanalysis and Molecular Imaging, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan

³ Department of Biological Science, Graduate School of Science, Osaka Prefecture University, 1-2, Gakuen-cho, Naka-ku, Sakai, Osaka, 599-8570, Japan

⁴ Central Institutes of Isotope Science, Hokkaido University, Kita 15 Nishi 7, Kita-ku, Sapporo, Hokkaido 060-0815, Japan




Figure S1. ¹H NMR spectra of (a) C-den, (b) Chex-den, (c) C-Phe-den and (d) Chex-Phe-den.