

Supporting information for

Characterization of thermoresponsive poly-N-Vinylcaprolactam for biological applications

Lorenzo Marsili^{1,2,*}, Michele Dal Bo¹, Giorgio Eisele³, Ivano Donati⁴, Federico Berti² and Giuseppe Toffoli⁵

¹ CRO National Cancer Institute; Via Franco Gallini 2, 33081 Aviano, Italy, lorenzo.marsili@phd.units.it, mdalbo@cro.it, gtoffoli@cro.it

² Department of Chemical and Pharmaceutical Sciences, University of Trieste, Via Licio Giorgieri 1, 34127 Trieste, Italy; fberti@units.it

³ Ronzoni Institute; Via G. Colombo 81, 20133 Milano, Italy, eisele@cat-rozoni.it

⁴ Department of Life Sciences, University of Trieste, Via Licio Giorgieri 5, 34127 Trieste, Italy; idonati@units.it




Figure S1 ¹H spectrum of PNVCL_122.




Figure S2 Carboxylic termination group signals in ¹H NMR spectrum of PNVCL_305 dissolved in d₆-DMSO.




Figure S3 ¹³C spectrum of PNVCL_122.




Figure S4 HSQC-DEPT spectrum of PNVCL₁₂₂.




Figure S5 Comparison between NVCL and PNVCL-COOH spectrum (PNVCL₃₀₅).




Figure S6 Reduced viscosity reported as a function of PNVC_244 (black squares), PNVC_305 (red circles) and PNVC_1220 (blue triangles) concentration. The value of η_{red} were calculated according to the Huggins method by using the calibration lines reported in the figure.




Figure S7 Displacement of the correlation curve of PNVC_1220 solution (0.5 w%) associated to LCST transition.




Figure S8 Variation of UV-VIS spectrum of a solution of PNVCL_244 in human plasma (5mg/ml) with temperature.




Figure S9 Variation of UV-VIS spectrum of a solution of PNVCL_244 in human plasma (5mg/ml) with temperature.




Figure S10 Conductimetric titration of PNVCL_305.