

Supplementary Information

Isolation of *Microbulbifer* sp. SOL66 with High Polyhydroxyalkanoate-Degrading Activity from the Marine Environment

Sol Lee Park¹, Jang Yeon Cho¹, Soo Hyun Kim¹, Shashi Kant Bhatia^{1,2}, Ranjit Gurav¹, See-Hyoung Park³, Kyungmoon Park³ and Yung-Hun Yang^{1,2,*}

¹ Department of Biological Engineering, College of Engineering, Konkuk University, Seoul 05029, Korea; shckd20206@naver.com (S.L.P.); whwkddus1123@gmail.com (J.Y.C.); gsm06136@naver.com (S.H.K.); shashibiotechhpu@gmail.com (S.K.B.); rnjtgurav@gmail.com (R.G.)

² Institute for Ubiquitous Information Technology and Applications, Konkuk University, Seoul 05029, Korea

³ Department of Biological and Chemical Engineering, Hongik University, Sejong 30016, Korea; shpark74@hongik.ac.kr (S.-H.P.); pkm2510@hongik.ac.kr (K.P.)

* Correspondence: seokor@konkuk.ac.kr; Tel.: +82-2-450-3936

Figure S1: Surface change of the PHB pellet after 18 days cultivation.




Figure S2: GC-MS data of the supernatant of culture medium of *Microbulbifer* sp. SOL66 with P(3HB-*co*-4HB).

