

Experimental Design and Optimization of Nano-Transfersomal Gel to Enhance the Hypoglycemic Activity of Silymarin

Marwa H. Abdallah ^{1,2,*}, Amr S. Abu Lila ², Seham Mohammed Shawky ³, Khaled Almansour ¹, Farhan Alshammari ¹, El-Sayed Khafagy ^{4,5} and Tarek Saad Makram ⁶

¹ Department of Pharmaceutics, College of Pharmacy, University of Ha'il, Ha'il 81442, Saudi Arabia;

kh.almansour@uoh.edu.sa (K.A.); frh.alshammari@uoh.edu.sa (F.A.)

² Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Zagazig University,

Zagazig 44519, Egypt; a.abulila@uoh.edu.sa (A.S.A)

³ Department of Pharmaceutics and Pharmaceutical Technology, Faculty of Pharmacy, Al-Azhar University, Cairo 11651, Egypt; sehamshawky@azhar.edu.eg (S.M.S)

⁴ Department of Pharmaceutics, College of Pharmacy, Prince Sattam Bin Abdulaziz University,

Al-Kharj 11942, Saudi Arabia; e.khafagy@psau.edu.sa (E.-S.K)

⁵ Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Suez Canal University, Ismailia 41552, Egypt

⁶ Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, October 6 University,

Al Mehwar Al Markazi 12511, Egypt; tareksaadmakram@yahoo.com (T.S.M)

* Correspondence: mh.abdallah@uoh.edu.sa (M.H.A)




Figure S1. Schematic diagram of locally fabricated diffusion cell.