

Correction

Correction: Yang et al. Construction and Evaluation of Chitosan-Based Nanoparticles for Oral Administration of Exenatide in Type 2 Diabetic Rats. *Polymers* 2022, 14, 2181

Jian-Miao Yang ^{1,2,†}, Lin-Jie Wu ^{3,†}, Meng-Ting Lin ³, Yi-Ying Lu ³, Tian-Tian Wang ^{1,3}, Min Han ^{3,*}, Bin Zhang ^{4,*} and Dong-Hang Xu ^{1,*} 

- ¹ Department of Pharmacy, The Second Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou 310009, China; yangjm@enzemed.com (J.-M.Y.); 15574881635@163.com (T.-T.W.)
² Taizhou Hospital of Zhejiang Province, Zhejiang University, Taizhou 317099, China
³ Institute of Pharmaceutics, College of Pharmaceutical Sciences, Zhejiang University, Hangzhou 310058, China; 22119114@zju.edu.cn (L.-J.W.); 15700060560@163.com (M.-T.L.); 21919089@zju.edu.cn (Y.-Y.L.)
⁴ Zhejiang Strong Pharmaceutical Co., Ltd., Hangzhou 311500, China
* Correspondence: hanmin@zju.edu.cn (M.H.); zhangbin1@cnstrong.cn (B.Z.); xudonghang@zju.edu.cn (D.-H.X.); Tel.: +86-571-88208437 (M.H.); +86-571-87784529 (D.-H.X.)
† These authors contributed equally to this work.

The authors wish to make the following corrections to this paper: [1].

There was a mistake in Figure 3 as published. Figure 3c must be substituted with the following one. The corrected Figure 3 is presented below.



Citation: Yang, J.-M.; Wu, L.-J.; Lin, M.-T.; Lu, Y.-Y.; Wang, T.-T.; Han, M.; Zhang, B.; Xu, D.-H. Correction: Yang et al. Construction and Evaluation of Chitosan-Based Nanoparticles for Oral Administration of Exenatide in Type 2 Diabetic Rats. *Polymers* 2022, 14, 2181. *Polymers* 2023, 15, 2852. <https://doi.org/10.3390/polym15132852>

Received: 21 June 2023
Accepted: 25 June 2023
Published: 28 June 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 3. (a,b) The cytotoxicity of CS-TPP-ALG NPs, CS-TPP-ALG carrier, and exenatide in Caco-2 cells after 24 h (a) and 72 h (b). (c) Intracellular distribution of Rho-CS-TPP-ALG NPs in Caco-2 cells. Cells were exposed to Rhodamine B, Rho-CS, and Rho-CS-TPP-ALG NPs at 37 °C for 4 h, the concentration of fluorescence was 1 µg/mL.

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Reference

1. Yang, J.-M.; Wu, L.-J.; Lin, M.-T.; Lu, Y.-Y.; Wang, T.-T.; Han, M.; Zhang, B.; Xu, D.-H. Construction and Evaluation of Chitosan-Based Nanoparticles for Oral Administration of Exenatide in Type 2 Diabetic Rats. *Polymers* **2022**, *14*, 2181. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.