

Supplementary Material for Article

Effect of Different Cover Crops, Mass-Trapping Systems and Environmental Factors on Invertebrate Activity in Table Olive Orchards—Results from Field Experiments in Crete, Greece

Nikolaos Volakakis ^{1,2,3,*}, Emmanouil Kabourakis ¹, Leonidas Rempelos ^{3,4}, Apostolos Kiritsakis ⁵
and Carlo Leifert ^{6,7,*}

¹ Olive and Agroecological Production Systems Lab (EOPS), Hellenic Mediterranean University, 714 10 Iraklio, Greece

² Geokomi Plc, Sivas Festos, P.O. Box 21, 70200 Crete, Greece

³ Nafferton Ecological Farming Group, Newcastle University, Newcastle upon Tyne NE1 7RU, UK


⁴ Lincoln Institute for Agri-Food Technology, University of Lincoln, Riseholme Park, Lincoln LN2 2LG, UK

⁵ International Observatory of Oxidative stress for Health and Agricultural Products, 21 Kar. Diehl Str, 54623 Thessaloniki, Greece


⁶ SCU Plant Science, Southern Cross University, Military Rd., Lismore, NSW 2480, Australia

⁷ Department of Nutrition, IMB, University of Oslo, 0372 Oslo, Norway

* Correspondence: nvolakakis@gmail.com (N.V.); carlo.leifert@gmail.com (C.L.)



Supplementary Figure S1. Dimensions and arrangement of olive trees in an individual cover crop plot; each plot had 12 olive trees and a size of 432 m². , location of individual olive trees.



Supplementary Figure S2. Average mean daily temperature and total rainfall per months in the three cover-crop growing seasons monitored. P1, P2 and P3 indicate planting dates for cover crops; A1 and A2 indicate the dates on which invertebrate activity assessments were carried out in 2007 and 2008; H1 and H2 indicate the two olive harvests in 2006 and 2008.