

Table S1. Sequence of primers used in the qRT-PCR amplifications.

Gene name	Primer name	Primer sequence
<i>Pb4CL2</i>	<i>Pb4CL2-F</i>	AGTGTGGGTCAGGGTTATG
	<i>Pb4CL2-R</i>	AATCTCGCCTGGTTGGTTAT
<i>PbCAD</i>	<i>PbCAD -F</i>	TAGTTGGTGTAGTGACAGAG
	<i>PbCAD -R</i>	TGGAATGCGGACTATGAA
<i>PbPOD7</i>	<i>PbPOD7-F</i>	CGTGGTGTTCATGTGCTGATA
	<i>PbPOD7-R</i>	AACCAAGTCTCTGGTGGAAAG
<i>PbPOD17</i>	<i>PbPOD17-F</i>	CAACTCCGCCACGGTTATTA
	<i>PbPOD17-R</i>	ACCTTCTCCAAGGCTTCTTC
<i>PbCAD1</i>	<i>PbCAD1-F</i>	CTTCTCTACCAACCCATCTTC
	<i>PbCAD1-R</i>	CTTCTCTACCAACCCATCTTC
<i>PbCAD1-1</i>	<i>PbCAD1-1-F</i>	ACTCGAGCCGTACCTATCTT
	<i>PbCAD1-1-R</i>	ACCTCTCGAAAGCTGTGTTTAT
<i>Pb4CL11</i>	<i>Pb4CL11-F</i>	TTGGACTCCGATGGGTATTTG
	<i>Pb4CL11-R</i>	GAGCCTTCTCTGGGATTATGG