

## Article

# Effects of Pruning on Tea Tree Growth, Soil Enzyme Activity and Microbial Diversity

Qi Zhang <sup>1</sup>, Ying Zhang <sup>1,2</sup>, Pengyao Miao <sup>1,3</sup>, Meihui Chen <sup>1,2</sup>, Mengru Du <sup>1</sup>, Xiaomin Pang <sup>4</sup>, Jianghua Ye <sup>1</sup>, Haibin Wang <sup>1,\*</sup> and Xiaoli Jia <sup>5,\*</sup>

<sup>1</sup> College of Tea and Food Science, Wuyi University, Wuyishan City 354300, China; zhangqi1113@wuyiu.edu.cn (Q.Z.)

<sup>2</sup> College of Horticulture, Fujian Agriculture and Forestry University, Fuzhou 350002, China

<sup>3</sup> College of Resources and Environment, Fujian Agriculture and Forestry University, Fuzhou 350002, China

<sup>4</sup> Information Technology and Laboratory Management Center, Wuyi University, Wuyishan City 354300, China

<sup>5</sup> College of Ecology and Environmental Engineering, Wuyi University, Wuyishan City 354300, China

\* Correspondence: w13599084845@sina.com (H.W.); jiaxl2010@wuyiu.edu.cn (X.J.)

**Abstract:** In order to investigate the effect of pruning on the soil environment in which tea trees grow and the growth of tea trees, this study used Wuyi Meizhan (*Camellia sinensis*) as a research object and measured its growth indexes, soil physicochemical indexes, soil enzyme activity and microbial functional diversity to analyze the effects of pruning treatments on the growth of tea trees, soil enzyme activity and soil microbial functional diversity and the correlation between them. The results of the analysis of tea tree growth indexes showed that the hundred-bud weight, leaf area and yield in the pruning treatment were significantly higher than those in the unpruned treatment. The results of soil physicochemical index analysis showed that pH, available phosphorus, available potassium and organic matter were significantly higher in the pruning treatment than in the unpruned treatment ( $p < 0.05$ ), while available nitrogen and total phosphorus were significantly lower than in unpruned treatment ( $p < 0.05$ ). The results of soil enzyme activities showed that only polyphenol oxidase and catalase activities were significantly higher in the pruning than in the unpruned treatment, while urease, protease, acid phosphatase, asparaginase and glutaminase activities were significantly lower than in the unpruned treatment ( $p < 0.05$ ). Biolog analysis showed that the utilization of microbial carbon sources, especially amino acid and amine, increased in the rhizosphere soil of the pruned tea tree, while there was a significant decrease ( $p < 0.05$ ) in microbial diversity. It is evident that pruning promoted tea tree growth and some enzyme activity, while inhibiting the activity of enzymes associated with the nitrogen cycle, and the utilization of microbial carbon sources increased, but their diversity decreased. This study provides a theoretical basis for the daily management of tea plantation after pruning.

**Keywords:** pruning; Wuyi Meizhan; soil enzyme activity; growth index; microbial diversity; Biolog



**Citation:** Zhang, Q.; Zhang, Y.; Miao, P.; Chen, M.; Du, M.; Pang, X.; Ye, J.; Wang, H.; Jia, X. Effects of Pruning on Tea Tree Growth, Soil Enzyme Activity and Microbial Diversity. *Agronomy* **2023**, *13*, 1214. <https://doi.org/10.3390/agronomy13051214>

Academic Editors: Gianpiero Vigani, Maurizio Badiani and Georgia Ntatsi

Received: 30 March 2023

Revised: 20 April 2023

Accepted: 24 April 2023

Published: 25 April 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Tea tree pruning is an important management method in the process of tea tree planting. Tea tree pruning can stimulate the growth of lateral buds, and the increase in lateral branches, which in turn can be beneficial for the increase in tea yield. At the same time, tea tree pruning could reduce the labor force and improve the efficiency of tea harvesting [1–3]. In the process of tea tree growth, light pruning generally consists of cutting off 3–5 cm of the green leaf layer on the crown surface of a tea tree, which can promote bud tip germination and reduce leaf pinching; heavy pruning consists of pruning branches with aging trees, more dead branches, diseases and pests, and weak bud-rearing ability, and to cut 1/3–1/2 of the original tree height according to the degree of aging; generally, the older the trees, the more they are cut off [4,5]. Regardless of the degree of pruning, pruned tea trees need to be subjected to greater physical stress than

pruning tea trees due to the large loss of plant biomass [6]. Relevant research showed that pruned treatment led to changes in root secretions of tea trees [7]. Therefore, this study hypothesized that after being subjected to pruning stress, tea trees would produce a series of responses to adapt to pruning stress, and these defense mechanisms could be transmitted to the soil at the roots of tea trees.

Soil enzymes are important biocatalysts in the soil and are involved in all material circulation in the soil [8]. Relevant research showed that the degradation of organic matter and animal and plant residues in soil required the involvement of a large number of soil enzymes, including soil urease, sucrase, acid phosphatase, protease and catalase [9,10]. The pruning of tea trees produced large amounts of litter, the degradation of which required the involvement of a large number of enzymes. At the same time, tea litter led to the accumulation of allelochemicals such as polyphenols, flavonoids, and alkaloids in the soil, so the entire tea plantation contained a large number of secondary metabolites, which in turn affected soil health, soil microbial activity, and biogeochemical cycles [11].

Soil microorganisms play an important role in soil nutrient metabolism and affect soil physicochemical properties [12–14]. Relevant studies have shown that root secretions induced microorganisms to constantly change and adapt to the changing root soil environment [15]. The plant-soil feedback mechanism affected the composition of root secretions and metabolites in the soil, which in turn regulated the structure and function of the soil microbial community. Relevant research showed that soil microorganisms played an important role in promoting tea tree growth, improving tea quality, and protecting tea trees from pathogens [16]. Li et al. [17] showed that soil bacterial diversity tended to decrease with increasing years of tea planting, especially decreasing the relative abundance of some beneficial bacteria. However, there are few reports of changes in rhizosphere soil enzymes and soil microbial communities of perennial plants such as tea trees affected by physical stress from pruning [18,19]. In this study, the effects of pruning on tea tree growth index, soil physicochemical index, soil enzyme activity, and microbial functional diversity were analyzed with Wuyi rock tea Meizhan as the research object, aiming to reveal the effects of pruning stress on the soil environment of the root system of tea trees and provide a theoretical basis for the healthy management of tea plantations.

## 2. Materials and Methods

### 2.1. Experimental Site and Soil Sample Collection

The experimental site was located in the area of Foguo Yan in the Wuyi Mountain Scenic Area (117.99° E, 27.72° N) at an altitude of 239 m (Figure 1). The experimental tea cultivar was Meizhan (*Camellia sinensis* cv. Meizhan), with a planting area of about 0.7 ha. The tea plants were nine years old and had not been pruned before. In August 2021, the test area was evenly divided into six areas, three of which had tea trees pruned for treatment (MX), i.e., three replicates, and the other three areas had tea trees unpruned, i.e., controls (MC). Tea trees were pruned by trimming 3–5 cm from the tea tree canopy, and pruning litter was left on the surface of the original tea tree soil. Each year in late October, 700 kg/ha of compound fertilizer (N:P:K = 21:8:16) was applied to tea plantations in each treatment. Other tea plantation management measures, such as weeding and watering, were the same for both pruned and unpruned treatments.

### 2.2. Determination of the Growth Index of Tea Trees

Growth indexes were measured on 27 April 2022. Leaf area, hundred-bud weight, chlorophyll content, nitrogen content of tea leaves, and yield of pruned and unpruned treatments were measured [20]. Leaf area: 20 mature shoots were randomly selected to measure the leaf length and width, and leaf area was calculated based on length × width × 0.7. One hundred (100)-bud weight: 100 standard bud tips with three leaves were randomly selected, weighed and replicated three times. Chlorophyll and nitrogen content: chlorophyll and leaf nitrogen content of leaves were determined by chlorophyll meter (TYS-N), and the second functional leaf of new shoots was repeated eight times. Tea yields: three rows of tea

trees were randomly selected to harvest fresh leaves, respectively [20]. Three leaves of standard standing buds were picked and weighed. Tea yield = tea weight per row/area per row.



**Figure 1.** Tea plantations of Meizhan (*Camellia sinensis* cv. Meizhan). The upper part shows unpruned treatment and the lower part shows pruned treatment.

### 2.3. Soil Sample Collection and Determination of Physicochemical Indexes

Soil sampling was conducted on 27 April 2022. Equidistant soil sampling was used to divide the strip planting area into five equal divisions. Five tea trees from each equal division center were taken as one sample, and a total of three soil samples were taken [20]. The roots were dug up near the roots of the tea bushes, and the fine roots were collected at a depth of 5–20 cm of soil. Rhizosphere soil on fine roots was collected by shaking off. Soil samples collected from the pruned region were denoted MX and those collected from the unpruned region were denoted MC. Fresh soil samples were used for enzymatic and microbial functional diversity analyses, while air-dried soil samples were analyzed for soil physicochemical indexes.

Part of the air-dried soil was taken, crushed, screened by 2 mm, and sampled according to the quartering method. Soil pH, total nitrogen (TN), total phosphorus (TP), total potassium (TK), available nitrogen (AN), available phosphorus (AP), available potassium (AK), and organic matter (OM) were determined using the Soil Agricultural Chemical Analysis method [21]. Analyses were repeated three times.

### 2.4. Determination of Soil Enzyme Activity

The soil enzymes measured in this study mainly include urease, sucrase, polyphenol oxidase, catalase, acid phosphatase protease, asparaginase, glutaminase, and nitrate reductase. The determination methods all refer to the experimental manual of ‘Soil and environmental microbiology research method’ [22]. Briefly, urease activity was estimated colorimetrically by sodium phenol-sodium hypochlorite. Sucrase activity was determined colorimetrically by nitrosalicylic acid. Polyphenol oxidase activity was determined colorimetrically by pyrogallol. Catalase activity was determined by titration of potassium permanganate. Acid phosphatase activity was determined colorimetrically by disodium phenyl phosphate. The protease activity was determined colorimetrically by ninhydrin. The activities of asparaginase and glutaminase were determined colorimetrically by Nye’s reagent. Nitrate reductase activity was determined colorimetrically by phenol disulfonic acid.



### 2.5. Determination of Functional Diversity of Microorganisms

Soil microbial functional diversity was determined by BIOLOG ECO microplate method [23]. Briefly, 10 g of fresh soil was taken, placed in a 250 mL sterilized conical bottle, filled with 90 mL sterilized normal saline, sealed, shaken at 120 r/min for 10 min and left to stand for 2 min. In an ultra-clean workbench, 5 mL of supernatant was taken and put into 45 mL of sterile water to obtain a 1:100 dilution; the dilution was repeated once to obtain 1:1000 extract to be set aside. One hundred and fifty (150)  $\mu\text{L}$  of the diluted soil suspension was transferred to each well, and 150  $\mu\text{L}$  of sterile water was added to wells A1, A5 and A9 as control. After scanning the absorbance at 590 nm on a microplate reader (Nanjing Detie Experimental Equipment Co., Ltd., Nanjin, Jiangsu province, China) (recorded as the absorbance of 0 h), incubation was carried out in the dark at 28 °C. The absorbance at 590 nm of each well was then scanned at regular intervals each day for 7 days. Average well color development (AWCD) in BIOLOG ECO plate was used to express the results of ELISA reaction of soil microbial communities.  $\text{AWCD} = [\sum(C - R)]/31$ , where C is the absorbance of 31 carbon source wells and R is the absorbance of the control well. According to the classification of 31 single carbon sources in BIOLOG ECO microplate, carbon sources are divided into six categories: sugar, amino acids, carboxylic acids, amines, polymers and phenols.

### 2.6. Data Analysis

Excel 2017 software was used for data sorting; IBM SPSS Statistics 21.0 software (Chicago, IL, USA) was used for data T-test and correlation analysis; Excel 2017 and Rstudio 3.3 software (Boston, MA, USA) were used for OPLS-DA simulation, box diagram, heat map and redundancy analysis.

## 3. Results

### 3.1. Analysis of Tea Tree Growth Indexes

The results of pruning on tea tree growth index showed (Figure 2) that there was no significant difference between MC and MX in chlorophyll content and leaf nitrogen content. The leaf area and yield of MX were significantly higher than MC ( $p < 0.01$ ), which were 1.32 and 1.84 times higher, respectively. The hundred-bud weight of MX was significantly higher than that of MC ( $p < 0.05$ ), which was 1.40 times higher. Overall, pruning resulted in more vigorous growth and higher tea yields.


### 3.2. Analysis of Soil Physicochemical Index

The contents of AN and TP in MX were significantly lower than those in MC ( $p < 0.05$ ), by 6.15% and 29.13%, respectively (Table 1). After pruning, the pH, AP, AK and OM contents in tea soil were significantly higher than those in MC ( $p < 0.05$ ), by 10.94%, 17.22%, 9.74% and 5.65%, respectively. The content of TN and TK in MX was not significantly different from MC. This suggests that pruning could affect the available nutrient cycle in the rhizosphere soil of tea trees, especially N and P.

**Table 1.** Effect of pruning on soil physicochemical indexes of tea trees ( $n = 3$ ).

Index	MC	MX	Index	MC	MX
pH	4.02 $\pm$ 0.01 <sup>b</sup>	4.46 $\pm$ 0.01 <sup>a</sup>	TN (g/kg)	1.15 $\pm$ 0.02 <sup>a</sup>	1.10 $\pm$ 0.04 <sup>a</sup>
AN (mg/kg)	122.10 $\pm$ 1.03 <sup>a</sup>	114.59 $\pm$ 1.82 <sup>b</sup>	TP (g/kg)	1.03 $\pm$ 0.03 <sup>a</sup>	0.73 $\pm$ 0.02 <sup>b</sup>
AP (mg/kg)	13.24 $\pm$ 0.48 <sup>b</sup>	15.52 $\pm$ 0.62 <sup>a</sup>	TK (g/kg)	5.62 $\pm$ 0.33 <sup>a</sup>	5.84 $\pm$ 0.47 <sup>a</sup>
AK (mg/kg)	101.41 $\pm$ 2.81 <sup>b</sup>	111.29 $\pm$ 1.10 <sup>a</sup>	OM (g/kg)	20.01 $\pm$ 0.02 <sup>b</sup>	21.14 $\pm$ 0.09 <sup>a</sup>

Note: AN: available nitrogen, AP: available phosphorus, AK: available potassium, TN: total nitrogen, TP: total phosphorus; TK: total potassium, OM: organic matter. MC: Meizhan unpruned, MX: Meizhan pruned, lowercase letter represents the significant difference at the 0.05 level.




**Figure 2.** Effect of pruning on growth indexes of Meizhan tea trees. MC: Meizhan unpruned; MX: Meizhan pruned; the dotted line is average value; the green delineation shows the range of values; \*\*: significant difference at 0.01 level; ns: not significant. (A) Effect of pruning treatment on tea leaf area; (B) effect of pruning treatment on tea leaf nitrogen content; (C) effect of pruning treatment on tea hundred-bud weight; (D) effect of pruning treatment on tea Chlorophyll II content; (E) effect of pruning treatment on tea yield.

### 3.3. Analysis of Soil Enzyme Activity


Soil enzymes are important for nutrient cycle in soil. The analysis of soil enzyme activity showed (Figure 3) that the activities of polyphenol oxidase and catalase were significantly higher in MX than in MC ( $p < 0.05$  or  $p < 0.01$ ), 1.29 and 1.43 times higher, respectively; the activities of acid phosphatase, protease, urease, asparaginase and glutaminase were significantly higher in MC than in MX ( $p < 0.05$  or  $p < 0.01$ ), by 1.49, 1.05, 2.47, 1.10 and 1.11 times, respectively. This revealed that pruning could affect the activity of some enzymes in rhizosphere soil of tea trees, particularly those related to the nitrogen cycle.

### 3.4. Correlation Analysis

The correlation results showed that tea tree growth indexes were significantly and positively correlated with soil physicochemical indexes, except for available nitrogen (Figure 4). Tea tree growth indexes were positively correlated with polyphenol oxidase, catalase and nitrate reductase, and negatively correlated with other enzymes. With the exception of total nitrogen and available nitrogen, soil physical and chemical indexes were positively correlated with polyphenol oxidase, catalase and nitrate reductase, but negatively correlated with other enzymes. Total nitrogen and available nitrogen were negatively correlated with polyphenol oxidase, catalase and nitrate reductase, but positively correlated with other enzymes.



**Figure 3.** Effect of pruning treatment on soil enzyme activity of Meizhan tea trees ( $n = 3$ ). MC: Meizhan unpruned; MX: Meizhan pruned; \*: significant difference at 0.05 level; \*\*: significant difference at 0.01 level; ns: not significant. (A) Effect of pruning treatment on acid phosphatase in rhizosphere soil of tea trees; (B) Effect of pruning treatment on asparaginase in rhizosphere soil of tea trees; (C) Effect of pruning treatment on catalase in rhizosphere soil of tea trees; (D) Effect of pruning treatment on glutaminase in rhizosphere soil of tea trees; (E) Effect of pruning treatment on nitrate reductase in rhizosphere soil of tea trees; (F) Effect of pruning treatment on polyphenol oxidase in rhizosphere soil of tea trees; (G) Effect of pruning treatment on protease in rhizosphere soil of tea trees; (H) Effect of pruning treatment on sucrase in rhizosphere soil of tea trees; (I) Effect of pruning treatment on urease in rhizosphere soil of tea trees.




**Figure 4.** Correlation analysis between growth index, soil physicochemical index and soil enzyme activity. LA: leaf area, CC: chlorophyll content, LNC: leaf nitrogen content, HBW: hundred-bud weight, YI: yields, TN: total nitrogen, TP: total phosphorus, TK: total potassium, AN: available nitrogen, AP: available phosphorus, AK: available potassium, OM: organic matter, POL: polyphenol oxidase, CAT: catalase, ACP: acid phosphatase, SUC: sucrase, PRO: protease, URE: urease, NIT: nitrate reductase, ASP: asparaginase, GLU: glutaminase. Red is negatively correlated and blue is positively correlated. The color area is the correlation coefficient.

### 3.5. Analysis of Microbial Functional Diversity in Rhizosphere Soil Microorganisms of Tea Trees


The results of carbon source metabolism characteristics analysis of rhizosphere soil microorganisms of tea trees showed (Figure 5) that the degree of carbon source utilization by microorganisms in tea rhizosphere soil increased with the increase in culture time. From 24 h onwards, microbial carbon source utilization in tea rhizosphere soil was higher in MX than in MC.

The results of the utilization degree of six carbon sources categories showed that (Figure 6) the utilization degree of amino acids and amines by soil microorganisms was significantly higher in MX than in MC ( $p < 0.01$ ), by 1.64 and 2.15 times, respectively, whereas the utilization of carboxylic acids, carbohydrates, polymers and phenols did not differ significantly.

The diversity analysis showed (Table 2) that the Simpson diversity index of soil microorganisms was significantly lower in MX than in MC ( $p < 0.05$ ). However, the Shannon diversity index and Chao1 in MX were not significantly different from MC.



**Figure 5.** Changes in AWCD values of carbon sources utilized by soil microorganisms. MC: Meizhan unpruned; MX: Meizhan pruned.



**Figure 6.** Utilization rate of carbon sources by soil microorganisms. MC: Meizhan unpruned; MX: Meizhan pruned; \*\*: significant difference at 0.01 level; ns: not significant; the dotted line is average value; the green delineation shows the range of values; (A) Utilization rate of amino acids by soil microorganisms; (B) Utilization rate of amines by soil microorganisms; (C) Utilization rate of polymers by soil microorganisms; (D) Utilization rate of phenols by soil microorganisms; (E) Utilization rate of carboxylic acids by soil microorganisms; (F) Utilization rate of carbohydrates by soil microorganisms.

**Table 2.** Microbial diversity index in rhizosphere soil of tea tree.


Treatment	Simpson Index	Shannon Index	Chao 1
MC	0.999 ± 0.001 <sup>a</sup>	4.525 ± 0.009 <sup>a</sup>	0.913 ± 0.002 <sup>a</sup>
MX	0.983 ± 0.001 <sup>b</sup>	4.510 ± 0.028 <sup>a</sup>	0.910 ± 0.006 <sup>a</sup>

Note: MC: Meizhan unpruned; MX: Meizhan pruned; lowercase letter represents a significant difference at 0.05 level.



### 3.6. Carbon Source Screening for Microbial Differences in Rhizosphere Soil of Tea Trees

OPLS-DA can be used to model the correlation between carbon source utilization and sample type, and to screen key carbon sources that can characterize sample differences by variable importance projection value (VIP value). Meanwhile, in order to test the reliability of the OPLS-DA model, permutation testing was usually used to verify the model, so as to evaluate the accuracy of the model. The results of the analysis of the OPLS-DA model showed that the goodness of fit  $R^2Y$  value of the OPLS-DA model for pruned and unpruned samples was 0.999 ( $p < 0.01$ ), and the predictive  $Q^2$  value for predictability was 0.892 ( $p < 0.01$ ) (Figure 7A). It can be seen that both  $R^2Y$  and  $Q^2$  values of the model reached significant levels, and the fit of the model was good and credible for further analysis. The results of the OPLS-DA score map showed (Figure 7B) that OPLS-DA could effectively distinguish pruned and unpruned samples in different areas. In conclusion, there were significant differences in the utilization degree of carbon sources by soil microorganisms between pruned and unpruned treatments. The S-plot analysis showed (Figure 7C,D) that 15 key carbon sources (VIP > 1) distinguished MC and MX samples, among which pruned treatment increased the utilization of 13 carbon sources, and both amino acids and carboxylic acids accounted for more than 30% of the 15 key carbon sources.




**Figure 7.** OPLS-DA model and S-plot of soil microorganisms after tea tree pruning. (A) OPLS-DA models of the fitting degree test of MX and MC; (B) analysis of carbon source utilization of MX and MC

by OPLS–DA model; (C) OPLS–DA loading diagram for carbon source utilization of MX and MC, the red dots represent different substances of MX, and the green dots represent different substances of MC,  $p[1]$  represents the cocorrelation coefficient between the principal component and the index,  $p(\text{corr})[1]$  represents the correlation coefficient between the principal component and the index; (D) heat maps for carbon source utilization of MX and MC, the red is up-regulated and the green is down-regulated; the darker the color, the greater the utilization rate.

### 3.7. Redundancy Analysis

The results of redundancy analysis (RDA) showed (Figure 8) that differences in soil microorganisms' use of carbon sources in rhizosphere soil of pruned and unpruned tea trees could be well distinguished using the AWCD 72 h values. RDA analysis of Biolog data and growth indicators showed (Figure 8A) that RDA1 could explain 63.16% of the variation and RDA2 could explain 23.81% of the variation. The carbon source with the highest positive correlation coefficient with RDA1 was  $\beta$ -methyl-D-glucoside, and the carbon source with the highest negative correlation coefficient with principal component 1 was itaconic acid. Further correlation analysis showed that the growth indexes were positively correlated with MX. RDA results of Biolog data and soil enzyme activities showed (Figure 8B) that RDA1 could explain 65.76% of the variation, and RDA2 could explain 22.79% of the variation. The carbon source with the highest positive correlation coefficient with RDA1 was  $\beta$ -methyl-D-glucoside and the carbon source with the highest negative correlation coefficient with RDA1 was itaconic acid. Further correlation analysis showed that polyphenol oxidase, catalase and nitrate reductase were positively correlated with MX, while protease, urease, sucrase, acid phosphatase, asparaginase and glutaminase were positively correlated with MC.



**Figure 8.** BIOLOG cluster and redundancy analysis of soil microorganisms of tea trees. (A) redundancy analysis of BIOLOG data and growth indices; (B) redundancy analysis of BIOLOG data and soil enzyme activity; MC: Meizhan unpruned; MX: Meizhan pruned; LA: leaf area, CC: chlorophyll content, LNC: leaf nitrogen content, HBW: hundred-bud weight, YI: yields, POL: polyphenol oxidase, CAT: catalase, ACP: acid phosphatase, SUC: sucrase, PRO: protease, URE: urease, NIT: nitrate reductase, ASP: asparaginase, GLU: glutaminase; 1, 2, 3: the mean with replicate experiments; the dotted line show the horizontal and vertical axes; the red lines show environmental factors.

## 4. Discussion

Pruning is a cultivation management method that is usually performed as a routine step in tea cultivation to maintain the height of the leaf-picking table, increase the branches of tea trees, and improve tea productivity [24,25]. Zhang et al. [3] reported that summer pruning improved branch growth and tea quality. Yin et al. [26] reported that light pruning after spring could improve the yield and quality of the following spring tea. Lu et al. [5]

reported that heavy pruning of tea trees could increase hundred-buds weight and thicker and shorter stems of tea leaves. The results of this study showed (Figure 2) that the hundred-bud weight, leaf area and yield of pruned treatment were significantly higher than those of unpruned treatment, but there were no significant differences in chlorophyll content and leaf nitrogen content. This suggested that tea yield increased by increasing leaf area after pruning.

Plant growth depends on the presence and availability of water and nutrients in the soil [7]. Agricultural management measures and other factors, including abiotic conditions (pH, light, temperature) and biological conditions (varieties, planting years) affected soil physicochemical indexes [27,28]. Tea trees absorb a large amount of nutrients from the environment, especially the soil, during growth. In this study, it was shown that pruning significantly increased soil pH (Table 1). After pruning, the absorption of nitrogen by tea trees increased, resulting in a significant decrease in the nitrogen content of rhizosphere soil. The tea tree was stimulated by exogenous factors after pruning, and absorbed mineral elements efficiently, resulting in increased yield. For example, Mg or Mn could promote the enhancement of photosynthesis in tea trees. Cao et al. [29] indicated that in low-yield and inefficient tea plantations, applying organic fertilizer to tea trees after pruning could improve tea quality. The results suggested that the application of nitrogen fertilizer should be increased after pruning, which could increase the yield of tea.

Soil enzyme activity is a characterization and indicator of its biological condition and fertility [30,31]. Changes in the soil environment can lead to differences in soil enzyme activity. Correlation results showed that polyphenol oxidase and catalase activities were positively correlated with pH, and total phosphorus, available phosphorus, available potassium, and organic matter contents (Figure 4). Pruning increased soil polyphenol oxidase activity and favored the degradation of soil acidic substance and the increase in soil pH. Sun et al. [32] showed that soil pH and nitrogen were the main factors affecting catalase activity. Therefore, the increase in catalase activity after pruning may be due to an increase in soil pH. In this study, tea trees were treated with fertilizer after pruning to provide sufficient nitrogen sources for their growth, when there were sufficient nutrients in the soil and it was the soil environment that tea trees needed to improve in order to facilitate their own growth. Further research found that the increased activity of rhizosphere soil polyphenol oxidase and catalase after pruning provided a good environment for nitrogen uptake by tea trees, ensuring their own growth.

In general, inadequate fertilizer quantity stimulated and increased soil enzyme activity and promoted soil nutrient cycle, while enzyme activity was relatively low when the fertilizer was adequate. In this study, urease, protease, glutaminase and asparaginase activities related to the nitrogen cycle were significantly lower in the pruned treatment than in the unpruned treatment (Figure 3). On the one hand, the reduction in soil available nitrogen content after pruning proved that tea trees have a high demand for nitrogen (Table 1), while the content of available nitrogen in the soil was sufficient for the growth of tea trees (up to the Chinese national standard for soil fertility level 1), therefore, the activity of enzymes related to nitrogen cycle in the soil did not need to be increased. On the other hand, Arafat et al. [7] showed that degradation of tea litter led to the accumulation of allelochemicals such as polyphenols, flavonoids and alkaloids in the soil, which might inhibit the activity of these enzymes, as shown by the decrease in enzyme activity after pruning.

Soil microorganisms are highly sensitive to changes in the surrounding environment, and are often considered biological indicators for assessing soil health [33]. Relevant research showed that organic compounds secreted by roots trigger soil microorganisms to constantly change their surroundings [15,34]. Pramanik et al. [18] showed that pruning increased the number of bacteria, fungi and actinomycetes in the rhizosphere soil of tea trees, but reduced the number of denitrifying bacteria. The results of this study showed that Simpson diversity index was significantly lower in pruning treatment than in unpruned treatment, and the efficiency of carbon source utilization by tea rhizosphere microorganisms was higher than that in unpruned treatment, indicating that pruning treatment reduced

the functional diversity of tea rhizosphere microorganisms, but enhanced the functions of specific species (Table 2, Figure 6). At the same time, litter degradation due to pruning increased amino acids and carboxylic acids of carbon sources in the soil, and promoted the growth of microorganisms that used the relevant carbon sources (Figure 7). The research of Bora et al. [35] showed that long-term pruning reduced the richness of microorganisms in the rhizosphere soil of tea trees, while soil polyphenols accumulation promoted the growth and abundance of some microorganisms. This may be the reason pruned treatment reduced rhizosphere microbial diversity, but the utilization rate of carbon source was higher than that of unpruned treatment.

## 5. Conclusions

In conclusion, the pruned treatment stimulated the growth of tea trees and increased tea yield. The activity of some enzymes in tea rhizosphere soil was inhibited by the effect of tea leaf litter, especially those related to nitrogen metabolism, and microbial diversity was reduced. In order to prevent the accumulation of allelochemicals in the soil and to promote the growth of core microorganisms, the carbon source utilization of soil microorganisms was higher in the pruned treatment than in the unpruned treatment. As tea trees grow vigorously after pruning, they required large amounts of soil nitrogen, but soil enzyme activities related to the nitrogen cycle were affected by allelopathic substances. Therefore, it is recommended to add nitrogen fertilizer after pruning to meet the growth needs of tea trees and increase tea yield. Microbial diversity in the soil is reduced. Tea trees promote the growth of core microorganisms in defense against the accumulation of chemosensitive substances in the soil, resulting in a higher utilization of carbon sources by soil microorganisms between the roots of tea trees in pruning treatment than in unpruned treatment. As a result of pruning, tea trees grow more strongly and therefore require large amounts of soil nitrogen, but soil enzyme activities associated with nitrogen cycling are affected by chemosensitive substances. It is recommended that nitrogen fertilization should be supplemented in time after pruning to meet the growth requirements of tea trees in order to improve tea production.

**Author Contributions:** Q.Z. performed the experiments, carried out most data analysis, and wrote the manuscript. Y.Z., P.M., M.C. and M.D. advised on the experiments and data analysis. J.Y. and X.P. project administration. X.J. and H.W. designed the entire experiment and corrected the manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Scientific Research Project of Introducing Talents, Wuyi University (YJ201909); Innovation and Entrepreneurship Training Program for College Students (S202210397033); Construction of first-class undergraduate specialty (tea science) in Fujian Province (SJZY2019004); Fujian Province Young and Middle-aged Teacher Education Research Project (Science and Technology) (JAT200664), Faculty and students co-creation team of Wuyi University (2021-SSTD-05); Special for science and technology commissioner of Nanping City (NP2021KTS05).

**Data Availability Statement:** All data generated or analyzed during this study are included in this published article.

**Conflicts of Interest:** The authors declare that there are no competing interest in the submission of this manuscript.

## References

1. Wijeratn, M.A. Planting and cultivation of tea. In *Global Tea Science: Current Status and Future Needs*, 1st ed.; Sharma, Gunasekare, K., Eds.; Burleigh Dodds Science Publishing: London, UK, 2018; p. 32.
2. Tian, R.Q.; Lv, R.Q.; Fang, L.; Shi, Y.Z. Study on the influence of different pruning methods on tree crown formation in tea plantation with high quality tea machine. *Chin. Tea* **2015**, *37*, 16–19.
3. Zhang, L.; Li, M.; Li, X.; Yan, P.; Zhang, L.; Han, W. Summer pruning improves the branch growth and tea quality of tea trees (*Camellia sinensis*). *Acta Physiol. Plant.* **2021**, *43*, 65. [[CrossRef](#)]
4. Mamun, M.; Hoque, M.M.; Ahmed, M. Effects of different pruning operations on the incidence of red spider mite of tea in bangladesh. *Bangl. J. Bot.* **2016**, *45*, 247–251.



5. Lu, L.; Luo, W.; Zheng, Y.; Jin, J.; Liu, R.; Lv, Y.; Ye, Y.; Ye, J. Effect of different pruning operations on the plant growth, phytohormones and transcriptome profiles of the following spring tea shoots. *Beverage Plant Res.* **2022**, *2*, 12. [\[CrossRef\]](#)
6. Kandiah, S. Studies on the physiology of pruning tea 1. turnover of resources in relation to pruning. *Tea Quart.* **1971**, *42*, 89–100.
7. Arkorful, E.; Hu, S.; Zou, Z.; Yu, Y.; Chen, X.; Li, X. Metabolomic Analyses Provide New Insights into Signaling Mechanisms for Nutrient Uptake by Lateral Roots of Pruned Tea Plant (*Camellia sinensis*). *J. Agric. Food Chem.* **2020**, *68*, 7890–7903. [\[CrossRef\]](#)
8. Sun, X.; Ye, Y.; Ma, Q.; Guan, Q.; Jones, D.L. Variation in enzyme activities involved in carbon and nitrogen cycling in rhizosphere and bulk soil after organic mulching. *Rhizosphere* **2021**, *19*, 100376. [\[CrossRef\]](#)
9. Salam, A.K.; Katayama, A.; Kimura, M. Activities of some soil enzymes in different land use systems after deforestation in hilly areas of West Lampung, South Sumatra, Indonesia. *Soil Sci. Plant Nutr.* **1998**, *44*, 93–103. [\[CrossRef\]](#)
10. Elkhilfi, Z.; Iftikhar, J.; Sarraf, M.; Ali, B.; Saleem, M.H.; Ibranshabib, I.; Bispo, M.D.; Meili, L.; Ercisli, S.; Kayabasi, E.T.; et al. Potential Role of Biochar on Capturing Soil Nutrients, Carbon Sequestration and Managing Environmental Challenges: A Review. *Sustainability* **2023**, *15*, 2527. [\[CrossRef\]](#)
11. Arafat, Y.; Din, I.U.; Tayyab, M.; Jiang, Y.; Chen, T.; Cai, Z.; Zhao, H.; Lin, X.; Lin, W.; Lin, S. Soil Sickness in Aged Tea Plantation Is Associated With a Shift in Microbial Communities as a Result of Plant Polyphenol Accumulation in the Tea Gardens. *Front. Plant Sci.* **2020**, *11*, 601. [\[CrossRef\]](#)
12. Nacke, H.; Thürmer, A.; Wollherr, A.; Will, C.; HoDac, L.; Herold, N.; Schöning, I.; Schrupf, M.; Daniel, R. Pyrosequencing-Based Assessment of Bacterial Community Structure Along Different Management Types in German Forest and Grassland Soils. *PLoS ONE* **2011**, *6*, e17000. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Romaniuk, R.; Giuffré, L.; Costantini, A.; Nannipieri, P. Assessment of soil microbial diversity measurements as indicators of soil functioning in organic and conventional horticulture systems. *Ecol. Indic.* **2011**, *11*, 1345–1353. [\[CrossRef\]](#)
14. Lan, Y.; Wang, S.; Zhang, H.; He, Y.; Jiang, C.; Ye, S. Intercropping and nitrogen enhance eucalyptus productivity through the positive interaction between soil fertility factors and bacterial communities along with the maintenance of soil enzyme activities. *Land Degrad. Dev.* **2023**. [\[CrossRef\]](#)
15. Huang, X.-F.; Chaparro, J.M.; Reardon, K.F.; Zhang, R.; Shen, Q.; Vivanco, J.M. Rhizosphere interactions: Root exudates, microbes, and microbial communities. *Botany* **2014**, *92*, 267–275. [\[CrossRef\]](#)
16. Bag, S.; Mondal, A.; Banik, A. Exploring tea (*Camellia sinensis*) microbiome: Insights into the functional characteristics and their impact on tea growth promotion. *Microbiol. Res.* **2022**, *254*, 126890. [\[CrossRef\]](#)
17. Li, Y.; Li, Z.; Li, Z.; Jiang, Y.; Weng, B.; Lin, W. Variations of rhizosphere bacterial communities in tea (*Camellia sinensis* L.) continuous cropping soil by high-throughput pyrosequencing approach. *J. Appl. Microbiol.* **2016**, *121*, 787–799. [\[CrossRef\]](#)
18. Pramanik, P.; Phukan, M.; Ghosh, S.; Goswami, A.J. Pruned tea bushes secrete more root exudates to influence microbiological properties in soil. *Arch. Agron. Soil Sci.* **2018**, *64*, 1172–1180. [\[CrossRef\]](#)
19. Jiang, Y.; Lin, X.; Khan, M.U.; Jiang, W.; Xu, Y.; Li, Z.; Lin, W. Tea pruning for the umbrella-shaped canopy can alleviate rhizosphere soil degradation and improve the ecosystem functioning of tea orchards. *Catena* **2023**, *222*, 106885. [\[CrossRef\]](#)
20. Ye, J.H.; Hu, W.W.; Zhang, Q.; Wang, H.B.; Jia, X.L.; He, H.B. Correlation between soil characteristics of tea plantations and the growth and fresh leaf quality of wuyi tea (*Camellia sinensis* cv. Shuixian). *Chin. J. Trop Crop* **2020**, *41*, 1838–1846.
21. Lu, K. *Methods of Soil Agricultural Chemical Analysis*; China Agricultural Science and Technology Press: Beijing, China, 2000.
22. Li, Z.G.; Luo, Y.M.; Teng, Y. *Soil and Environmental Microbiology Research Method*; Science Press: Beijing, China, 2008.
23. Choi, K.H.; Dobbs, F.C. Comparison of two kinds of Biolog microplates (GN and ECO) in their ability to distinguish among aquatic microbial communities. *J. Microbiol. Meth.* **1999**, *36*, 203–213. [\[CrossRef\]](#)
24. Barua, D.N. Seasonal Dormancy in Tea (*Camellia sinensis* L.). *Nature* **1969**, *224*, 514. [\[CrossRef\]](#)
25. Kumar, R.; Bisen, J.S.; Singh, M.; Bera, B. Effect of pruning and skiffing on growth and productivity of Darjeeling tea (*Camellia sinensis* L.). *Int. J. Tech. Res. Appl.* **2015**, *3*, 28–34.
26. Yin, L.Q.; Xiao, X.; Luo, Q.X.; Xia, L.F.; Yang, Q.; Liang, M.Z. Effect of light pruning in different period on tea yield and quality of yunnan big-leaf varieties. *Southwest China J. Agric. Sci.* **2019**, *32*, 1034–1038.
27. Tang, W.; Yang, H.; Wang, W.; Wang, C.; Pang, Y.; Chen, D.; Hu, X. Effects of Living Grass Mulch on Soil Properties and Assessment of Soil Quality in Chinese Apple Orchards: A Meta-Analysis. *Agronomy* **2022**, *12*, 1974. [\[CrossRef\]](#)
28. Meddich, A.; Oufdou, K.; Boutasknit, A.; Raklami, A.; Tahiri, A.; Ben-Laouane, R.; Ait-El-Mokhtar, M.; Anli, M.; Mitsui, T.; Wahbi, S.; et al. Use of Organic and Biological Fertilizers as Strategies to Improve Crop Biomass, Yields and Physicochemical Parameters of Soil. In *Nutrient Dynamics for Sustainable Crop Production*; Meena, R., Ed.; Springer: Singapore, 2020; pp. 247–288.
29. Cao, Y.; Cui, X.M.; Luo, X.Y. Effects of combine trimming with fertilizing on tea quality of yield and low performing tea garden. *J. Anhui Agric. Sci.* **2012**, *40*, 13311–13312.
30. Zuber, S.M.; Villamil, M.B. Meta-analysis approach to assess effect of tillage on microbial biomass and enzyme activities. *Soil Biol. Biochem.* **2016**, *97*, 176–187. [\[CrossRef\]](#)
31. Zielewicz, W.; Swędrzyński, A.; Dobrzyński, J.; Swędrzyńska, D.; Kulkova, I.; Wierzchowski, P.S.; Wróbel, B. Effect of Forage Plant Mixture and Biostimulants Application on the Yield, Changes of Botanical Composition, and Microbiological Soil Activity. *Agronomy* **2021**, *11*, 1786. [\[CrossRef\]](#)
32. Sun, H.; Zhang, J.F.; Hu, Y.; Wang, L.; Chen, G.C. Response of soil catalase to cover of different stands. *Bull. Soil Sci.* **2016**, *47*, 605–610.
33. Mn, N.; Winding, A. *Microorganisms as Indicators of Soil Health*; National Environmental Research Institute: Aarhus, Denmark, 2002.

34. Badri, D.V.; Vivanco, J.M. Regulation and function of root exudates. *Plant Cell Environ.* **2009**, *32*, 666–681. [[CrossRef](#)]
35. Bora, S.S.; Hazarika, D.J.; Gogoi, R.; Dullah, S.; Gogoi, M.; Barooah, M. Long-term pruning modulates microbial community structure and their functional potential in Tea (*Camellia sinensis* L.) soils. *Appl. Soil Ecol.* **2022**, *176*, 104483. [[CrossRef](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.