

Article

Effect of Ulvan on the Biocontrol Activity of *Debaryomyces hansenii* and *Stenotrophomonas rhizophila* against Fruit Rot of *Cucumis melo* L.

Tomas Rivas-García ¹, Bernardo Murillo-Amador ¹, Alejandra Nieto-Garibay ¹ ,
Roberto G. Chiquito-Contreras ², Gabriel Rincon-Enriquez ³  and
Luis G. Hernandez-Montiel ^{1,*} 

¹ Centro de Investigaciones Biológicas del Noroeste, 23096 La Paz, Mexico; eltom_r@hotmail.com (T.R.-G.); bmurillo04@cibnor.mx (B.M.-A.); anieto04@cibnor.mx (A.N.-G.)

² Facultad de Ciencias Agrícolas Universidad Veracruzana, 91090 Xalapa Veracruz, Mexico; contrerasch@yahoo.com

³ Centro de Investigaciones en Tecnología y Diseño del estado de Jalisco, 44270 Guadalajara Jalisco, Mexico; grincon@ciatej.mx

* Correspondence: lhernandez@cibnor.mx; Tel.: +52-612-123-8484

Received: 18 October 2018; Accepted: 19 November 2018; Published: 22 November 2018



Abstract: In the present study, the following was investigated: (a) The effect of ulvan on in vivo and in vitro biocontrol of *Debaryomyces hansenii* and *Stenotrophomonas rhizophila* against *Fusarium proliferatum* and (b) the effect of ulvan on in vivo and in vitro growth of *D. hansenii* and *S. rhizophila* and muskmelon quality parameters. The results showed that the biocontrol activity of *D. hansenii* and *S. rhizophila* could be enhanced by ulvan (5 g/L). The combination of ulvan and *S. rhizophila* resulted in a more effective control of fruit rot in comparison to fungicide benomyl. On in vitro growth of *F. proliferatum*, individual treatments of *D. hansenii* and *S. rhizophila* inhibited spore germination and mycelial growth with no statistical difference with the combined treatments. Ulvan does not have a direct effect on the in vivo and in vitro growth of *D. hansenii* and *S. rhizophila*. Furthermore, the combined treatments improve the natural disease incidence and quality parameters like weight, firmness, total soluble solids (TSS), and pH. These results suggest that the use of ulvan may be an effective method to improve the biological activity of *D. hansenii* and *S. rhizophila*.

Keywords: biocontrol; *Debaryomyces hansenii*; *Stenotrophomonas rhizophila*; ulvan; muskmelon; *Fusarium* spp.

1. Introduction

Muskmelon fruit (*Cucumis melo* L.) is commercialized worldwide because of its flavor and nutritional content [1]. During fruit ripening, it is easily perishable and susceptible to fungal pathogens during storage, transportation, and commercialization [2]. Fruit rot caused by *Fusarium* spp. is one of the most serious diseases of melon fruit, and it is generally controlled by applying synthetic fungicides [3]. However, indiscriminate use of synthetic fungicides causes environmental problems, puts humans at risk, and may proliferate fungicide resistance [4].

Biological control of postharvest disease is an effective and nonchemical alternative. It relies on the use of antagonist microorganisms which limit or stop the development of fungal pathogens [5]. In previous studies, the yeast *Debaryomyces hansenii* and the bacteria *Stenotrophomonas rhizophila* have showed good results and promissory characteristics as biological control agents. *D. hansenii* and *S. rhizophila* have significantly inhibited pathogens like *Aspergillus* spp., *Fusarium* spp., *Colletotrichum* spp.,

and *Penicillium* spp., among others [6–10]. *D. hansenii* has been considered as a potential biocontrol agent due to volatile organic compound (VOC) production, β -1, 3 glucanase and protease activity, inhibition of spore germination, and the competition for nutrients like saccharose, glucose, fructose and total carbohydrates [11]. *S. rhizophila* is another potential biocontrol agent because it produces lithic enzymes, siderophores, and secondary metabolites, which act as antifungal compounds [12,13].

Microbial antagonists, when applied individually, usually have a much lower level of effectiveness compared to that of synthetic fungicides [5]. Nonetheless, their activity can be enhanced by manipulation of the environment, using mixtures of beneficial organisms and physiological and genetic enhancement or biocontrol with other methods, such as low doses of fungicides and other chemicals [14]. Improvement of the biocontrol agents effect may result in direct inhibition of the pathogen, elicitation of systemic acquired resistance in the host tissue, and stimulation of the microbial antagonists [15]. Biological control agents (BCAs) combined with chemicals like sodium bicarbonate, harpin, quitosan, and ulvan have been demonstrated to provide enhanced characteristics in controlling fruit decay [16–19].

Ulvan is a polysaccharide isolated from green algae of the genus *Ulva* and it has been used as an alternative treatment for chemical fungicides [20]. These sulfated heteropolisaccharide can reduce the disease severity of many pathogens of plants at concentrations of 5 g L^{-1} or less [19–25]. In apple, ulvan reduces the mycelial growth of *Colletotrichum gloeosporioides*, decreases disease severity to 66%, and increases peroxidase and glucanase activity in the host [22]. In *Arabidopsis thaliana*, ulvan increased nicotinamide adenine dinucleotide hydrogen (NADPH) oxidase activity and hydrogen peroxide levels [23]. In *Medicago truncatula*, ulvan is an efficient elicitor of resistance, which confers protection against *Colletotrichum trifolii* [24]. In tomato, oligoulvan reduced the severity caused by *Fusarium oxysporum* f. sp. *lycopersici*, stimulating phenyl alanine ammonia lyase, increasing the phenolic compounds, and inducing salicylic acid synthesis [25].

To our knowledge, no study has been conducted to determine the ability of ulvan to improve the biocontrol activity of the yeast *D. hansenii* and the bacteria *S. rhizophila*. In particular, the objectives of this study were to evaluate (a) the effect of ulvan on in vivo and in vitro control of *D. hansenii* and *S. rhizophila* against *F. proliferatum* and (b) the effect of ulvan on in vitro and in vivo growth of *D. hansenii* and *S. rhizophila* and muskmelon quality parameters.

2. Materials and Methods

2.1. Microorganisms and Fruit Materials

2.1.1. Fruit

Muskmelon (*Cucumis melo* L. var. *reticulatus*) fruit were collected from a commercial orchard located in El Pescadero, Baja California Sur, México. Fruit of uniform size at commercial maturity stage were selected and transported immediately to the laboratory. Fruit without physical damage or symptoms of fruit rot were disinfected with 2% (v/v) sodium hypochlorite for 2 min, washed with sterile distilled water, air dried at room temperature, and placed in plastic containers prior to use.

2.1.2. Pathogen Inoculum

The pathogen *Fusarium proliferatum* was previously isolated from infected melon fruits and maintained in potato dextrose agar (PDA, at dose of 39 g L^{-1}) plates at $4 \text{ }^\circ\text{C}$ for storage [26]. To reactivate the culture and verify their pathogenicity, the pathogen was inoculated into wounded melon fruits and re-isolated onto PDA after infection was established. Spore suspension was obtained from 10 days old cultures PDA at $25 \text{ }^\circ\text{C}$, and spore concentration was determined using a hemocytometer and adjusted to 10^4 spores/mL with sterile distilled water prior to use.

2.1.3. Antagonist Microorganisms

The antagonist microorganisms were obtained from the Centro de Investigaciones Biológicas del Noroeste (CIBNOR), La Paz Baja California Sur, México, and were originally isolated from the Ojo de Liebre hyperhaline lagoon (27°35' and 27°52' north latitude and 113° 58' and 114°0' west latitude). *D. hansenii* and *S. rhizophila* were maintained in PDA and tripticase soy agar (TSA, at dose of 40 g L⁻¹) plates respectively at 4 °C for storage. Liquid cultures of *D. hansenii* and *S. rhizophila* were grown in 250 mL Erlenmeyer flasks containing 50 mL of potato dextrose broth (PDB, at dose of 39 g L⁻¹) and tripticase soy broth (TSB, at dose of 40 g L⁻¹), respectively; both microorganisms had been inoculated with a loop of each culture and were incubated on a rotary shaker at 180 rpm and 27 °C. *D. hansenii* concentration was adjusted to 1 × 10⁶ CFU mL⁻¹ with a hemocytometer and the cells suspension of *S. rhizophila* was adjusted to 1 × 10⁸ CFU mL⁻¹ using a UV/V spectrophotometer (HACH, Dusseldorf, Germany) at 660 nm and absorbance of 1.

2.1.4. Chemical Treatments

Ulvan (OligoTech®, Elicityl Ltd., Crolles, France) solution was prepared at 5 g L⁻¹ using sterile deionized water. The synthetic fungicide used in this study was benomyl at 1000 ppm and synthetic bactericide bactrol at 500 ppm.

2.2. Effect of Ulvan, *D. hansenii*, and *S. rhizophila* against *F. proliferatum* In Vivo

In this experiment, six equidistant 3 mm wounds in diameter were performed in each fruit and were inoculated with 20 µL of the following: (1) 1 × 10⁶ CFU mL⁻¹ *D. hansenii*; (2) 1 × 10⁸ CFU mL⁻¹ *S. rhizophila*; (3) 5 g L⁻¹ ulvan; (4) 1 × 10⁶ CFU mL⁻¹ *D. hansenii* + 5 g L⁻¹ ulvan; (5) 1 × 10⁸ CFU mL⁻¹ *S. rhizophila* + 5 g L⁻¹ ulvan; (6) sterile distilled water (control); and (7) 1000 ppm benomyl. They were left to dry for 2 h and then a suspension (20 µL) of 1 × 10⁴ spores mL⁻¹ *F. proliferatum* was inoculated into each wound. Fruits were placed in plastic containers at 27 °C and 90% of relative humidity (RH) for 7 days. Disease control and lesion diameter (mm) were measured. Disease control (DC) was calculated by the formula: $100 - ([100 \times Fi]/Tf)$, where F_i = number of infected fruits in each treatment and T_f = total of infected fruits in control treatment. Each treatment consisted of three fruits and was replicated ten times.

2.3. Effect of Ulvan, *D. hansenii*, and *S. rhizophila* against *F. proliferatum* In Vitro

2.3.1. Effect on Mycelial Growth

The effects of ulvan, *D. hansenii*, and *S. rhizophila* on the mycelial growth of *F. proliferatum* were assessed as described by Zhou et al. [27]. A 10 mm diameter hole was made in a 90 mm diameter plate containing 20 mL of nutrient broth (NB, at dose of 31 g/L). As treatments, 100 µL of: (1) 1 × 10⁶ CFU mL⁻¹ *D. hansenii*; (2) 1 × 10⁸ CFU mL⁻¹ *S. rhizophila*; (3) 5 g L⁻¹ ulvan; (4) 1 × 10⁶ CFU mL⁻¹ *D. hansenii* + 5 g L⁻¹ ulvan; (5) 1 × 10⁸ CFU mL⁻¹ *S. rhizophila* + 5 g L⁻¹ ulvan; (6) sterile distilled water (control); and (7) 1000 ppm benomyl were deposited into the holes. After 2 h, 100 µL 1 × 10⁴ spores mL⁻¹ suspension of *F. proliferatum* was deposited into each hole. The plates were incubated at 27 °C for 7 days. The mycelial radial growth was measured with the ImageJ® program, which measured the relative area of the calibration parameter and the lesion site in pixels and converted the measurement of lesion site in mm² based on the known value of the calibration. Inhibition percentage (I%) was calculated by the following equation: $I\% = (G_c - G_t)/G_c \times 100$, in which G_c means the radial growth of the pathogen in the control treatments and G_t means the radial growth of the pathogen with the treatments. Each treatment was replicated ten times.

2.3.2. Effect on Spore Germination

The effect of ulvan, *D. hansenii*, and *S. rhizophila* on spore germination was assayed according to the method of Mattiuz et al. [28]. We deposited 50 μL 1×10^4 spores mL^{-1} suspension of *F. proliferatum* into an Eppendorf® tube. Then, 50 μL of: (1) 1×10^6 CFU mL^{-1} *D. hansenii*; (2) 1×10^8 CFU mL^{-1} *S. rhizophila*; (3) 5 g L^{-1} ulvan; (4) 1×10^6 CFU mL^{-1} *D. hansenii* + 5 g L^{-1} ulvan; (5) 1×10^8 CFU mL^{-1} *S. rhizophila* + 5 g L^{-1} ulvan; (6) sterile distilled water (control); and (7) 1000 ppm benomyl were deposited into the Eppendorf® tubes. After incubation at 27 °C on a rotary shaker set at 180 rpm, an aliquot of 20 μL was taken every 12 h to observe the spore germination rate with an optical microscope (CARL ZEISS, Primo Star, Oberkochen, Germany). The experiment finished when control treatment reached a 100% germinated spores. A spore was considered as being germinating if its germ tube was longer than the spore itself. The germination inhibition was obtained by counting the number of germinated spores (NGS) among the first 100 spores observed. Each treatment was replicated ten times and inhibition ratio was calculated as being $(100 - \text{NGS}) \times 100/100$, and was expressed as a percentage (%).

2.4. Effect of Ulvan on Populations of *D. hansenii* and *S. rhizophila* in Vivo

The fruit were disinfected and wounded as described before. Then, the wounds were treated with 20 μL of a cell suspension of: (1) 1×10^6 CFU mL^{-1} *D. hansenii*; (2) 1×10^8 CFU mL^{-1} *S. rhizophila*; (3) 1×10^6 CFU mL^{-1} *D. hansenii* + 5 g L^{-1} ulvan; and (4) 1×10^8 CFU mL^{-1} *S. rhizophila* + 5 g L^{-1} ulvan, (5) 1×10^6 CFU mL^{-1} *D. hansenii* + 1000 ppm benomyl, and (6) 1×10^8 CFU mL^{-1} *S. rhizophila* + 500 ppm bactrol. Sterile distilled water was used as control. Fruits were placed in plastic containers at 27 °C and 90% RH. *D. hansenii* and *S. rhizophila* were recovered from the wounds 1 h after inoculation (time 0) and after 1, 2, 3, and 4 days. Wounded tissue was removed with an ethanol-flamed, 5 mm cork borer and ground in sterile mortar with 5 mL of sterile 0.05 M phosphate buffer (pH 7.0). The number of CFU of the yeast and bacteria was determined using the dilution plating technique. The results were expressed as Log_{10} CFU/wound. Each treatment consisted of three fruits and was replicated ten times.

2.5. Effect of Ulvan on the Growth of *D. hansenii* and *S. rhizophila* In Vitro

The experiment was conducted in petri plates (90 mm) of PDA for *D. hansenii* and TSA for *S. rhizophila* (20 mL per plate). One hundred milliliters of: 1×10^6 CFU mL^{-1} *D. hansenii*; or 1×10^8 CFU mL^{-1} *S. rhizophila* was spread with an ethanol-flamed glass rod. After 2 h, 4 equidistant holes (10 mm diameter) were made and 100 μL of: (1) 5 g L^{-1} ulvan for both microorganisms, (2) 1000 ppm benomyl for *D. hansenii*, or (3) 500 ppm bactrol for *S. rhizophila* were deposited in 4 holes. Sterile distilled water was used as control. The plates were incubated at 27 °C for 2 days. The number of CFU was determined by colony density and inhibition percentage (I%) was calculated by the following equation: $I\% = (\text{Ga} - \text{Gu})/\text{Gc} \times 100$, in which Ga means the CFU of the antagonist in the control treatments and Gu means the CFU of the antagonist with chemicals. Each treatment was replicated ten times.

2.6. Effect of *D. hansenii* or *S. rhizophila* in Combination with Ulvan on Natural Disease Incidence

Intact fruit were saturated in treatment solutions as follows: (1) 1×10^6 CFU mL^{-1} *D. hansenii*; (2) 1×10^8 CFU mL^{-1} *S. rhizophila*; (3) 1×10^6 cells mL^{-1} *D. hansenii* + 5 g L^{-1} ulvan; (4) 1×10^8 CFU mL^{-1} *S. rhizophila* + 5 g L^{-1} ulvan; (5) sterile distilled water; and (6) 1000 ppm benomyl. Each fruit was kept soaked for 2 min, air dried at room temperature (25 °C) for 2 h, and packed in plastic containers at 27 °C and 90% relative humidity (RH) for 7 days. The numbers of decayed fruits were recorded, the incidence of decayed fruit was evaluated, and the quality parameters were determined. Each treatment was replicated ten times.

Determination of Quality Parameters

The weight loss (%), fruit firmness (N), total soluble solids (%), and pH were measured to evaluate the effect of ulvan, *D. hansenii*, and *S. rhizophila* on quality parameters. For weight loss, fruit from plastic

containers were weighed before and after storage. Firmness values were determined by compression after application of a load of 9.8 N using the GY- texture Analyzer at two opposite sides of the equatorial region of the fruit. A homogeneous sample of fruit was prepared by crushing it in the mortar and pestle and a few drops of fruit juice were used for the total soluble solids (TSS) and pH. TSS was determined with a digital Abbe refractometer (PR—32, Atago Co., Tokyo, Japan) at room temperature. A few drops of the fruit juice were placed on the refractometer for measurement of total soluble solids percentage.

2.7. Statistical Analysis

To assess the advantage of in vivo combined postharvest treatments (BCAs + ulvan) with respect to the same treatments applied alone (BCAs or ulvan), the type of interaction (additive, synergistic, or antagonistic) was evaluated. The synergy factor (SF) was calculated according to the Abbott's formula [29]: $SF = EO/EE$; where EO and EE are, respectively, the observed and expected biocontrol percentage (C%) of the combination. EE was calculated as follows: $(Ea + Eb) - (Ea \times Eb/100)$, where $Ea = C\%$ of postharvest treatment (BCAs); $Eb = C\%$ of postharvest treatment b (ulvan). If $SF = 1$, the interaction between the combination treatments was identified as additive; if $SF < 1$, the interaction was antagonistic, and if $SF > 1$, the interaction was synergistic.

The data were processed by one-way analysis of variance (ANOVA). Statistical data analyses were performed using the software program STATISTICA 10.0, and the post hoc least significant difference Fisher test ($p \leq 0.05$) was used for comparison of the means. When it was necessary, data were transformed into arcsine square root values to normalize distribution before analysis of variance.

3. Results

3.1. Effect of Ulvan, *D. hansenii*, and *S. rhizophila* on Muskmelon Fruit Rot by *F. proliferatum*

D. hansenii, *S. rhizophila*, and ulvan had a significant effect on disease control of fruit rot caused by *F. proliferatum* on muskmelon fruit stored at 27 °C for 7 days (Table 1). While ulvan treatment had a slight effect on disease control with 14.3%, in combination with *S. rhizophila*, it had the most effective disease control and was significant similar to benomyl treatment (64.3%). According to Abbott's formula [29], the treatments *D. hansenii* + ulvan and *S. rhizophila* + ulvan had synergistic effect in comparison with their individual treatments. On lesion diameter (Figure 1), single treatment of *S. rhizophila* was better than *D. hansenii* + ulvan treatment. *S. rhizophila* + ulvan were the most effective treatment on reducing lesion diameter, but not better than benomyl. These results suggest that ulvan enhances the biocontrol activity of *D. hansenii* and *S. rhizophila* against fruit rot caused by *F. proliferatum* in muskmelon fruit.

Table 1. Effect of *Debaryomyces hansenii*, *Stenotrophomonas rhizophila*, and ulvan on disease control of muskmelon fruit rot by *Fusarium proliferatum* incubated at 27 °C for 7 days.

Treatments	Disease Control (%)	Ee	SF
<i>D. hansenii</i>	28.6 ± 3.1 ^a *	-	-
<i>S. rhizophila</i>	35.7 ± 2.8 ^b	-	-
ulvan	14.3 ± 3.9 ^c	-	-
<i>D. hansenii</i> + ulvan	57.1 ± 4.3 ^d	38.8	1.5
<i>S. rhizophila</i> + ulvan	64.3 ± 4.1 ^e	44.9	1.4
Benomyl	64.3 ^a ± 2.4 ^e	-	-

Disease control: Biocontrol percentage for each treatment was calculated by comparison with the control (0%) (*F. proliferatum* + water). Ee: Expected effect or expected biocontrol percentage, SF: Synergy factor, see Materials and Methods. * The values are means ± standard deviation of ten replicates (three fruits each). Different letters indicate significant difference ($p \leq 0.05$) according to Fisher test.




Figure 1. Effects of the yeast *Debaryomyces hansenii*, the bacteria *Stenotrophomonas rhizophila*, and ulvan on lesion diameter (mm) of muskmelon fruit by *Fusarium proliferatum* during incubation at 27 °C for 7 days. Vertical bars are means \pm standard deviation of ten replicates (three fruits each) Bars with different letters are significantly different ($p \leq 0.05$) according to Fisher test.

3.2. Effect of *D. hansenii*, *S. rhizophila*, and *Ulvan* on Mycelial Growth and Spore Germination of *F. proliferatum*

On PDA media, ulvan had no inhibiting effect on the growth of *F. proliferatum* compared with the control (Figure 2a). Single treatments of *D. hansenii* and *S. rhizophila* significantly inhibited the growth of *F. proliferatum* in vitro. However, *D. hansenii* and *S. rhizophila* in combination with ulvan did not differ from single treatments with BCAs. The results on spore germination inhibition showed a similar pattern to mycelial growth inhibition against *F. proliferatum* (Figure 2b). These results suggest that ulvan does not have a direct effect on mycelial growth and spore germination inhibition of *F. proliferatum*.




Figure 2. Cont.




Figure 2. Inhibitory effect of the yeast *D. hansenii*, the bacteria *S. rhizophila*, and ulvan on (a) mycelial growth and (b) spore germination rate of *F. proliferatum* on potato dextrose agar (PDA). Vertical bars are means \pm standard deviation of ten replicates. Bars with different letters are significantly different ($p \leq 0.05$) according to Fisher test.

3.3. Effect of Ulvan on the In Vivo Population Dynamics or in the In Vitro Growth of *D. hansenii* and *S. rhizophila*

Ulvan exhibited no significant negative effects both on the in vivo populations (Figure 3) and on the in vitro growth of *D. hansenii* and *S. rhizophila* (Figure 4). Both BCAs grew efficiently on muskmelon tissue during the period of time quantified. After 3 days of incubation, there were significant differences between *S. rhizophila* and *S. rhizophila* + ulvan treatments. It was found that for the rest of treatments during the period of incubation, there were no significant differences between single and addition of ulvan on treatments. Chemicals totally inhibited the in vivo growth of BCAs. These results suggest that ulvan does not have a direct effect on the in vivo and in vitro development of both BCAs.




Figure 3. Population dynamics of the yeast *D. hansenii* and the bacteria *S. rhizophila* with and without ulvan in wounds at 27 °C. Chemical treatments totally inhibited biological control agents (BCAs) since the first day after inoculation. Sterile distilled water was used as control. The data presented are the means \pm standard deviation of ten replicates. Different letters are significantly different ($p \leq 0.05$) according to Fisher test.




Figure 4. Effect of ulvan on the yeast *D. hansenii* and the bacteria *S. rhizophila* on PDA plates for yeast and case soy broth (TSB) plates for bacteria incubated at 27 °C for 2 days. Chemical treatments consisted of benomyl for *D. hansenii* and bactrol for *S. rhizophila*. Sterile distilled water was used as control. Vertical bars are means \pm standard deviation of ten replicates. Bars with different letters are significantly different ($p \leq 0.05$) according to Fisher test.

3.4. Effect of *D. hansenii*, *S. rhizophila*, and Ulvan on Natural Disease Incidence and Quality Parameters

Fruit surfaces were inoculated to explore the preventive activity of *D. hansenii* and *S. rhizophila* alone or in combination with ulvan against fruit rot. Even though 70% of the fruits in the control developed decay symptoms after 7 days of storage, the disease incidence (DI) was significantly reduced with all treatments (Table 2). DI under treatment with *S. rhizophila* + ulvan (13.3%) was statistically similar to treatment with benomyl (10.0%). The combined treatments of BCAs with ulvan were more effective to reduce the disease incidence of muskmelon fruits than the single application. The quality parameters of three fruits from ten replicates were measured (Table 2). Treatment with benomyl and water (control) significantly increased weight loss and decreased the firmness of muskmelon fruit. On total soluble solids (TSS), no significant effects were observed. Single treatment of ulvan or the combination with *D. hansenii* or *S. rhizophila* decreased the pH level.

Table 2. Effect of *D. hansenii*, *S. rhizophila*, and ulvan on natural disease incidence and quality parameters of muskmelon fruit after 7 days of incubation at 27 °C.

Treatments	DI (%)	Weight Loss (gr)	Firmness (N)	TSS (%)	pH
<i>D. hansenii</i>	33.3 \pm 1.2 ^{a*}	0.30 \pm 0.02 ^a	4.2 \pm 0.5 ^a	9.2 \pm 0.08 ^{ac}	6.5 \pm 0.1 ^a
<i>S. rhizophila</i>	26.7 \pm 1.6 ^b	0.30 \pm 0.01 ^a	4.2 \pm 0.4 ^a	9.2 \pm 0.09 ^a	6.5 \pm 0.1 ^a
ulvan	23.3 \pm 0.8 ^c	0.24 \pm 0.03 ^a	4.2 \pm 0.4 ^a	9.3 \pm 0.06 ^b	6.1 \pm 0.1 ^b
<i>D. hansenii</i> + ulvan	20.0 \pm 1.2 ^c	0.21 \pm 0.03 ^a	4.3 \pm 0.3 ^b	9.3 \pm 0.08 ^{bc}	6.2 \pm 0.1 ^b
<i>S. rhizophila</i> + ulvan	13.3 \pm 0.7 ^d	0.22 \pm 0.02 ^a	4.3 \pm 0.3 ^b	9.3 \pm 0.06 ^b	6.1 \pm 0.1 ^b
Benomyl	10.0 \pm 0.4 ^d	0.68 \pm 0.05 ^b	4.1 \pm 0.8 ^c	9.2 \pm 0.08 ^a	6.6 \pm 0.1 ^c
Control (water)	70.0 \pm 1.4 ^e	1.06 \pm 0.08 ^c	4.0 \pm 0.6 ^d	9.2 \pm 0.07 ^a	6.6 \pm 0.08 ^c

* The values are means \pm standard deviation of ten replicates (three fruits each). Different letters indicate significant difference ($p \leq 0.05$) according to Fisher test. DI = disease incidence, TSS = Total soluble solids.

4. Discussion

The efficacy of biological control alternatives should be comparable to the level of control provided by conventional fungicides in order to obtain general acceptance [30]. Obtaining such high levels of control with biological control alternatives is difficult [15]. Thus, there is a tendency to promote an integrated system rather than a single one [31]. The results in the present study demonstrate that the combination of ulvan with *D. hansenii* or *S. rhizophila* can result in a significant enhanced control of fruit rot in muskmelon, compared with single treatments of *D. hansenii*, *S. rhizophila*, and ulvan (Figure 1) (Table 1). Previous studies have demonstrated that the biocontrol ability of different antagonist microorganisms to control postharvest diseases of fruits can be significantly improved by

combining alternative, but compatible treatments [32]. Chitosan possess antifungal properties and the ability to elicit host defense responses; hence, it has been suggested as an effective additive to improve the biocontrol performance of the antagonistic yeasts *Candida saitoana* and *C. laurentii* [33]. The combination of NaHCO_3 and the bacterial antagonist *Burkholderia spinosa* is effective on the suppression of anthracnose, crown rot, and blossom end rot on banana [34].

To understand the direct effect of ulvan alone or in combination with *D. hansenii* and *S. rhizophila* on the growth of *F. proliferatum*, we investigated the effects of ulvan on mycelial growth and spore germination in vitro. In previous studies, ulvan promotes the conidial germination and appressoria formation of *Colletotrichum gloeosporioides* [35,36]. Nonetheless, our results showed that ulvan did not influence the mycelial growth of *F. proliferatum*. However, it slightly inhibited the spore germination of *F. proliferatum*, probably due to chemical PDA alteration (Figure 2). By contrast, single treatments of *D. hansenii* or *S. rhizophila* were found to be similar to the combined treatments with ulvan on spore germination inhibition. *D. hansenii* and *S. rhizophila* have the capacity to alleviate unfavorable conditions and grow efficiently [11,13]. Both BCAs probably reduce the slight spore germination by inhibition due to chemical alteration of the medium. It has been reported that ulvan did not exhibit any direct fungal activity against *Alternaria brassicola*, *Colletotrichum lindemuthianum*, and *Uromyces appendiculatus* [37–39]. These different results might be related to different sensitivities of fungal species to ulvan and dissimilar testing methods might explain these apparently contradictory results on fungal species.

We propose that the efficacy of ulvan in combination with *D. hansenii* and/or *S. rhizophila* in controlling fruit rot of muskmelon by *F. proliferatum* might be linked to the fruit-mediated mechanisms which increase defense response, like priming, PR proteins, and oxidative burst. Jaulneau et al. [40] elucidated that the ulvan-induced defense response on *Medicago truncatula* is mediated by the jasmonic acid signaling pathway. On wheat and rice, ulvan has a priming effect by increasing the initial oxidative burst and by enhancing the resistance against powdery mildew [41]. Cluzet et al. [24] concluded by a microarray that ulvan increases the expression of codificant genes to phytoalexins, PR proteins, and structural proteins. Although ulvan has a small effect on disease control, its effect is attributed to systemic acquired resistance (SAR) mechanisms and priming, which acts after systemic induced resistance (ISR) mechanisms [24,40,41]. Further investigation needs to be carried out in order to elucidate which mechanisms of resistance are induced in muskmelon fruit by ulvan.

During the last two decades, numerous BCAs have been isolated, identified, and applied to control postharvest decay of different fruits and vegetables [32]. It is crucial for BCAs to colonize fruit tissue more efficiently than the pathogen to compete for space and nutrients [42]. Moreover, our data showed that *D. hansenii* and *S. rhizophila* grew rapidly in muskmelon wounds (Figure 3). However, the direct antifungal activity of BCAs is an important mechanism of action during the colonization time and the efficacy of this mechanism relies on the rapid colonization level of BCAs, thereby inhibiting the early pathogenic process by fungi [42]. Previous studies have shown that *D. hansenii* significantly inhibited the mycelial growth of *Monilinia fruticola* (74.4%) and *M. fructigena* (44.1%), and *S. rhizophila* significantly inhibited the mycelial growth of *C. gloeosporioides* (93%) [8,43]. According to previous results, *D. hansenii* and *S. rhizophila* control the in vitro growth of *F. proliferatum* by mechanisms such as mycoparasitism by lytic enzymes and secondary metabolites excretion, e.g., surfactants and volatile organic compounds [44]. In addition, ulvan did not have any influence on the growth of the BCAs in vivo or in vitro (Figures 3 and 4). To our knowledge, this is the first time that ulvan has been applied in combination with BCAs to control fungal diseases of fruit.

It is known that resistance in harvested fruit is associated with levels of senescence, as it drops considerably with the onset of tissue senescence [45]. In a previous study, the effect of combining *Pichia membranefaciens* and benzo-thiadiazole-7-carbothioic acid S-methyl ester (BTH) on the control of blue mold by *Penicillium expansum* in peach fruit showed that quality parameters were not impaired [31]. Ulvan in combination with both BCAs decreased natural disease incidence and significantly maintained the fruit firmness and weight (Table 2). An initial increase in the TSS content of fruit may be due to the

hydrolysis of insoluble polysaccharide into simple sugars, but subsequently, TSS content decreased as the storage period increased, which is related to a higher respiratory process [46]. In our results, ulvan by the combined treatments with BCAs showed the highest TSS values by probably delaying the respiratory process. During muskmelon fruit maturation, the pH increases from approximately 5.3 to approximately 6.8 [45]. The increase in pH promotes the activity of the poligalacturonase enzyme that is related to the pathogenicity and virulence of *F. proliferatum* [47,48]. In our results, ulvan and the combined treatments with BCAs maintained the lowest pH values.

5. Conclusions

The results presented in our study showed that ulvan enhances the effect of *D. hansenii* and *S. rhizophila* in controlling fruit rot in muskmelon, but ulvan does not have direct effect on BCA growth. Also elucidated was the effect of ulvan on pathogenicity of *F. proliferatum* and development of fruit rot disease of muskmelon. The mode of action of both treatments may have complemented each other, whereby ulvan probably provides protection by muskmelon resistance induction, and BCAs inhibit fungal growth colonization by competition for space and direct antifungal activity. However, the mechanism by which ulvan enhanced the biocontrol efficacy of BCAs is complex and may be a result of several different interactions among ulvan, BCAs (*D. hansenii* or *S. rhizophila*) pathogen, and fruits. In this study, we found a new methodology to improve the performance of the antagonistic *D. hansenii* and *S. rhizophila* for controlling fruit rot in muskmelon during postharvest stage.

Author Contributions: Conceptualization, L.G.H.-M. and B.M.-A.; methodology, L.G.H.-M. and T.R.-G.; software, G.R.-E. and A.N.-G.; validation, L.G.H.-M., R.G.C.-C., and T.R.-G.; formal analysis, B.M.-A. and A.N.-G.; investigation, T.R.-G., L.G.H.-M., and B.M.-A.; resources, G.R.-E. and R.G.C.-C.; data curation, B.M.-A. and T.R.-G.; writing—original draft preparation, L.G.H.-M. and T.R.-G.; writing—review and editing, L.G.H.-M., A.N.-G., and G.R.-E.; visualization, T.R.-G.; supervision, R.G.C.-C. and A.N.-G.; project administration, L.G.H.-M.; funding acquisition, L.G.H.-M.

Funding: This research was funded by the grant project SEP-CONACYT 181972 and grant project Problemas nacionales 2015-01-352 CONACYT (Consejo Nacional de Ciencia y Tecnología).

Acknowledgments: The CONACYT by grant awarded to T. Rivas Garcia and Michael Cordoba, a native English-speaking editor, for editing the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Sánchez-Estrada, A.; Tiznado-Hernández, M.E.; Ojeda-Contreras, A.J.; Valenzuela-Quintanar, A.I.; Troncoso-Rojas, R. Induction of enzymes and phenolic compounds related to the natural defense response of netted melon fruit by a bio-elicitor. *J. Phytopathol.* **2009**, *157*, 24–32. [[CrossRef](#)]
2. Huang, K.; Zou, Y.; Luo, J.; Liu, Y. Combining UV-C treatment with biocontrol yeast to control postharvest decay of melon. *Environ. Sci. Pollut. Res.* **2015**, *22*, 14307–14313. [[CrossRef](#)] [[PubMed](#)]
3. Jamalzadeh, M.; Etebarian, H.R.; Aminian, H.; Alizadeh, A. A review of mechanisms of action of biological control organisms against post-harvest fruit spoilage. *EPPO Bull.* **2011**, *41*, 65–71. [[CrossRef](#)]
4. Mari, M.; Di Francesco, A.; Bertolini, P. Control of fruit postharvest diseases: old issues and innovative approaches. *Stewart Postharvest Rev.* **2014**, *10*, 1–4.
5. Droby, S.; Wisniewski, M.; Macarasin, D.; Wilson, C. Twenty years of postharvest biocontrol research: Is it time for a new paradigm? *Postharvest Biol. Technol.* **2009**, *52*, 137–145. [[CrossRef](#)]
6. Medina-Córdova, N.; López-Aguilar, R.; Ascencio, F.; Castellanos, T.; Campa-Córdova, A.I.; Angulo, C. Biocontrol activity of the marine yeast *Debaryomyces hansenii* against phytopathogenic fungi and its ability to inhibit mycotoxins production in maize grain (*Zea mays* L.). *Biol. Control* **2016**, *97*, 70–79. [[CrossRef](#)]
7. Hernández-Montiel, L.G.; Ochoa, J.L.; Troyo-Diéguez, E.; Larralde-Corona, C.P. Biocontrol of postharvest blue mold (*Penicillium italicum* Wehmer) on Mexican lime by marine and citrus *Debaryomyces hansenii* isolates. *Postharvest Biol. Technol.* **2010**, *56*, 181–187. [[CrossRef](#)]

8. Hernandez-Montiel, L.G.; Zulueta-Rodriguez, R.; Angulo, C.; Rueda-Puente, E.O.; Quiñonez-Aguilar, E.E.; Galicia, R. Marine yeasts and bacteria as biological control agents against anthracnose on mango. *J. Phytopathol.* **2017**, *165*, 833–840. [[CrossRef](#)]
9. Hernandez-Montiel, L.G.; Gutierrez-Perez, E.D.; Murillo-Amador, B.; Vero, S.; Chiquito-Contreras, R.G.; Rincon-Enriquez, G. Mechanisms employed by *Debaryomyces hansenii* in biological control of anthracnose disease on papaya fruit. *Postharvest Biol. Technol.* **2018**, *139*, 31–37. [[CrossRef](#)]
10. Hernandez-Montiel, L.G.; Rivas-Garcia, T.R.; Romero-Bastidas, M.; Chiquito-Contreras, C.J.; Ruiz-Espinoza, F.H.; Chiquito-Contreras, R.G. Potencial antagonico de bacterias y levaduras marinas para el control de hongos fitopatogenos. *REMEXCA* **2018**, *20*, 4311–4321. [[CrossRef](#)]
11. Medina-Córdova, N.; Rosales-Mendoza, S.; Hernández-Montiel, L.G.; Angulo, C. The potential use of *Debaryomyces hansenii* for the biological control of pathogenic fungi in food. *Biol. Control* **2018**, *121*, 216–222. [[CrossRef](#)]
12. Berg, G.; Egamberdieva, D.; Lugtenberg, B.; Hagemann, M. Symbiotic plant–microbe interactions: Stress protection, plant growth promotion, and biocontrol by *Stenotrophomonas*. In *Symbioses and Stress*; Springer: Dordrecht, The Netherlands, 2010; pp. 445–460.
13. Schmidt, C.S.; Alavi, M.; Cardinale, M.; Müller, H.; Berg, G. *Stenotrophomonas rhizophila* DSM14405T promotes plant growth probably by altering fungal communities in the rhizosphere. *Biol. Fertil. Soils* **2012**, *48*, 947–960.
14. Nunes, C.A. Biological control of postharvest diseases of fruit. *Eur. J. Plant Pathol.* **2012**, *133*, 181–196. [[CrossRef](#)]
15. Dukare, A.S.; Paul, S.; Nambi, V.E.; Gupta, R.K.; Singh, R.; Sharma, K.; Vishwakarma, R.K. Exploitation of microbial antagonists for the control of postharvest diseases of fruits: A review. *Crit. Rev. Food Sci. Nutr.* **2018**, 1–16. [[CrossRef](#)] [[PubMed](#)]
16. Hong, P.; Hao, W.; Luo, J.; Chen, S.; Hu, M.; Zhong, G. Combination of hot water, *Bacillus amyloliquefaciens* HF-01 and sodium bicarbonate treatments to control postharvest decay of mandarin fruit. *Postharvest Biol. Technol.* **2014**, *88*, 96–102. [[CrossRef](#)]
17. Tang, J.; Liu, Y.; Li, H.; Wang, L.; Huang, K.; Chen, Z. Combining an antagonistic yeast with harpin treatment to control postharvest decay of kiwifruit. *Biol. Control* **2015**, *89*, 61–67. [[CrossRef](#)]
18. Yu, T.; Yu, C.; Chen, F.; Sheng, K.; Zhou, T.; Zunun, M.; Abudu, O.; Yang, S.; Zheng, X. Integrated control of blue mold in pear fruit by combined application of chitosan, a biocontrol yeast and calcium chloride. *Postharvest Biol. Technol.* **2012**, *69*, 49–53. [[CrossRef](#)]
19. Abouraicha, E.; El Alaoui-Talibi, Z.; El Boutachfai, R.; Petit, E.; Courtois, B.; Courtois, J.; El Modafar, C. Induction of natural defense and protection against *Penicillium expansum* and *Botrytis cinerea* in apple fruit in response to bioelicitors isolated from green algae. *Sci. Hort.* **2015**, *181*, 121–128. [[CrossRef](#)]
20. Stadnik, M.J.; Freitas, M.B.D. Algal polysaccharides as source of plant resistance inducers. *Trop. Plant Pathol.* **2014**, *39*, 111–118. [[CrossRef](#)]
21. Siah, A.; Magnin-Robert, M.; Randoux, B.; Choma, C.; Rivière, C.; Halama, P.; Reignault, P. Natural agents inducing plant resistance against pests and diseases. In *Natural Antimicrobial Agents*; Merillon, J.M., Riviere, C., Eds.; Springer: Cham, Switzerland, 2018; ISBN 978-3-319-67045-4.
22. Araujo, L.; Stadnik, M.J. Cultivar-specific and ulvan-induced resistance of apple plants to *Glomerella* leaf spot are associated with enhanced activity of peroxidases. *Acta Sci. Agron.* **2013**, *35*, 287–293. [[CrossRef](#)]
23. de Freitas, M.B.; Stadnik, M.J. Ulvan-induced resistance in *Arabidopsis thaliana* against *Alternaria brassicicola* requires reactive oxygen species derived from NADPH oxidase. *Physiol. Mol. Plant Pathol.* **2015**, *90*, 49–56. [[CrossRef](#)]
24. Cluzet, S.; Torregrosa, C.; Jacquet, C.; Lafitte, C.; Fournier, J.; Mercier, L.; Salamagne, S.; Briand, X.; Esquerre-Tugaye, M.T.; Dumas, B. Gene expression profiling and protection of *Medicago truncatula* against a fungal infection in response to an elicitor from green algae *Ulva* spp. *Plant Cell Environ.* **2004**, *27*, 917–928. [[CrossRef](#)]
25. El Modafar, C.; Elgadda, M.; El Boutachfai, R.; Abouraicha, E.; Zehhar, N.; Petit, E.; El Alaoui-Talibi, Z.; Courtois, J. Induction of natural defense accompanied by salicylic acid-dependent systemic acquired resistance in tomato seedlings in response to bioelicitors isolated from green algae. *Sci. Hort.* **2012**, *138*, 55–63. [[CrossRef](#)]

26. Rivas-Garcia, T.; Hernandez-Montiel, L.G.; Murillo-Amador, B.; Nieto-Garibay, A.; Chiquito-Contreras, R.G.; Rincon-Enriquez, G. Identification and characterization of *Fusarium* spp. from muskmelon in northwest Mexico. *Biotechnia* **2018**, *20*, 71–75.
27. Zhou, Y.; Zhang, L.; Zeng, K. Efficacy of *Pichia membranaefaciens* combined with chitosan against *Colletotrichum gloeosporioides* in citrus fruits and possible modes of action. *Biol. Control* **2016**, *96*, 39–47. [[CrossRef](#)]
28. Mattiuz, B.H.; Ducamp-Collin, M.N.; Mattiuz, C.F.M.; Vigneault, C.; Marques, K.M.; Sagoua, W.; Montet, D. Effect of propolis on postharvest control of anthracnose and quality parameters of 'Kent' mango. *Sci. Hort.* **2015**, *184*, 160–168. [[CrossRef](#)]
29. Levy, Y.; Benderly, M.; Cohen, Y.; Gisi, U.; Bassand, D. The joint action of fungicides in mixtures: Comparison of two methods for synergy calculation. *EPPO Bulletin* **1986**, *16*, 651–657. [[CrossRef](#)]
30. Blackburn, D.; Shapiro-Ilan, D.I.; Adams, B.J. Biological control and nutrition: Food for thought. *Biol. Control* **2016**, *97*, 131–138. [[CrossRef](#)]
31. Cao, S.; Yang, Z.; Hu, Z.; Zheng, Y. The effects of the combination of *Pichia membranaefaciens* and BTH on controlling of blue mould decay caused by *Penicillium expansum* in peach fruit. *Food Chem.* **2011**, *124*, 991–996. [[CrossRef](#)]
32. Sharma, R.R.; Singh, D.; Singh, R. Biological control of postharvest diseases of fruits and vegetables by microbial antagonists: A review. *Biol. Control* **2009**, *50*, 205–221. [[CrossRef](#)]
33. Meng, X.H.; Qin, G.Z.; Tian, S.P. Influences of preharvest spraying *Cryptococcus laurentii* combined with postharvest chitosan coating on postharvest diseases and quality of table grapes in storage. *LWT-Food Sci. Technol.* **2010**, *43*, 596–601. [[CrossRef](#)]
34. De Costa, D.M.; Gunawardhana, H.M.D.M. Effects of sodium bicarbonate on pathogenicity of *Colletotrichum musae* and potential for controlling postharvest diseases of banana. *Postharvest Biol. Technol.* **2012**, *68*, 54–63. [[CrossRef](#)]
35. Araújo, L.; Gonçalves, A.E.; Stadnik, M.J. Ulvan effect on conidial germination and appressoria formation of *Colletotrichum gloeosporioides*. *Phytoparasitica* **2014**, *42*, 631–640. [[CrossRef](#)]
36. Gonçalves, A.E.; Stadnik, M.J. Interferência de ulvana no desenvolvimento e melanização de apressórios de *Colletotrichum gloeosporioides*. *Trop. Plant Pathol.* **2012**, *37*, 431–437. [[CrossRef](#)]
37. de Freitas, M.B.; Ferreira, L.G.; Hawerth, C.; Duarte, M.E.R.; Nosedá, M.D.; Stadnik, M.J. Ulvan induce resistance against plant pathogenic fungi independently of their sulfation degree. *Carbohydr. Polym.* **2015**, *133*, 384–390. [[CrossRef](#)] [[PubMed](#)]
38. Paulert, R.; Talamini, V.; Cassolato, J.E.F.; Duarte, M.E.R.; Nosedá, M.D.; Smania, A.; Stadnik, M.J. Effects of sulfated polysaccharide and alcoholic extracts from green seaweed *Ulva fasciata* on anthracnose severity and growth of common bean (*Phaseolus vulgaris* L.). *J. Plant Dis. Protect.* **2009**, *116*, 263–270. [[CrossRef](#)]
39. Borsato, L.C.; Di Piero, R.M.; Stadnik, M.J. Mecanismos de defesa eliciados por ulvana contra *Uromyces appendiculatus* em três cultivares de feijoeiro. *Trop. Plant Pathol.* **2010**, *35*, 318–322. [[CrossRef](#)]
40. Jaulneau, V.; Lafitte, C.; Jacquet, C.; Fournier, S.; Salamagne, S.; Briand, X.; Esquerre-Tugaye, M.T.; Dumas, B. Ulvan, a sulfated polysaccharide from green algae, activates plant immunity through the jasmonic acid signaling pathway. *J. Biomed. Biotechnol.* **2010**. [[CrossRef](#)] [[PubMed](#)]
41. Paulert, R.; Ebbinghaus, D.; Urllass, C.; Moerschbacher, B.M. Priming of the oxidative burst in rice and wheat cell cultures by ulvan, a polysaccharide from green macroalgae, and enhanced resistance against powdery mildew in wheat and barley plants. *Plant Pathol.* **2010**, *59*, 634–642. [[CrossRef](#)]
42. Janisiewicz, W.J.; Korsten, L. Biological control of postharvest diseases of fruits. *Annu. Rev. Phytopathol.* **2002**, *40*, 411–441. [[CrossRef](#)] [[PubMed](#)]
43. Grzegorzczak, M.; Żarowska, B.; Restuccia, C.; Cirvilleri, G. Postharvest biocontrol ability of killer yeasts against *Monilinia fructigena* and *Monilinia fructicola* on stone fruit. *Food Microbiol.* **2017**, *61*, 93–101. [[CrossRef](#)] [[PubMed](#)]
44. Rivas-Garcia, T.; Hernandez-Montiel, L.G.; Murillo-Amador, B.; Nieto-Garibay, A.; Chiquito-Contreras, R.G.; Rincon-Enriquez, G. Enhanced biocontrol of fruit rot on muskmelon by marine *Debaryomyces hansenii* and *Stenotrophomonas rhizophila* and their potential modes of action. *Postharvest Biol. Technol.* in press.
45. Pech, J.C.; Bouzayen, M.; Latché, A. Climacteric fruit ripening: Ethylene-dependent and independent regulation of ripening pathways in melon fruit. *Plant Sci.* **2008**, *175*, 114–120. [[CrossRef](#)]

46. Gupta, N.; Jain, S.K. Storage behavior of mango as affected by postharvest application of plant extracts and storage conditions. *J. Food Sci. Technol.* **2014**, *51*, 2499–2507. [[CrossRef](#)] [[PubMed](#)]
47. Zhang, J.; Bruton, B.D.; Biles, C.L. *Fusarium solani* endo-polygalacturonase from decayed muskmelon fruit: Purification and characterization. *Physiol. Mol. Plant Pathol.* **1999**, *54*, 171–186. [[CrossRef](#)]
48. Leslie, J.F.; Summerell, B.A. *The Fusarium Laboratory Manual*, 2nd ed.; Blackwell Pub.: Ames, IA, USA, 2006; p. 387.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).