

HIF-1-dependent induction of $\beta 3$ adrenoceptor: evidence from the mouse retina

Rosario Amato^{1,*}, Francesco Pisani^{2,*}, Emiliano Laudadio^{3,*}, Maurizio Cammalleri¹, Martina Lucchesi¹, Silvia Marracci¹, Luca Filippi⁴, Roberta Galeazzi⁵, Maria Svelto^{2,6,7}, Massimo Dal Monte^{1,§,§} and Paola Bagnoli^{1,§}

¹ Department of Biology, University of Pisa, Pisa, Italy; rosario.amato@biologia.unipi.it (R.A.); maurizio.cammalleri@unipi.it (M.C.); martina.lucchesi@student.unisi.it (M.L.); silvia.marracci@unipi.it (S.M.); massimo.dalmonate@unipi.it (M.D.M.); paola.bagnoli@unipi.it (P.B.)

² Department of Biosciences, Biotechnologies and Biopharmaceutics, University of Bari Aldo Moro, Bari, Italy; francesco.pisani@uniba.it (F.P.)

³ Department of Materials, Environmental Sciences and Urban Planning, Polytechnic University of Marche, Ancona, Italy; e.laudadio@staff.univpm.it (E.L.)

⁴ Department of Clinical and Experimental Medicine, Division of Neonatology and NICU, University of Pisa, Pisa, Italy; luca.filippi@unipi.it (L.F.)

⁵ Department of Life and Environmental Sciences, Polytechnic University of Marche, Ancona, Italy; r.galeazzi@staff.univpm.it (R.G.)

⁶ Institute of Biomembranes and Bioenergetics, National Research Council, Bari, Italy; maria.svelto@uniba.it (M.S.)

⁷ National Institute of Biostructures and Biosystems (INBB), Rome, Italy; maria.svelto@uniba.it (M.S.)

* Correspondence: contributing authors

§ Equal contributing senior authors

§ Correspondence: massimo.dalmonate@unipi.it (M.D.M.); Tel.: +39-050-2211426

Supplementary information




Figure S1. Close superimposition of the heterodimeric interface from 4zpr (orange) with the HIF1 α homodimer model (green): full structure and focus on DNA binding domain (boxes).