

Review

Genetic, Genomics, and Responses to Stresses in Cyanobacteria: Biotechnological Implications

Corinne Cassier-Chauvat , Victoire Blanc-Garin and Franck Chauvat * 

Institute for Integrative Biology of the Cell (I2BC), Université Paris-Saclay, CEA, CNRS, 91198 Gif-sur-Yvette, France; corinne.cassier-chauvat@cea.fr (C.C.-C.); victoire.blanc-garin@cea.fr (V.B.-G.)
* Correspondence: franck.chauvat@cea.fr; Tel.: +33-1-69-08-78-11

Abstract: Cyanobacteria are widely-diverse, environmentally crucial photosynthetic prokaryotes of great interests for basic and applied science. Work to date has focused mostly on the three non-nitrogen fixing unicellular species *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002, which have been selected for their genetic and physiological interests summarized in this review. Extensive “omics” data sets have been generated, and genome-scale models (GSM) have been developed for the rational engineering of these cyanobacteria for biotechnological purposes. We presently discuss what should be done to improve our understanding of the genotype-phenotype relationships of these models and generate robust and predictive models of their metabolism. Furthermore, we also emphasize that because *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002 represent only a limited part of the wide biodiversity of cyanobacteria, other species distantly related to these three models, should be studied. Finally, we highlight the need to strengthen the communication between academic researchers, who know well cyanobacteria and can engineer them for biotechnological purposes, but have a limited access to large photobioreactors, and industrial partners who attempt to use natural or engineered cyanobacteria to produce interesting chemicals at reasonable costs, but may lack knowledge on cyanobacterial physiology and metabolism.

Keywords: *Synechocystis* PCC 6803; *Synechococcus* PCC 7942; *Synechococcus* PCC 7002; biodiversity; genotype-phenotype relationships; conjugation; transformation; RSF1010 plasmid; neutral cloning sites



Citation: Cassier-Chauvat, C.; Blanc-Garin, V.; Chauvat, F. Genetic, Genomics, and Responses to Stresses in Cyanobacteria: Biotechnological Implications. *Genes* **2021**, *12*, 500. <https://doi.org/10.3390/genes12040500>

Academic Editors: Denis Baurain and Luc Cornet

Received: 26 January 2021
Accepted: 25 March 2021
Published: 29 March 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Cyanobacteria are ancient Gram-negative prokaryotes that perform oxygenic photosynthesis and are phylogenetically close to recently discovered non-photosynthetic bacteria termed Melainabacteria and Sericytochromatia [1,2]. Cyanobacteria are regarded as the producer of the atmospheric oxygen (O₂) of Earth [3] and the ancestors of the plant chloroplast [4]. Cyanobacteria capture solar energy at high efficiencies (3–9%) [5] to power up their efficient photoautotrophic metabolism, which fixes huge amounts of inorganic carbon (CO₂, NaHCO₃, and Na₂CO₃) and nitrogen (N₂, NH₄, NO₂, NO₃, or urea) [6,7], into an enormous biomass [8] that supports a large part of the food chain.

By colonizing aquatic ecosystems (fresh, brackish, and marine waters) and soils (including deserts) of our planet, cyanobacteria are inevitably exposed to multiple stresses such as solar ultraviolet radiations and variations in light intensity and quality, inorganic-nutrients availabilities, temperatures (high and low), salinity, pH (acidic and basic), drought, and pollutants (herbicides and heavy-metals). In addition, cyanobacteria are involved in numerous interactions with competitors, predators, or symbiotic hosts [9]. Consequently, it is not surprising that cyanobacteria have evolved as a widely diverse organisms, which are of high interest for basic and applied research [10]. They display various metabolisms and morphologies [11,12], and numerous species can differentiate cells, akinetes (spores) and/or heterocysts, which are dedicated to growth or survival under adverse conditions [13,14].

Thus, cyanobacteria are good model organisms to study the impact of environmental conditions on the physiology, metabolism, and morphology of microbial cells. Furthermore, cyanobacteria synthesize a wide variety of bioactive metabolites (Figure 1), many of which being of interest for human health [15–18], and they are regarded as promising cell factories for the production of chemicals (fuels and biodegradable bioplastics) from highly abundant natural resources: solar energy, water (not necessarily potable), CO₂, and minerals, thanks to their active photosynthesis and the synthetic biology tools of model species [10,19,20].




Figure 1. Cyanobacteria can synthesize a wealth of biotechnologically interesting products.

Most of our knowledge on cyanobacteria came from studying the three non-nitrogen fixing, unicellular cyanobacteria *Synechocystis* PCC 6803, *Synechococcus* PCC 7942 (formerly named “*Anacystis nidulans*”) and *Synechococcus* PCC 7002 (formerly named “*Agmenellum quadruplicatum* PR6”) that are (i) straightforward to culture under laboratory conditions, (ii) amenable to genetic manipulation, and (iii) freezable for long-term storage. In this review, we will summarize the genetic and physiological properties of these models, emphasizing on their tolerance to stresses, and the recent progress in their genetics. We will also put forward that in spite of more than three decades of intensive research the genomes of these models still contain a large number of genes and small RNAs (sRNAs) of unknown function. Furthermore, many of the genes annotated “by sequence analogy” with those genes characterized in intensively studied, non-photosynthetic, models such as *Escherichia coli* might in fact have a different function in cyanobacteria. This situation makes comparative genomics and metabolic modeling difficult. Consequently, we will discuss that to better understand and exploit the wide biodiversity of cyanobacteria, strong efforts should be put in large-scale analysis of genes and sRNAs functions in model cyanobacteria. Finally, we will emphasize that we need to identify and thoroughly study new cyanobacteria endowed with natural properties of interest for basic or applied researches, and test whether the synthetic biology tools developed for model strains can be used to facilitate the engineering of these newly identified cyanobacteria so as to turn their promises into industrial realities.

2. Cyanobacteria Being Inevitably Exposed to Photo-Oxidative Stress Have Developed the Evolutionary-Conserved Glutathione System

Having evolved the oxygenic photosynthesis [21], cyanobacteria were the first organisms to be exposed to the toxic reactive oxygen species (ROS): singlet oxygen ($^1\text{O}_2$), superoxide anion (O_2^-), hydrogen peroxide (H_2O_2), and hydroxyl radical (OH) [22] that are generated when the light-driven electron transport exceeds what is needed for nutrients assimilation [23]. Accordingly, cyanobacteria represent a major source of ROS in natural aquatic environments [24].

To cope with ROS, cyanobacteria have developed the glutathione system [3]. It comprises the glutathione tripeptide (gamma-glutamyl-cysteinyl-glycine) and numerous glutathione-dependent enzymes [25], which have been conserved during evolution [26–29]. Glutathione is abundant in all organisms (concentration ranging from 0.1 to about 10 mM) where it plays pleiotropic roles in cell life and resistance to stresses [25,30]. Supporting the notion that cyanobacteria have evolved the glutathione system, the glutathione synthesis enzyme GshB and several glutathione transferases of the model cyanobacterium *Synechocystis* PCC 6803 were shown to play a prominent role in the tolerance to O_2 and photo-oxidative stress [31,32].

Glutathione occurs under two redox forms: the reduced (GSH) and the oxidized, dimeric (GSSG) forms. GSH, the major form, maintains the intracellular compartment in a reduced state. It provides electrons to the GSH-dependent enzymes glutaredoxins and glutathione transferases that operate in defenses against photo-oxidative stress and heavy metal pollutants [32–34]. These toxic-only metals inactivate many enzymes, by reacting with the thiol-group of their cysteines and/or displacing their normal metal cofactors [35]. To protect cells from metals, GSH can chelate them outside the cells under the form of large complexes that cannot penetrate into the cells [36]. GSH can also form a Fe(II)GSH complex, which likely supplies Fe for the synthesis of the Fe cofactor or (Fe-S) clusters of many metalloenzymes operating in photosynthesis, respiration, or other cellular functions [30,37,38].

Oxidative stress can also generate disulfides bridges between two cysteinyl-residues of either the same or two proteins, or a protein and a molecule of glutathione, which otherwise affect protein activity [33,39]. The glutathionylation/deglutathionylation process has been studied in the best studied unicellular cyanobacterium *Synechocystis* PCC 6803 where about 400 proteins can be glutathionylated in vitro [40]. These proteins participate in a wide range of cellular and metabolic processes including carbon and nitrogen metabolisms, cell division, stress responses and hydrogen production [40]. The glutathionylation/deglutathionylation control of the antioxidant peroxiredoxin II, the mercuric reductase, the AbrB2 transcription factor, and the metabolic enzyme 3-phosphoglycerate dehydrogenase was confirmed by biochemical studies with the purified recombinant proteins [34,40,41]. Attesting its importance for living organisms, the redox glutathionylation/deglutathionylation process and the glutathione transferase and glutaredoxin enzymes involved in its control have been conserved during evolution from cyanobacteria to plants and human [26,29,40,42,43].

Upon cell detoxication, GSH is oxidized for the glutathione disulfide form, GSSG, which can be reduced back to GSH by various factors, such as the NADPH-using glutathione reductase enzyme (GR), which occurs in most cyanobacteria [25] and other aerobic organisms [27,44], except the model cyanobacteria *Synechocystis* PCC 6803 [45] and *Synechococcus* PCC 7002 [25].

3. Cyanobacteria Possess Widely Diverse Genomes That Contain a Wealth of Unknown Genes and Poorly Characterized Multigene Families

The first articles reporting the sequence of an entire bacterial genome (*Hemophilus influenzae*, 1,830,137 bp; *Mycoplasma genitalium*, 580,070 bp) appeared in 1995 [46,47]. Shortly thereafter, the sequence of the 3,573,470-bp chromosome of the best studied unicellular cyanobacterium *Synechocystis* PCC 6803 was reported [48]. Currently, about 2000 complete or draft sequences of cyanobacterial genomes are accessible in public data bases such

as the DOE joint genome institute (<https://genome.jgi.doe.gov/portal/>, accessed on 15 March 2021) and the Microbial Genome Database for Comparative Analysis (<http://mbgd.genome.ad.jp/>, accessed on 15 March 2021). This number is steadily increasing in the frame of metagenomic analyses [49–52].

The genome of cyanobacteria differs in size, ranging from 1.44 Mbp [53] to 12.07 Mbp [54], and GC content (30–60%), probably as a result from gene gains and losses [55] transferred by plasmids, insertion sequences [56], and/or cyanophages [57–59]. Cyanophages infecting marine cyanobacteria contain genes important for photosynthesis, which could facilitate their transfer among marine cyanobacteria [60]. Instances of horizontal gene transfer are inferred on the basis of observations of patchy phylogenetic distribution and/or atypical nucleotide composition of several genes [55,59].

All cyanobacteria possess a circular chromosome and, depending on the species, a linear chromosome (rarely) and/or one to several circular plasmids (frequently), excepted the marine species UCYN-A, *Prochlorococcus* and *Synechococcus*, which have a small chromosome (1.44–2.7 Mbp), but no plasmids [53,61]. Some cyanobacterial plasmids contain genes likely involved in important processes, such as cysteine biosynthesis [62], thermal tolerance [63], resistance to metal stress [64], or DNA repair [58]. The marine cyanobacterium *Acarvochloris marina* MBIC1101, which has a 6.5 Mbp circular chromosome, has 25% of its genes located on its nine large (≥ 100 kbp) single-copy plasmids [65].

An unusual genome organization was observed in two strains of the unicellular genus *Cyanothece*. The genome of *Cyanothece* ATCC51142 (5.46 Mbp) harbors two chromosomes (one circular of 4.39 Mbp; and one linear of 0.4 Mbp) and four plasmids (10–39 kbp) [66,67]. *Cyanothece* PCC 7822 has a genome of 7.84 Mbp that comprises one circular chromosome (6.09 Mbp), three linear chromosomal elements (880, 474, and 14 kbp) and three plasmids [67]. These findings are interesting because a few bacteria possess a linear chromosome, such as the (non-photosynthetic) soil bacteria of the genus *Streptomyces*, which synthesize a wide variety of bioactive secondary metabolites [68], like cyanobacteria [16,17,69]. In *Streptomyces*, the linear chromosome exhibits a remarkable genetic organization with grossly a central conserved region flanked by variable chromosomal arms [70]. Based on these observations, it would be interesting in the future to compare the plasticity of linear versus circular chromosomes of cyanobacteria.

4. Interest and Current Limitations of Comparative Genomics

The comparative analysis of cyanobacterial genomes allows to determine which genes are present in any particular genome and which ones are absent. These findings serve to determine the pan-genome, which describes the entire genes set of all cyanobacteria analyzed. It includes (i) a core-genome, which comprises the sets of genes that exist in each of the strains analyzed [71] and (ii) the dispensable-genome, which comprises the genes present in a subset of the studied strains and species-specific genes [72]. For such analyses, strains can be selected on the basis of their natural habitats [73], their physiology and/or morphology [74], and/or their phylogenetic position determined by 16S-rRNA sequence comparison [75]. Such comparative genome analyses revealed habitat-specific (or heterocyst-specific) cyanobacterial proteins [72]. It was also revealed that cyanobacteria occupy a unique position among prokaryotes as a hub between anaerobes and obligate aerobes [76] and that the earliest cyanobacteria were small and unicellular, while filamentous forms appeared shortly afterwards [77]. Comparative genome studies were also used to discuss plastid evolution [78]. When applied to two closely related cyanobacteria, comparative genomics can be used to predict the behavior of one strain relative to the other one. As a rare example of such comparison, *Synechocystis* PCC 6803 was found to be more resistant to zinc excess than *Synechocystis* PCC 6714 as it was predicted [79]. Comparative genomic information is also crucial for genome-based reconstruction of an organism's metabolism as done for *Synechocystis* PCC 6803 to predict which metabolic reaction or pathway should be engineered to increase the production of biotechnologically interesting chemicals [80–85].

4.1. Danger of Genome Annotation Based Only on Sequence Comparison

Generating robust and accurate genome-scale models (GSMs) is an iterative process dependent on the expansion and updating of draft simulation models with available experimental data obtained after measurement of fluxes and metabolic pool sizes [80,86,87]. However, GSMs are largely based on gene function predicted “by sequence analogy” with genes characterized in other model organisms, which may behave differently in the studied organism [88]. In *Synechocystis* PCC 6803, the best studied model, only ca. 1050 coding sequences (~30%) have assigned functions [80,89], which may turn out to differ from those determined in well-studied heterotrophic models. For examples, the *Synechocystis* PCC 6803 LexA protein regulates the expression of genes involved in carbon assimilation [90], not DNA repair as it occurs in *E. coli*. Furthermore, the NAD(P)H dehydrogenase transcription factor (NdhR, Sll1594), homologous to the proteobacterial regulators of the Calvin–Benson–Bassham metabolic pathway (CbbR, hereafter referred to as Calvin cycle), does not regulate the transcription of Calvin cycle genes [91]. Instead, NdhR regulates numerous genes involved in the transport and assimilation of inorganic carbon (CO₂ and bicarbonate) [6,92,93].

In addition, a single cyanobacterial protein can have more than one function. The fructose-1,6/sedoheptulose-1,7-bisphosphatase (FBP/SBPase) operates in both the Calvin cycle and gluconeogenesis [94]. The CrtLdiox protein is also dual-function enzyme with both lycopene cyclase and dioxygenase activity [95]. The IsiA protein is active in both the light-harvesting ability of PSI and the photoprotection of PSII [96]. The AgrE enzyme catalyzes two sequential reactions in arginine catabolism, in converting arginine to ornithine, and then ornithine into proline [97,98]. The ApalaDH enzyme transforms pyruvate to L-alanine, L-alanine back to pyruvate, and glyoxylate to glycine [99]. The sucrose-synthesis SPS enzyme has sucrose-phosphate synthase and sucrose-phosphate phosphatase activities [100]. The KatG enzymes have both catalase and peroxidase activities, which are also capable of oxidizing chloride, bromide, and iodide compounds [101,102].

4.2. Identification of Essential Genes at the Level of a Whole Genome for a Better Understanding of the Genotype-Phenotype Relationships

To use genomic data for reconstruction of an organism’s metabolism it is also important to identify the comprehensive set of genes that are essential to cell growth in well-defined conditions. This has been addressed using transposon mutagenesis in *Synechocystis* PCC 6803 [103,104] and *Synechococcus* PCC 7942 [105–108]. For the same purpose, the construction of a whole genome library of gene insertion plasmids has been undertaken in *Synechocystis* PCC 6803 [7]. These works will certainly contribute to decrease the high number of proteins with still unknown function. In *Synechocystis* PCC 6803, transposon mutagenesis has been useful to study (i) photosynthesis [109–111], (ii) the transport of CO₂ or bicarbonate [112,113], (iii) the production of the poly-3-hydroxybutyrate (PHB) biodegradable bioplastic [114], and (iv) the NADPH:plastoquinone oxidoreductase complex operating in plastoquinone reduction and cyclic electron transfer (CET) around photosystem I [115–121]. Transposon mutagenesis has been also useful in *Synechococcus* PCC 7942 to identify a gene in fatty acid production [122].

4.3. Importance of Deciphering the Selectivity/Redundancy of Multiple Gene Families

To generate robust and predictive model of the metabolism of cyanobacteria, it is also important to unravel the redundancy/selectivity of multigene families, such as those that code for the stress-responsive redox proteins ferredoxins, glutaredoxins, and glutathione transferase.

Ferredoxins (Fed), are small (acidic) proteins present in most organisms. They use their iron-sulfur cluster (Fe-S) to distribute electrons to various metabolic pathways involved in nutrients assimilation [123]. The best studied cyanobacterium *Synechocystis* PCC 6803 possess nine Feds. The Fed1-6 proteins possess a (2Fe-2S) center, while Fed7 harbors a (4Fe-4S) cluster. In contrast both Fed8 and Fed9 have two clusters: (3Fe-4S) (4Fe-4S)

and (4Fe-4S) (4Fe-4S), respectively. The highly abundant Fed1 protein, essential to photosynthesis [124,125], is encoded by a light-inducible gene [126]. The low-abundant Fed2–Fed9 proteins are encoded by stress-responsive genes (light, carbon, herbicides, or heavy metals), which have differential importance (crucial/dispensable) for the photoautotrophic growth and/or the resistance to stresses [45,123,125,127]. Fed1, Fed7, and Fed9 participate in a ferredoxin–glutaredoxin–thioredoxin crosstalk pathway that operates in the protection against oxidative and metal stresses [45]. Fed7, but not Fed9, interacts with a DnaJ-like protein, an interaction that has been strengthened in photosynthetic eukaryotes in the form of a Fed7–DnaJ fusion protein [123]. Fed7 also has a regulatory role under photooxidative stress [128]. Conversely, Fed9, but not Fed7, interacts with the Flv3 flavodiiron protein involved in the photoreduction of O₂ to H₂O [123]. Other ferredoxin partners [129,130] should be studied to better understand the selectivity/redundancy of ferredoxins in *Synechocystis* PCC 6803. Attesting their importance for the life of cyanobacteria, the *fed* genes were highly conserved in cyanobacteria. For example, the other models *Synechococcus* PCC 7002 and *Synechococcus* PCC 7942 have, respectively, nine and six *fed* genes, while the symbiotic strain *Acaryochloris marina* MBIC11017 endowed with a large genome (8.36 Mb) possesses sixteen *fed* genes [123].

Like ferredoxins, the evolutionary-conserved enzymes glutaredoxins (Grxs) are widely distributed in cyanobacteria [25,131]. Grxs use electrons provided by glutathione (GSH), or the thioredoxin reductase enzyme [45,132], to reduce the oxidative-stress-generated disulfides occurring between two cysteinyl-residues of either the same or two proteins, or a protein and a molecule of glutathione, which otherwise affect protein activity [33,39]. The dithiol Grxs, which possess a CXXC redox center (C and X stand for cysteine and any other amino acid, respectively), catalyze the reduction in protein disulfides or glutathione–protein mixed disulfides (the latter activity is named “deglutathionylation”). The monothiol Grxs, which have a CXXS redox center (S stands for serine), operates in iron sensing and trafficking the biogenesis of iron–sulfur clusters of proteins and deglutathionylation [30]. All cyanobacteria possess a monothiol Grx-encoding gene and a variable number of dithiol Grx genes [25]. *Synechocystis* PCC 6803 possesses three Grxs, which are all dispensable to the standard photoautotrophic growth [25,133]. The dithiol enzymes Grx1 and Grx2, which can interact together [45], operate in tolerance to H₂O₂ [45,133], arsenate [134,135], selenate [45], mercury, and uranium [34]. Grx1 can reactivate the oxidized (glutathionylated) form of the mercuric reductase enzyme by catalyzing its deglutathionylation [34]. The monothiol Grx3 (CGFS redox center) enzyme forms a homodimer bridged by a glutathione-ligated (2Fe-2S) cluster [26]. This feature, which has been conserved in Grx3 orthologs from cyanobacteria to plants and mammals, likely operates in Fe sensing and distribution of (2Fe-2S) cluster [26,30,136].

Another important multiple genes family encodes the evolutionary-conserved glutathione transferase (GT) enzymes, which can conjugate glutathione (GSH) on diverse toxics (oxidants, chemicals, and heavy metals) thereby generating water-soluble complexes that can then be degraded or excreted out of the cell [33,137,138]. GST also operate in the glutathionylation/deglutathionylation process [42,43]. Glutathione-S-transferases (GST) are commonly divided in three different families: (i) cytosolic GSTs (the largest family), (ii) mitochondrial GSTs, and (iii) microsomal (membranous) GSTs designated as MAPEGs (membrane-associated protein involved in ecosanoïd and glutathione metabolism) [139]. Little is known about GSTs in cyanobacteria though they are regarded as having evolved the oxygen-generating photosynthesis, as well as GSH and GSH-dependent enzymes to protect themselves against the toxic ROS (reactive oxygen species) massively produced by their active photosynthesis [3]. Based on phylogenetic tree analyses, 12 GST classes were identified in cyanobacteria [140,141]. Cyanobacterial GSTs have been studied mostly in *Synechocystis* PCC 6803, which possesses six GST, namely, Sll0067, Sll1147, Sll1545, Sll1902, Slr0236, and Slr0605 [142–144]. While the role of Sll1902 and Slr0605 is still unknown, Sll1545 and Slr0236 were shown to operate in the protection against photo-oxidative stress triggered by high light or H₂O₂ [32]. Sll1147 plays a prominent role in tolerance to mem-

brane stresses triggered by heat, cold, and lipid peroxidation [29]. Sll0067 operates in the protection against methylglyoxal (MG), a toxic metabolite by-product of the catabolism of sugars, lipids, and amino-acids, which causes diabetes in human. Sll0067 catalyzes the conjugation of GSH with MG, the first step in MG detoxification catalyzed by the glyoxalase enzymes [145].

5. A Few Cyanobacteria Are Currently Amenable to Gene Manipulation, Leaving the Wide Biodiversity of Cyanobacteria Largely Unexplored

Historically, three non-nitrogen fixing unicellular cyanobacteria *Synechocystis* PCC 6803 (euryhaline strain), *Synechococcus* PCC 7942 (freshwater strain formerly named “*Anacystis nidulans*”) and *Synechococcus* PCC 7002 (marine strain formerly named “*Agmenellum quadruplicatum* PR6”) were chosen as model because they possess a simple morphology [146,147], a small genome (see above) and the important natural capability to be genetically transformed (see below). In addition, the freshwater strain *Anabaena* (*Nostoc*) PCC 7120 was also chosen as a genetically manipulable representative of filamentous (pluricellular) cyanobacteria [148]. This latter model is well used to analyze the differentiation of heterocysts, the cells dedicated to the fixation of atmospheric nitrogen, and their communication with vegetative cells [14], which are not the focus of the present review.

Although these four species are phylogenetically distant, they represent only a limited part of the wide biodiversity of cyanobacteria. This is evident from the comparison of the size and organization of their genomes (2.7–7.2 Mbp, absence of linear chromosome) with those of many other cyanobacteria (1.44–12M bp see above, presence of a linear chromosome in a few species).

New rapidly growing unicellular cyanobacteria have been recently described, namely, the freshwater strains *Synechococcus* UTEX 2973 [149], *Synechococcus* PCC 11801 [150], and *Synechococcus* PCC 11802 [151] and the marine strain *Synechococcus* PCC 11901 [80,152]. *Synechococcus* PCC 11901 tolerates high temperatures, up to 43 °C, which are non-optimal for *Synechococcus* UTEX 2973 [150,153], *Synechococcus* PCC 11801 [150,151], and *Synechococcus* PCC 7942 [154], and even detrimental to *Synechocystis* PCC 6803 [29,155]. *Synechococcus* PCC 11801 [150], *Synechococcus* PCC 11802 [151], and *Synechococcus* PCC 11901 [152] are naturally transformable, unlike *Synechococcus* UTEX2973 [150]. The latter strain was made transformable by cloning the *Synechococcus* PCC 7942 *pilN* gene encoding the Tfp pilus assembly protein [156].

Unfortunately, *Synechococcus* UTEX2973, *Synechococcus* PCC 11801, and *Synechococcus* PCC 11802 are very closely related to the freshwater model *Synechococcus* PCC 7942, in sharing, respectively, 99.8% and about 83% of genome identity with it [149–151,157]. Similarly, *Synechococcus* PCC 11901 shares 96.7% sequence identity with the marine model *Synechococcus* PCC 7002 [87,152]. Thus, future studies carried out with these new model strains will not significantly increase our understanding of the cyanobacterial biodiversity.

6. Properties of the Intensively Studied Unicellular Cyanobacteria *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

The three non-nitrogen fixing models *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002 have specific physiological properties that can influence their potential to serve as cell factories for biotechnological projects. *Synechocystis* PCC 6803 in being one of the few cyanobacteria capable to grow under photoautotrophic, mixotrophic, or heterotrophic conditions [147] has become the best studied cyanobacterium. It serves as model to study the photoautotrophic metabolism [6,7,93], the crucial importance of the carbon/nitrogen metabolic balance [6,93], and the responses to stresses [35]. Such investigations are also carried out, to a lesser extent, with *Synechococcus* PCC 7002 and *Synechococcus* PCC 7942. The latter model is well used to investigate cell division [158], the carbon/nitrogen metabolic crosstalk [93], the circadian rhythm [159], and the biogenesis of the CO₂-concentrating carboxysome organelle [160].

6.1. Genome Organization in the Intensively Studied Unicellular Cyanobacteria *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

Synechocystis PCC 6803, the best studied model, harbors a 3.57 Mbp circular chromosome [48], which has a copy number of about 10–22 per cell [161,162], and seven plasmids of sizes 2.3 [163,164], 2.4 [165], 5.2 [166], 44, 103, 106, and 120 kbp [167]. The ploidy of the three smaller plasmids 2.3–5.2 kbp was shown to be similar and increased from 3 to 8 copies per chromosome in cells reaching the stationary phase of growth. In contrast, the ploidy of the four larger plasmids, ranging from ~0.3 to 1.2 per chromosome depending on the studied plasmid, varied little with the growth phase [168]. The 5.2 kbp plasmid appeared to be dispensable to the standard photoautotrophic growth of *Synechocystis* PCC 6803 [169]. In contrast, large plasmids appeared to be essential (cells could not lose these plasmids), likely because they encode several toxin–antitoxin systems mediating plasmid maintenance [170,171] and they operate in beneficial functions such as potassium transport [172], copper resistance [173], and the metabolic tricarboxylic acid cycle [174].

The other model cyanobacteria *Synechococcus* PCC 7942 and *Synechococcus* PCC 7002 possess a circular chromosome of, respectively, 2.69 Mbp and 3.0 Mbp, each occurring at two to five copies per cell [175–177]. In addition, *Synechococcus* PCC 7942 has two plasmids of 7.84 [178] and 46.4 kbp [179], while *Synechococcus* PCC 7002 contains six plasmids [180] from 4.8 [181] to 186 kbp. The 7.84 kbp plasmid of *Synechococcus* PCC 7942 is not essential to its photoautotrophic growth [182,183]. In comparison, the well-studied filamentous cyanobacterium *Anabaena* (*Nostoc*) PCC 7120 has a 7.2 Mbp genome that comprises a circular chromosome of 6.41 Mbp and six plasmids of sizes 408.10, 186.614, 101.96, 55.41, 40.34, and 5.58 kbp [184].

6.2. Physiological Properties of *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002 and Biotechnological Implication

In agreement with them having different genome size and organization, the three model cyanobacteria also show different physiologies. *Synechococcus* PCC 7002, which has a doubling time of ~2.6 h [185], grows 2–7 times faster than *Synechococcus* PCC 7942 and *Synechocystis* PCC 6803 depending on the environmental conditions [149,183,186]. However, *Synechococcus* PCC 7002 requires vitamin B12 (cobalamin) to grow [146], unlike *Synechocystis* PCC 6803 and *Synechococcus* PCC 7942 [147]. Thus, the cost of vitamin B12 supplementation should be considered when *Synechococcus* PCC 7002 is to be used for biotechnological purposes requiring large-scale cultures. The vitamin B12 auxotrophy of *Synechococcus* PCC 7002 is due to the fact that it uses a cobalamin-dependent methionine synthase (MetH) for the synthesis of methionine, though it cannot synthesize cobalamin de novo. Recently, a cobalamin-independent methionine synthase *metE* gene from *Synechococcus* PCC 73109 was expressed in *Synechococcus* PCC 7002 to relieve its cobalamin auxotrophy [187], but this modified *Synechococcus* PCC 7002 sub-strain has been little employed yet.

All three model cyanobacteria are growing well on ammonium and nitrate, the usual nitrogen sources for cyanobacteria. In addition, both *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 can grow on urea (a frequent pollutant) as the sole nitrogen source [188–190], unlike *Synechococcus* PCC 7942 [190]. Furthermore, *Synechococcus* PCC 7002 and *Synechocystis* PCC 6803 are salt resistant (*Synechococcus* PCC 7002 is a marine strain), unlike the freshwater strain *Synechococcus* PCC 7942 [191]. Thus, *Synechococcus* PCC 7942 is not a suitable cell factory for future projects aiming at the photosynthetic production of chemicals in waters polluted by urea and/or salt to save the costs of potable waters.

As iron is an essential enzyme cofactor for oxygenic photosynthesis, cyanobacteria utilize multiple strategies to maintain iron levels within a desired range. One of them is the synthesis, export, and re-import of ferric ion chelators called siderophores. Both *Synechococcus* PCC 7942 and *Synechococcus* PCC 7002 can synthesize siderophore, unlike *Synechocystis* PCC 6803 that can only import the siderophore produced by other organisms [192]. These three model cyanobacteria have other metabolic differences. The RbcX chaperone operating in assembly of the CO₂-fixing RubisCO enzyme is essential in *Synechococcus* PCC 7002 [193], whereas it is dispensable in *Synechococcus* PCC 7942 [194] and *Synechocystis* PCC 6803 [195]. In addition, *Synechocystis* PCC 6803 has four flavodiiron proteins (Flv1–Flv4) [196], which

function as heterodimers Flv1/3 and Flv2/4. Flv1/3 catalyzes the NAD(P)H-driven reduction in oxygen to water on the acceptor side of PSI [196–198], while Flv2/4 operates both in the photoprotection of the photosystem II [199] and in an oxygen-dependent alternative electron flow [200]. Unlike *Synechocystis* PCC 6803, both *Synechococcus* PCC 7942 and *Synechococcus* PCC 7002 possess only Flv1/3, not Flv2/4 [201].

Finally, both *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 secrete extracellular polymeric substances mainly composed of exopolysaccharides, which act in the formation of biofilm and the protection against salt and metals stresses [202–205], unlike *Synechococcus* PCC 7942 that do not normally form biofilms [206].

6.3. Comparison of the Stress-Responsive Glutathione and DNA Repair Systems of the Model Cyanobacteria *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

Attesting the importance of the glutathione system for the tolerance to photo-oxidative stress and cell detoxication, most cyanobacteria possess the glutathione system. Its composition varies depending on the species and the environmental challenges they face. Most cyanobacteria have a glutathione reductase (GR) enzyme as *Synechococcus* PCC 7942, unlike both *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 [25,45]. Furthermore, *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 possess three glutaredoxins (Grxs) operating in tolerance to oxidative and metal stresses [34,45,133–135], whereas *Synechococcus* PCC 7942 has only two Grxs [25,131]. Another common feature shared by *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 is the fact that they both have an orange carotenoid protein (OCP) operating in photoprotection, whereas *Synechococcus* PCC 7942 has no OCP [207]. *Synechocystis* PCC 6803 and *Synechococcus* PCC 7942 have a MAPEG-type glutathione transferase enzyme operating in the tolerance to temperature stresses [29], which could occur during culture in open ponds, unlike *Synechococcus* PCC 7002.

Concerning the also important DNA repair system, which has been poorly studied in cyanobacteria, it is worth noting that the three model species have the following common and specific features. *Synechocystis* PCC 6803 possesses the *recD* gene, which is absent in *Synechococcus* PCC 7002 and *Synechococcus* PCC 7942 [58]. Furthermore, *Synechocystis* PCC 6803 has *recB* and *recJ*, which are duplicated in *Synechococcus* PCC 7002 and *Synechococcus* PCC 7942.

Synechocystis PCC 6803 and *Synechococcus* PCC 7942 have the *umuC* and *umuD* genes (*umuC* is duplicated in *Synechocystis* PCC 6803), unlike *Synechococcus* PCC 7002 that has no *umuC* and *umuD* genes [58]. Furthermore, *Synechocystis* PCC 6803 and *Synechococcus* PCC 7942 *recQ*, which is duplicated in *Synechococcus* PCC 7002.

Synechococcus PCC 7942 has *mutT* and *mutY*, whereas *Synechococcus* PCC 7002 lacks *mutT* and *Synechocystis* PCC 6803 lacks *mutY* [58].

Synechocystis PCC 6803 and *Synechococcus* PCC 7002, which can both use exogenous carbohydrates to accelerate their growth [6], possess *lexA*, which encodes a transcription regulator involved in carbon assimilation, not DNA repair [90]. In contrast, the obligate photoautotroph *Synechococcus* PCC 7942 has no *lexA* [58].

Synechocystis PCC 6803, *Synechococcus* PCC 7002, and *Synechococcus* PCC 7942 have also different features in the simplest DNA repair system, which removes only the base-modifying agent in one single step catalyzed by the AlkB demethylase, the Ogt alkyltransferase, and the Phr photolyase. All three model cyanobacteria possess *phr*, whereas *Synechocystis* PCC6803 has *alkB* but not *ogt*, *Synechococcus* PCC7942 has *ogt* (duplicated) but not *alkB*, and *Synechococcus* PCC7002 has neither *alkB* nor *ogt* [58].

6.4. Comparative Analysis of the Growth and Response to Stresses of the Three Model Cyanobacteria *Synechocystis* PCC 6803, *Synechococcus* PCC 7002, and *Synechococcus* PCC 7942

To better characterize the common and different physiological features exhibited by these three model cyanobacteria, it would be very interesting to perform simultaneous analyses and comparisons of their growth and tolerance to stresses in the same laboratories. This task is of high interest for not only basic science but also for biotechnological projects, which use of robust cell factories capable to withstand the environmental challenges

imposed by the industrial process and the possible toxicity of the intended products. So far, a very limited number of comparative studies have been carried out. *Synechocystis* PCC 6803 was shown to be more tolerant than *Synechococcus* PCC 7942 to UV (and gamma radiations) [90] and to the undecane hydrocarbon [208]. Reciprocally, *Synechococcus* PCC 7942 appeared to be more tolerant to chromate than *Synechocystis* PCC 6803, possibly because the sulfate transporters of *Synechococcus* PCC 7942 have lower affinity to chromate than those of *Synechocystis* PCC 6803 [209]. Furthermore, chromate generated more ROS (reactive oxygen species) in *Synechocystis* PCC 6803, as compared to *Synechococcus* PCC 7942, likely because *Synechocystis* PCC 6803 has intrinsic levels of superoxide dismutase, catalase and 2-Cys-peroxiredoxin than *Synechococcus* PCC 7942 [210]. Additionally, interestingly, *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 were reported to be more tolerant to ethanol than *Synechococcus* PCC 7942 [208,211].

Another important aspect to study is the genetic stability of the recombinant cyanobacteria generated for applied research. It has been reported that many engineered cyanobacteria appeared to be genetically unstable [58]. In one *Synechocystis* PCC 6803 recombinant strain, the instability was caused by the transposition of an IS5 insertion sequence leading to the inactivation of a regulatory gene [56]. Thus, it is of interest to note that the genomes of *Synechococcus* PCC 7942 and *Synechococcus* PCC 7002 are predicted to contain one and approximately 10 transposase encoding genes, whereas *Synechocystis* PCC 6803 possesses a single transposase gene.

6.5. As Observed in *Synechocystis* PCC 6803, the Sub-Strains of a Single Cyanobacterium Cultivated in Various Laboratories Can Have Different Behaviors

Another level of complexity is the un-surprising finding that a single cyanobacterium cultivated for some times in different laboratories tend to develop genetic and physiological differences from one laboratory to another. In the case of *Synechocystis* PCC 6803, originally isolated from a fresh water lake in California [146] and deposited in both the American Type Culture Collection (strain number ATCC 27184) and the Pasteur Culture Collection (strain number PCC 6803), it was reported that several sub-strains cultivated in different laboratories varied in genotypes and/or phenotypes. As compared to the original genome sequence [48], several specific mutations were identified in genes related to photosynthesis, transport, or motility [212–218]. Furthermore, variability in genome copy number was also observed [162]. Moreover, phenotypic variations were also reported including photosynthesis [217], cell size [219], motility [212,218], capability to grow on glucose as the carbon source [212,220], or resistance to temperature and salt (NaCl) stresses [221].

7. Genetic Characteristics of the Model Cyanobacteria *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

Shortly after their selection based on their robustness and simple (unicellular) morphologies [146,147] *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002 were found to be naturally competent for genetic transformation (see below).

7.1. *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002 Are Naturally Competent for Genetic Transformation

Natural competence for transformation of prokaryotes refers to their capability to take up DNA from their environment and incorporate it into their own genome. Originally, antibiotic resistant mutants, generated after UV or chemical mutagenesis, were selected, and their genomic DNA was isolated and used to transform wild type cells selecting for antibiotic resistance. Three naturally transformable species have emerged from such studies, namely, *Synechococcus* PCC 7942 [222], *Synechococcus* PCC 7002 (*Agmenellum quadruplicatum* PR6) [223,224], and *Synechocystis* sp. PCC 6803 [225]. In *Synechococcus* PCC 7942, the efficiency of transformation was found to decrease at pH lower than 7.0 or temperature ≥ 40 °C, two conditions being also unfavorable for growth. Furthermore, transformation was best effective with cells reaching the transition from the first to the second exponential phases of growth, supporting the notion that transformation depends upon the physio-

logical stage of the culture [183,226]. In addition, transformation appeared to be more efficient when expression of the transferred genes was allowed for 24 h on non-selective solid medium, prior to introducing the selective antibiotic underneath the agar of the plate for its slow diffusion toward the cells for gentle selection of the transformants [183]. Additionally, interestingly, the transformation to *Synechocystis* PCC6803 was reported to be strongly stimulated after the deletion of the exonuclease *recJ* gene [227]. Recently, the natural competence for transformation was shown to involve pilus appendages [228] and to be regulated by the circadian clock [229].

In *Synechococcus* PCC 7002, it was shown that single-stranded DNA cannot transform competent cells. Furthermore, cells in the stationary phase of growth or deprived of nitrogen or light before exposure to donor DNA tend to lose their competence for transformation. In contrast, significant improvement in transformation frequency were achieved by increasing the nitrate content of the culture medium or lowering the temperature from 39 °C (the optimal temperature for growth) to 30 °C before exposure to donor DNA [230].

Later, it was established in *Synechococcus* PCC 7942 [231], *Synechococcus* PCC 7002 [232], and *Synechocystis* PCC 6803 [161,163,233] that the donor DNA is integrated in the chromosome, or an endogenous plasmid (see below) of the recipient cells, through homologous recombination (double crossing-over or gene conversion) occurring on each side of the transferred DNA sequence in the region of homology between both the donor DNA and the recipient DNA. This process, similar to what had been described earlier in *Bacillus subtilis* [234], was extensively used for deleting cyanobacterial genes (or a part of them) through their targeted replacement by an easy selectable antibiotic resistance marker. Soon after its discovery, natural transformation of cyanobacteria was used in pioneering in vivo analyses of proteins involved in photosynthesis and/or stress resistance, which were performed with *Synechocystis* PCC 6803 [235–240], *Synechococcus* PCC 7942 [241–245], and *Synechococcus* PCC 7002 [246,247].

The molecular mechanism of natural transformation has been recently described in *Synechocystis* PCC 6803 [248]. First, DNA uptake from the environment is mediated by binding to the (type IV) pili appendages, which are also involved in cell adhesion and biofilm formation as well as twitching motility. During pilus retraction, the DNA is pulled into the periplasmic space, where one DNA strand is degraded while the other is translocated further across the cytoplasmic membrane by the Com (competence) proteins. The single-stranded DNA arrived in the cytoplasm is protected from nucleases and incorporated into the genome of recipient cells via homologous recombination.

7.2. Interest and Limitation of the Polyploidy of *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

In cyanobacteria, genomic modification is a time-consuming process because these organisms are polyploid. For example, *Synechococcus* PCC 7942, *Synechococcus* PCC 7002 and *Synechocystis* PCC 6803 harbor about 2–5 and 10–12 chromosome copies per cell [161,175–177]. Thus, to create a homoploid mutant, a segregation procedure must be applied to ensure that all chromosome copies in the transformants carry only the modified DNA. This requires multiple rounds of culture streaking in the presence of the selective antibiotic, which can last several weeks. However, the polyploidy of cyanobacteria is not a merely negative trait. It allows to study genes that are essential to cell life. In such cases, we obtain the corresponding heteroploid mutants, which possess both mutant and WT chromosome copies (with and without the studied vital gene). Such mutants survive because they retain a limited but sufficient amount of the studied essential protein, and they usually have a phenotype different from the wild-type strain. This difference often serves as a guide to infer a role of the studied crucial protein.

7.3. Utilization of Neutral Genome Sites for Gene Manipulation in *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

Transformation has been used extensively for the introduction of endogenous or heterologous genes into neutral chromosomal or plasmid sites, i.e., loci that can be disrupted

with no negative effect on cellular viability. Many neutral sites localized inside a dispensable gene or intergenic region have been identified in the chromosome or in an endogenous plasmid of *Synechocystis* PCC 6803 [163,249–252], *Synechococcus* PCC 7942 [253–255] and *Synechococcus* PCC 7002 [256–258]. These neutral cloning sites were frequently used for cloning and expression of endogenous, or heterologous, genes involved in cell metabolism, stress responses, or the engineering of recombinant strains for the photosynthetic production of chemicals (see Tables 1–3).

In *Synechocystis* PCC 6803, the frequently used neutral loci are (i) the gene *slr0168* or the intergenic region between *slr2030* and *slr2031*, which have no known function; (ii) the *cpcB* gene involved in the synthesis of the photosynthetic pigment phycocyanin; and (iii) the three *psbA* genes (Table 1). Although *psbA1* is not expressed [235,259], *psbA2* and *psbA3*, encoding the D1 protein subunit of the photosystem II, are expressed and dispensable, but they cannot be inactivated simultaneously [259]. The deletion of *psbA2* gene is compensated by an up-regulation of *psbA3* [260]. Neutral sites on the endogenous plasmids pCC5.2 and pCA2.4 have also been identified and evaluated for genetic integration and expression [251,261]. One neutral site on pCC5.2 was used for cloning the limonene synthase genes from *Mentha spicata* or *Citrus limon*, which directed the production of limonene [262]. Interestingly, the production of fluorescent proteins directed from neutral sites in pCC5.2 or pCA2.4 were, respectively, 14- or 100-fold higher than those observed after chromosomal integration [249,261,263,264]. These data are consistent with the finding that the endogenous small plasmids have a higher copy number than the chromosome [168]. These findings suggest that to be well-expressed genes should be cloned preferentially in a small endogenous plasmid than in the chromosome, but this assumption remain to be verified with other genes.

Table 1. Literature on the utilization of neutral chromosomal cloning sites in *Synechocystis* PCC 6803.

Neutral Site and Objective of the Gene Manipulation	References
<i>psbA1</i>: a silent gene	[235]
Photoproduction of extra bicarbonate transporters to increase biomass	[265]
Increase carbon import to improve growth	[266]
Photoproduction of isobutanol	[267]
<i>psbA2</i> (slr1311): gene encoding the D1 protein of the PSII	[259]
Photoproduction of zeaxanthin	[268]
Analysis of a thioredoxin-interacting LuxR-like regulator	[269]
Photoproduction of beta-caryophyllene	[270]
Photoproduction of D1 protein of the PSII	[271]
Photoproduction of polyhydroxybutyrate (PHB) biodegradable bioplastics	[272]
Photoproduction of isoprene	[273–277]
Photoproduction of lipids	[278]
Photoproduction of aromatic amino-acids	[279]
Photoproduction of tryptophan	[280]
Analysis of endogenous flavodiiron proteins	[281,282]
<i>cpcB</i>: phycocyanin synthesis gene	
Photoproduction of isoprene	[283,284]
Photoproduction of geranylinalool	[285]

Table 1. Cont.

Neutral Site and Objective of the Gene Manipulation	References
<i>glpK</i>, the gene encoding the glycerol kinase enzyme	
Photoproduction of lipid	[286]
Combination <i>psbA2</i> and <i>cpcB</i>	
Cloning of various genes for the photoproduction of isoprene	[287]
Photoproduction of β -phellandrene	[288–293]
<i>slr0646</i> encoding the PBP5 dispensable penicillin binding protein	[294]
Photoproduction of the p-coumaric acid	[295]
<i>ndhB</i> (slr0223): encoding subunit 2 of the NAD(P)H-dehydrogenase	
Analysis of the circadian expression of the DnaK (heat-shock protein) encoding gene	[296]
Construction of bioluminescent reporter strains for metal detection	[297]
Analysis of the light regulation of the photosystem I genes	[298,299]
<i>slr0168</i>	[220]
Analysis of tolerance to stresses	[300–302]
Analysis of endogenous and heterologous Fe- or Cu/Zn superoxide dismutase	[302]
Promoter analysis	[303]
Analysis of fatty-acids synthesis	[304]
Photoproduction of ethylene	[305,306]
Photoproduction of lactate	[307–310]
Photoproduction of 2,3-butanediol	[311]
Photoproduction of sucrose	[312]
Development of a marker-less gene replacement tool	[313]
Photoproduction of fatty alcohol	[314]
Photoproduction of isoprene	[275]
Photoproduction of ethanol	[315–317]
Photoproduction of glycerol	[318]
Photoproduction of <i>n</i> -butanol	[319]
Analysis of the regulation of Rubisco	[320]
Promoter analysis	[321]
Photoproduction of 1,2-propanediol	[322]
Analysis of alka(e)ne turnover	[323]
Photoproduction of mannitol	[324]
Photoproduction of bisabolene	[325]
<i>slr0168</i> and <i>slr1193</i>	
Photoproduction of ethanol	[326]
<i>slr0168</i> and <i>slr1556</i>	
Photoproduction of alkanes	[327]
<i>ndhB</i> and <i>slr0168</i>	
Photoproduction of ethanol	[315]
<i>psbA1</i> and <i>slr0168</i>	
Analysis of promoters	[328]
<i>psbA2</i> and <i>slr0168</i>	
Photoproduction of fatty-acids	[329]
Analysis of the cyanobacterial iron superoxide dismutase SOD	[302]
Photoproduction of ethylene	[330]
<i>phaCE</i> genes operating in the synthesis of polyhydroxybutyrate (PHB) biodegradable bioplastics	[331]
Photoproduction of acetone	[332]

Table 1. Cont.

Neutral Site and Objective of the Gene Manipulation	References
Intergenic region between <i>slr2030</i> and <i>slr2031</i>	
Analysis of glutathione synthesis	[333]
Analysis of heme oxygenase encoding genes	[334]
Photoproduction of poly-hydroxybutyrate (PHB) biodegradable bioplastics	[335]
Analysis of Flv3 flavodiiron protein	[336]
Photoproduction of pinene	[337]
Analysis of promoters and ribosome binding sites	[338]
Development of the CRISPR technologies for gene deletion or silencing	[339]
Photoproduction of ethylene	[340]
<i>slr0846</i> and <i>slr2030-slr2031</i> intergenic region	
Photoproduction of glutamate, linalool, and valencene	[341]
Intergenic regions between <i>sll0821-slr0846</i> and <i>slr2030-slr2031</i>	
Photoproduction of limonene	[342]
<i>slr0168</i> and <i>slr1704-sll1575</i> intergenic region	
Photoproduction of 3-hydroxypropionic acid	[343]
<i>slr0168</i>, <i>psbA2</i>, and <i>slr2030-Slr2031</i> intergenic region	
Photoproduction of the manoyl oxide terpene	[344]
Photoproduction of ethanol and butanol	[345]
Intergenic regions between <i>slr1495-sll1397</i>, <i>slr1362-sll1274</i>, <i>slr1828-sll1736</i>, and <i>slr1992-phaA2</i>	
Photoproduction of 3-hydroxybutyrate the precursor of the synthesis of PHB	[346]

The chromosomal cloning sites, written in bold cases are highlighted in grey color. The *cpcB* gene operates in the synthesis of the phycocyanin pigment. The *glpK* gene encodes the glycerol kinase. *slr0646* encodes the PBP5 dispensable penicillin binding protein [294]. The gene *ndhB* (sll0223) encodes a subunit of a NAD(P)H-dehydrogenase enzyme. The gene *slr0168* has no known function [220]. The *phaCE* genes operate in the synthesis of polyhydroxybutyrate (PHB) biodegradable bioplastics [331]. While *psbA1* is a silent gene [235], its homologue *psbA2* (slr1311) encodes the D1 protein of the photosystem II [259].

In *Synechococcus* PCC 7942 (Table 2), the most frequently employed chromosomal neutral sites are NSI (GenBank accession n° U30252), NSII (GenBank accession U44761) [253]), and NSIII (GenBank accession ABB56771.1) [254,255].

Table 2. Literature on the utilization of neutral chromosomal cloning sites in *Synechococcus* PCC 7942.

Neutral Site and Objective of the Gene Manipulation	References
NSI	
Photoproduction of 1,2-propanediol	[347]
Photoproduction of glycerol	[348]
Photoproduction of lactate	[349]
Photoproduction of succinate	[350]
Photoproduction of ethylene	[351]
Photoproduction of ethanol	[352]
NSII	
Photoproduction of free fatty acids	[353–355]
Photoproduction of B12 vitamin	[356]
NSIII	
Photoproduction of isobutyraldehyde	[357]
Photoproduction of carboxysome proteins	[358]

Table 2. Cont.

Neutral Site and Objective of the Gene Manipulation	References
NSI and NSII	
Analysis of the circadian rhythm	[359]
Photoproduction of isobutyraldehyde	[360]
Photoproduction of 1-butanol	[361,362]
Analysis of carboxysomes	[363,364]
Photoproduction of isopropanol	[364]
Photoproduction of isobutanol	[365]
Photoproduction of 3-hydroxypropionic acid	[366]
Photoproduction of 1,3-propanediol	[367]
Photoproduction of amorphaadiene and squalene	[368]
Photoproduction of limonene	[369]
Photoproduction of acetone	[370]
Photoproduction of isoprene	[371]
Analysis of gene-expression control systems	[372]
Photoproduction of 2,3-butanediol	[373]
Photoproduction of farnesene	[374,375]
Photoproduction of lactate	[376]
NSI and NSIII	
Photoproduction of biomass and sucrose export	[377]
Photoproduction of a synthetic CO ₂ -fixing photorespiratory bypass	[378]
Photoproduction of 2,3 butanediol	[379]
Photoproduction of amorphaadiene or squalene	[380]
NSI, NSII, and NSIII	
Overproduction of transporters to facilitate sugar export	[254]
Promoter analysis	[381]
Photoproduction of polyketides	[382]
Analysis of the influence of pilus biogenesis on the natural transformation	[229]
<i>psbA1</i>	
Photoproduction of ethylene	[383]
<i>glgC</i>	
Photoproduction of isobutanol	[366]
Intergenic region <i>Synpcc7942_0893</i> and <i>Synpcc7942_0894</i>	
Photoproduction of 2,3-butanediol	[384]

The chromosomal cloning sites, written in bold cases are highlighted in grey color.

In *Synechococcus* PCC 7002, the few neutral cloning sites employed (Table 3) are mainly the chromosomal genes *glpK*, which had been thought for some time to harbor a frameshift mutation preventing the production of a functional glycerol kinase enzyme [385], and *acsA*, which encodes an acetyl-CoA ligase, the inactivation of which conferred resistance to (3-hydroxy)propionate [386].

Table 3. Literature on the utilization of chromosomal neutral cloning sites in *Synechococcus* PCC 7002.

Neutral Cloning Sites	Objective of the Gene Manipulation and References
<i>glpK</i> (SYNPCC7002_A2842) encoding the glycerol kinase [385,386]	Removal of carboxysomes for containment of genetically modified strains [387]
<i>acsA</i> gene (SYNPCC7002_A1838) encoding an acetyl-CoA ligase and <i>glpK</i>	Analysis of an organic acid-based counter selection system [386]
<i>acsA</i> and <i>glpK</i>	Analysis of promoters and ribosome binding sites [388]
Integration between SYNPCC7002_A0935 and SYNPCC7002_A0936	Photoproduction of bisabolene and limonene [388]
<i>glpK</i> and <i>desB</i> (SYNPCC7002_A0159 encoding a ω 3 acyl-lipid desaturase [389] and integration between SYNPCC7002_A0935 and SYNPCC7002_A0936	Development of genetic tools [390]
<i>glpK</i> and integration between A0935-A0936	Engineering a strain for melamine degradation [391]
Intergenic regions between SYNPCC7002_A0932 and SYNPCC7002_A0933, SYNPCC7002_A1202 and SYNPCC7002_A1203, SYNPCC7002_A1778 and SYNPCC7002_A1779	Development of genetic tools [257]
SYNPCC7002_A1838, SYNPCC7002_A2542, and SYNPCC7002_A2842	Photoproduction of L-lysine [392]

The inactivation of the glycerol kinase *glpK* gene (SYNPCC7002_A2842), which was mistakenly annotated as having a frameshift mutation preventing the production of a functional protein [385], has no influence on the physiology of *Synechococcus* PCC 7002 [386]. Similarly, the inactivation of *desB* (SYNPCC7002_A0159, ω 3 acyl-lipid desaturase) has no detrimental influence at temperature above 22 °C [389]. The inactivation of the *acsA* gene (SYNPCC7002_A1838) encoding an acetyl-CoA ligase confers resistance to 3-hydroxypropionate and propionate [386].

7.4. Development of Transformable Shuttle Vectors Based on the Endogenous Plasmids of *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002

Because transformation and autonomously replicating plasmids have played a crucial role for gene manipulation in *Escherichia coli*, several groups tried to introduce an *E. coli* plasmid (pBR322 and its pUC derivatives) into cyanobacteria by transformation. All attempts were unsuccessful [163,178,231], in spite of a single report [393] that was never confirmed thereafter. These findings indicated that these *E. coli* plasmids are not able to replicate in cyanobacteria. Consequently, chimeric plasmids capable to replicate both in *E. coli* and a transformable cyanobacteria were constructed by cloning a small (cryptic) cyanobacterial plasmid (or a part of it) into an *E. coli* plasmid [394]. This approach was initiated in *Synechococcus* PCC 7942 using its smaller 7.84 kbp endogenous plasmid [182,395]. Then, the transformation efficiency was increased by plating the transformation mixture (recipient cells plus transforming DNA) on solid medium and incubating the plate in standard condition for 24 h prior to adding the selective antibiotic underneath the agar of the plate [183]. The transformation was more efficient when the recipient cyanobacterial cells were taken at the transition from the first to the second exponential growth phases [183], similarly to what has been observed for the transformation with linear chromosomal DNA [226]. The influence of the growth phase on transformation, which was not confirmed by other workers [396], was also observed in *Synechocystis* PCC 6803 [163].

Synechococcus PCC 7942 was found to be transformed more efficiently by chimeric-plasmid DNA isolated directly from this cyanobacterium rather than from *E. coli* [182]. This was explained by assuming that *Synechococcus* PCC 7942 and *E. coli* DNA are differently modified, for instance, by *dam*- or *dcm*-like DNA methylation systems [182]. Similarly, *Synechococcus* PCC 7002 was more efficiently transformed by its chimeric plasmid when it had been isolated directly from this host rather than from *E. coli* [397]. Furthermore, the elimination from this *Synechococcus* PCC 7002 biphasic plasmid of the *AvaI* restriction site (cleaved by the *AvaI*-isoschizomer *AquI* endonuclease of *Synechococcus* PCC 7002) strongly increased the efficiency of transformation to *Synechococcus* PCC 7002 [397]. The importance

of DNA modification for genetic transformation and cell fitness of cyanobacteria were firmly established in *Synechocystis* PCC 6803 [398–401].

The biphasic plasmids autonomously replicating in *E. coli* and a specific cyanobacterium were improved by several groups through the addition of restriction sites for facile gene cloning and/or several antibiotic-resistance genes for effective selection in *Synechococcus* PCC 7942 [402–404], *Synechocystis* PCC 6803 [163], and *Synechococcus* PCC 7002 [405]. These shuttle vectors were used for complementation analyses selecting for both the antibiotic resistance of the vector and the wild-type phenotype [406]. This approach allowed the identification and analysis of genes encoding the key stress-defense proteins RecA [405] and the Mn SOD [407]. One *Synechocystis* PCC 6803 shuttle vector served for analyzing the activity of several *E. coli* promoters, such as the *tac* promoter which appeared to be as effective in *Synechocystis* PCC 6803 as in *E. coli* [408]. This promoter-probe vector was also used to show that the lambda phage *Ci₈₅₇* gene (encoding a temperature sensitive repressor) and associated P_R promoter can be employed for strong and tight temperature-controlled gene expression [408]. Latter these gene expression devices were also shown to work well in other cyanobacteria [409,410].

The *Synechococcus* PCC 7002 and *Synechococcus* PCC 7942 shuttle vectors were employed for the production of heterologous larvicidal proteins in cyanobacterial cells, which turned out to be toxic when ingested by mosquito larvae [411,412]. Other studies reported the cloning of heterologous genes in *Synechococcus* PCC 7942 to increase its resistance to cadmium [413] or salt [414,415]. Similarly, the *Zymomonas mobilis* genes encoding the pyruvate decarboxylase (*pdh*) and alcohol dehydrogenase II (*adh*) were cloned into a *Synechococcus* PCC 7942 shuttle vector to engineer an ethanol producer [416]. The same strategy of cloning heterologous genes into a shuttle vector was used to generate a *Synechococcus* PCC 7942 recombinant strain for the production of ethylene [417–419]. Chimeric shuttle vectors were also used to overexpress endogenous metabolic genes to improve CO₂-fixation and/or biomass production in *Synechococcus* PCC 7942 [420].

7.5. The Chimeric Shuttle Vectors Based on an Endogenous Cyanobacterial Plasmid Tend to Have a Narrow-Host-Range of Replication

The abovementioned biphasic shuttle plasmids based on a cyanobacterial replicon appeared to replicate in one or a few genetically manipulable cyanobacteria. For examples, the *Synechococcus* PCC 7942 chimeric plasmids could not transform *Synechocystis* PCC 6803 [163], and reciprocally, the *Synechocystis* PCC 6803 shuttle vectors could not transform *Synechococcus* PCC 7942 [163]. In contrast, a shuttle vector based on an endogenous plasmid from *Synechococcus* PCC 7942 was shown to replicate in *Anabaena* PCC 7120, but not in *Synechocystis* PCC 6803 [421].

Plasmid vectors that can be transferred to, but cannot replicate in, a host are interesting for transporting DNA that must either transpose (e.g., a transposon, useful for random mutagenesis) or by homologous recombination in order to be stably maintained.

7.6. The Development of Autonomously Replicating Vectors Derived from the Broad-Host-Range Conjugative Plasmid RSF1010 Has Boosted the Genetics of Cyanobacteria

The interest of chimeric plasmid vectors that uses two narrow-host-range replicons originating from *E. coli* and a cyanobacterium is limited by the fact that they can shuttle only from *E. coli* to mostly the corresponding cyanobacterium. Such narrow-host-range vectors are not suitable for rapid tests and comparisons of a gene function in diverse cyanobacterial hosts to select (and engineer) a cyanobacterial chassis for an effective photosynthetic production of an industrially interesting chemical. Thus, several groups turned their attention to the naturally-occurring RSF1010 plasmid [422], a member of the incQ incompatibility group that replicates in a wide range of Gram-negative bacterial genera (cyanobacteria are also Gram-negative), including *Acetobacter*, *Acinetobacter*, *Agrobacterium*, *Alcaligenes*, *Azotobacter*, *Escherichia*, *Klebsiella*, *Methylophilus*, *Providencia*, *Pseudomonadales*, *Rhizobiaceae*, *Rhodospirillum rubrum*, and *Salmonella* SP- and *Serratia*. The value of RSF1010 as a shuttle vector is further enhanced by its ability to be transferred by conjugation from

an *E. coli* strain also carrying a self-transmissible (incP group) plasmid, such as RP4. The RSF1010-derived plasmids can be used to carry novel genetic information to those bacteria that are not capable of transformation [423–425].

RSF1010 and its derivatives were shown to be efficiently transferred by conjugation from *E. coli* to cyanobacteria where they replicate autonomously, even though they contain no cyanobacterial replicon. These cyanobacteria were, namely, *Synechocystis* PCC 6803 [426], *Synechococcus* NKBG15041C, *Pseudanabaena* NKBG040605C [427], *Synechocystis* PCC 6714, *Synechococcus* strains PCC7942 and PCC6301 [428], *Thermosynechococcus elongatus* PB1 [429], marine *Synechococcus* strains sp. WH7803, WH8102 and WH8103 [430], *Gloeobacter violaceus* PCC 7421 [431,432], *Prochlorococcus* MIT9313 [433], *Nostoc* (*Anabaena*) PCC 7120, *Nostoc punctiforme* ATCC29133 (also registered as PCC 73102) [255,434,435], *Leptolyngbya* BL0902 [436] and *Cyanothece* PCC 7425 [410].

Shortly after, the *Synechocystis* PCC6803 pioneering report [426], RSF1010 was used for the development of the first conjugative plasmid vector for promoter analysis [409] and regulated protein production in cyanobacteria [437]. The promoter-probe vector used the promoter-less chloramphenicol-acetyl-transferase (*cat*) gene as the reporter. When expressed by a studied promoter, *cat* directs the production of the CAT enzyme, the activity of which can be monitored by a spectrophotometric assay and confers the resistance to chloramphenicol [409]. This vector was used for the analysis of constitutive or regulated promoters [126,438,439] and references therein. The conditional expression vector [437] harbors the lambda-phage gene *cI₈₅₇* encoding a temperature-sensitive repressor that tightly controls the activity of the otherwise strong *p_R* promoter. Together these elements allow a tight temperature-controlled expression of the studied genes (no production of the corresponding proteins at temperatures below 30 °C, moderate level at 34–36 °C, and high production at 39–40 °C). This vector and its derivative harboring the *gfp* gene encoding the green-fluorescent reporter protein have been employed for proteins involved in photosynthesis [125], response to stress [34], cell division [440–442], and biogenesis of the carboxysome [410].

Other RSF1010-derived plasmids vectors have been used to analyze ribosome binding sites and transcription terminators [264], light-emitting proteins GFP, YFP (yellow-fluorescent protein), and luciferase [255,264,432,434]. RSF1010-derived vectors were also employed to analyze the role of carbon stores glycogen and PHB (polyhydroxybutyrate) in the tolerance to stress [443] and systems for the control of gene expression (in *Synechococcus* PCC 7942) [372], as well as various proteins such as the *Synechocystis* PCC 6803 photolyase enzyme PhrA [444] and to improve carbon fixation [445].

RSF1010 derivative plasmids were also employed in many works aiming at producing biotechnologically interesting products including bisabolol and patchoulol [446], ethanol [447], ethylene [306,340,448–450], erythritol [451], hydrogen [452,453], isobutanol [454,455], lactate [308], limonene [410], n-butanol [319], and triterpenes (lupeol, marnerol, and hydroxy-marnerol) [456].

Interestingly, in *Synechocystis* PCC 6803, it has been shown that RSF1010 and the pCC5.2 endogenous plasmid could be used for cloning, respectively, two pentose phosphate pathway native genes and the limonene synthase genes (*lims*) from either *Mentha spicata* or *Citrus limon*, which directed the production of limonene [262].

8. Interest and Limitation of the CRISPR/Cas Genome Editing Technology

Recently, the CRISPR/Cas system (CRISPR stands for clustered regularly interspaced short palindromic repeats and Cas for CRISPR-associated endonuclease) has facilitated the way genomes are edited in cyanobacteria, such as *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, *Synechococcus* PCC 7002, *Synechococcus* UTEX 2973, and the filamentous strain *Nostoc* (*Anabaena*) PCC 7120 (for reviews see [80,83,84,339,457]). Briefly, CRISPR/Cas genome editing systems exploit the Class II family of Cas endonucleases, which have a site-specific RNA-guided DNA cleavage activity. As compared to the well-established gene deletions techniques based on homologous DNA recombination, the interest of

the CRISPR/Cas system are (i) CRISPR/Cas systems can allow the engineering of non-transformable cyanobacteria, providing they can be manipulated by conjugation; (ii) a marker-less mutation is generated at the DNA target site; and (iii) multiple DNA loci can be modified simultaneously, by co-expressing the appropriate guide RNAs and editing templates.

The limitation of the CRISPR/Cas technology are the potential toxicity of the Cas DNase and the time required to eliminate the CRISPR/Cas plasmid vector from the generated mutant. However, this curing step can be accelerated by the presence of a negative selection marker in the CRISPR/Cas vector [80].

A variant of the CRISPR/Cas system, the CRISPRi (CRISPR interference) system that make use of DNase-inactive variants of Cas, is especially relevant to repress (fully or not) the transcription of studied genes, including the essential ones that cannot be deleted. This strategy was used for the targeted repression of vital genes to arrest growth and increase carbon partitioning and biofuel titers in *Synechocystis* PCC 6803 [345]. The CRISPRi technology was also employed to generate mutants with increased yields of growth and lactate secretion [458].

9. Responses to Stresses: The Recent Progress in Omics Technics Are Limited by the Large Number of Genes of Still Unknown Function

Because of their oxygenic photosynthesis, which triggered oxygen-promoted changes in metal availability, and the fact that they colonized most aquatic biotopes of our planet, cyanobacteria have always been challenged by changes in light, metals, and nutrients availabilities [28]. The responses to these stresses have been well studied with omics techniques that measure the changes in abundance of transcripts (transcriptomics), proteins (proteomics), or metabolites (metabolomics). The available genome sequences facilitate the use of transcriptomic and proteomic approaches.

Initially, transcriptomics focused on *Synechocystis* PCC 6803, because its genome has been the first to be fully sequenced [48] and used to develop the first commercially available DNA microarrays (IntelliGene™ CyanoCHIP; Takara Bio Inc., Shiga, Japan), which only comprised probes for protein-coding genes deposited on a glass support. These CyanoCHIP were used to the transcriptional responses to high light [459], inhibitors of photosynthesis [460], depletion of the LexA regulator [90], salt [461–464], cold stress [465], acid stress [466,467], heat shock [468–470], osmotic stress [465,471,472], oxidative stresses triggered by methyl viologen [473], or H₂O₂ [474,475], Cd-, Fe-, or Zn-stresses [474].

Later, several genome-wide *Synechocystis* PCC 6803 microarrays were also developed, based on long oligonucleotide probes (60- to 70-mer) spotted on a glass support. Such oligonucleotides-based microarrays circumvented the labor-intensive and error-prone steps of probe amplification and purification. They were used to study cell responses to sulfur starvation [476] and the deletion of the AbrB2 transcription regulator [477].

As an alternative to microarray analyses, which are based on hybridization of mRNA to DNA probes, the direct sequencing of RNA (an approach designated as RNA-Seq) was adapted to cyanobacteria [478]. RNA-Seq rapidly became the standard method for cyanobacterial transcriptomics [250,479]. It revealed that the *Synechocystis* PCC 6803 transcriptome includes more than 4000 transcriptional units, half of which representing small RNAs (sRNAs), which often harbor a small protein-coding sequence of less than 100 amino acid residues, and non-coding RNAs (ncRNAs) [480]. These ncRNAs could not be detected by DNA microarrays that only comprised probes for protein-coding genes. The vast majority of ncRNAs are still uncharacterized, and most of them are antisense transcripts (asRNAs). The phylogenetic conservation of ncRNAs across genomes of relatively distant cyanobacteria and their regulated transcription in response to major stresses, such as, light, iron, carbon, or nitrogen availability, nitrogen starvation [481], and butanol or ethanol stress [482], suggest that many ncRNAs may be involved in regulation [250,479].

Proteomics was also used to study the cyanobacterial responses to stresses, again starting with *Synechocystis* PCC 6803 [483], the genome of which is predicted to contain 3672 putative open reading frames (ORFs, i.e., protein coding sequences). Of these,

3264 and 408 ORFs are located on the chromosome and the seven endogenous plasmids, respectively [484]. Traditionally, two-dimensional polyacrylamide gels (2D-PAGE) and utilization of different fluorescence dyes (difference gel electrophoresis; 2D-DIGE) were employed to estimate concentrations for each protein between stress versus unstressed conditions. Later strategies took advantage of the sensitivity of liquid chromatography (LC), coupled with tandem mass spectrometry (MS), known as LC-MS/MS, for quantitative proteomic analysis, using different tags such as the isobaric tags for relative and absolute quantitation (iTRAQ)-based quantitative proteomics [485,486]. This quantitative technique became the dominant proteomics method for the identification of differentially expressed proteins of *Synechocystis* PCC 6803 [485–488].

These proteome techniques were employed to study the responses to cold [489]; copper [490]; CO₂ limitation [491]; high light [492]; high or low temperature [469,492–494]; high or low pH [495–497]; nitrogen-, phosphate, or sulfate-starvations [481,498,499]; metal stress [498,500,501]; salt stress [502–507]; and UV-B stress [508]. They also served to study the tolerance of *Synechocystis* PCC 6803 to butanol, ethanol, or hexane biofuels [488,509–512], as well as to analyze cyanobacterial strains engineered for the production of butanol [511], ethanol [510], hexane [509], or 3-hydroxypropionic acid [513].

In addition to the quantification of proteins, proteome methods can be used to identify protein modifications, such as glutathionylation [40], lysine malonylation [514], lysine methylation [515], phosphorylation [516–518], which are potentially involved in controlling protein activities.

However, our current understanding of the transcriptome and proteome responses to various challenges is limited by the fact that a large number of the responsive genes or proteins have still an unknown function.

In addition to transcriptomics and proteomics, metabolomics that focuses on low-molecular-weight metabolites provides the most straightforward characterization of metabolic responses to environmental changes. Compared to other omics studies, a few metabolomic research studies have been performed in cyanobacteria, and again *Synechocystis* PCC 6803 has been the most studied model. As many metabolites turn over quickly, fast sampling through fast culture filtration appeared to be very important in metabolomic analyses [519,520]. The combination of gas chromatography or liquid chromatography with mass spectrometry permits quantitative analysis of more than 100 metabolites in cyanobacterial cells. In addition to metabolomics, which seeks comprehensive profiling of predominantly intra-organism compounds, volatilomics assesses those compounds released by an organism: the key components of chemically mediated inter-organismal communication [521,522]. The field of volatilomics grew out as advances in collection methods of volatile organic compounds and gas chromatography coupled with mass spectrometry.

Finally, in several studies, transcriptomics, proteomics, and/or metabolomics were integrated to better analyze the responses of *Synechocystis* PCC 6803 to environmental conditions [492,497,523–529], as well as the production of [530] or the tolerance to chemicals [531]. In some cases, it appeared that omics data at different levels do not necessarily correlate a finding that can be explained by regulations occurring at the levels of gene expression [6,93,479,532–534] and/or enzyme stability and activity [40,483,527].

10. Conclusions

Cyanobacteria are a widely-diverse photosynthetic prokaryotes of wide interest for basic and applied sciences. So far, cyanobacterial research has focused primarily on a few models, such as the three unicellular non-nitrogen fixing species *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002, which are straightforward to culture under laboratory conditions, easily amenable to genetic modification and can be frozen for long-term storage. Extensive “omics” data sets and many genetic tools and genome-scale metabolic models (GSM) have been generated to guide the engineering of these model cyanobacteria for the photosynthetic production of biotechnologically interesting chemicals. Interestingly, it has been put forward that GSM should take into account

and describe photon absorption and light-shading thereby addressing the challenge of accurately modeling light as a metabolite [535,536]. In addition, GSM should be validated with experimental data obtained after measurement of metabolic fluxes and metabolic pool sizes [80,86,87,536]. However, omics data interpretation and GSM metabolic designs are based on our currently limited understanding of the genotype–phenotype relationships of cyanobacteria. Thus, to generate robust and predictive GSM models of the cyanobacterial metabolism, it is important to continue the analysis of *Synechocystis* PCC 6803, *Synechococcus* PCC 7942 and *Synechococcus* PCC 7002, and increase the efforts to

- (i) verify the function of numerous genes that have been annotated merely by sequence analogy with those genes characterized only in intensively studied non-photosynthetic models (*E. coli*, yeast, etc.), which may have a different function in cyanobacteria;
- (ii) and analyze the specificity/redundancy of multiple gene families;
- (iii) characterize the function of the large number of as yet unknown genes and non-coding RNAs;
- (iv) identify the comprehensive set of genes that are essential to the growth of cells incubated in well-defined conditions.

Furthermore, most of the attempts to reprogram *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, or *Synechococcus* PCC 7002 for the photoproduction of chemicals have focused on increasing product synthesis by small-scale cultures growing under laboratory conditions because most academic researchers lack access to large-scale production systems that are necessary to evaluate the potential of engineered strains under realistic industrial conditions.

Moreover *Synechocystis* PCC 6803, *Synechococcus* PCC 7942, and *Synechococcus* PCC 7002 represent only a limited part of the wide biodiversity of cyanobacteria. This arguably limits fundamental discovery and applied research towards wider commercialization. Thus, new phylogenetically-distant candidate cyanobacteria should be isolated and developed from diverse environments with a robust growth and high tolerance to local conditions, so as to be used as chassis for the photosynthetic production of high-value chemicals in diverse industrial sites. We think that the genetic modifiability of such candidate strains using the conjugative transfer of RSF1010-derived broad-host-range plasmids will be key for such works.

To summarize, we recommend to strengthen the communication between academic researchers, who know well cyanobacteria and can manipulate them, but have a limited access to large photobioreactors and industrial partners, who attempt to use cyanobacteria to produce interesting chemicals at reasonable costs, but often lack knowledge on cyanobacterial genetics, physiology, and metabolism. Moreover, to minimize operation costs we need to develop robust cyanobacteria capable to grow on industrial waters and fumes, in huge photobioreactors, as well as well as efficient technologies to harvest the end products.

Author Contributions: Conceptualization, C.C.-C. and F.C.; writing—original draft preparation, F.C., C.C.-C., and V.B.-G.; writing—review and editing, F.C., C.C.-C., and V.B.-G.; supervision, F.C. and C.C.-C.; project administration, F.C. and C.C.-C.; funding acquisition, F.C. and C.C.-C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded in part by the ANR CalvinDesign (ANR-17-CE05-0001).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Soo, R.M.; Hemp, J.; Hugenholtz, P. Evolution of photosynthesis and aerobic respiration in the cyanobacteria. *Free Radic. Biol. Med.* **2019**, *140*, 200–205. [\[CrossRef\]](#)
2. Garcia-Pichel, F.; Zehr, J.P.; Bhattacharya, D.; Pakrasi, H.B. What's in a name? The case of cyanobacteria. *J. Phycol.* **2020**, *56*, 1–5. [\[CrossRef\]](#)
3. Hamilton, T.L.; Bryant, D.A.; Macalady, J.L. The role of biology in planetary evolution: Cyanobacterial primary production in low-oxygen Proterozoic oceans. *Environ. Microbiol.* **2016**, *18*, 325–340. [\[CrossRef\]](#)
4. Ponce-Toledo, R.I.; Deschamps, P.; López-García, P.; Zivanovic, Y.; Benzerara, K.; Moreira, D. An Early-Branching Freshwater Cyanobacterium at the Origin of Plastids. *Curr. Biol.* **2017**, *27*, 386–391. [\[CrossRef\]](#)
5. Ducat, D.C.; Way, J.C.; Silver, P.A. Engineering cyanobacteria to generate high-value products. *Trends Biotechnol.* **2011**, *29*, 95–103. [\[CrossRef\]](#)
6. Veaudor, T.; Blanc-Garin, V.; Chenebault, C.; Diaz-Santos, E.; Sassi, J.F.; Cassier-Chauvat, C.; Chauvat, F. Recent advances in the photoautotrophic metabolism of cyanobacteria: Biotechnological implications. *Life* **2020**, *10*, 71. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Mills, L.A.; McCormick, A.J.; Lea-Smith, D.J. Current knowledge and recent advances in understanding metabolism of the model cyanobacterium *Synechocystis* sp. PCC 6803. *Biosci. Rep.* **2020**, *40*. [\[CrossRef\]](#)
8. Garcia-Pichel, F.; Belnap, J.; Neuer, S.; Schanz, F. Estimates of global cyanobacterial biomass and its distribution. *Arch. Hydrobiol. Suppl. Algol. Stud.* **2009**, *109*, 213–227. [\[CrossRef\]](#)
9. Chang, A.C.G.; Chen, T.; Li, N.; Duan, J. Perspectives on endosymbiosis in coralloid roots: Association of cycads and cyanobacteria. *Front. Microbiol.* **2019**, *10*. [\[CrossRef\]](#) [\[PubMed\]](#)
10. Cassier-Chauvat, C.; Chauvat, F. Cyanobacteria: Wonderful Microorganisms for Basic and Applied Research. *eLS* **2018**, 1–11. [\[CrossRef\]](#)
11. Cassier-Chauvat, C.; Chauvat, F. Cell division in cyanobacteria. In *The Cell Biology of Cyanobacteria*; Flores, E., Herrero, A., Eds.; Caister Academic Press: Poole, UK, 2014.
12. Springstein, B.L.; Nürnberg, D.J.; Weiss, G.L.; Pilhofer, M.; Stucken, K. Structural determinants and their role in cyanobacterial morphogenesis. *Life* **2020**, *10*, 355. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Chauvat, F.; Corre, B.; Herdman, M.; Joset-Espardellier, F. Energetic and metabolic requirements for the germination of akinetes of the cyanobacterium *Nostoc* PCC 7524. *Arch. Microbiol.* **1982**, *133*, 44–49. [\[CrossRef\]](#)
14. Herrero, A.; Stavans, J.; Flores, E. The multicellular nature of filamentous heterocyst-forming cyanobacteria. *FEMS Microbiol. Rev.* **2016**, *40*, 831–854. [\[CrossRef\]](#)
15. Dittmann, E.; Gugger, M.; Sivonen, K.; Fewer, D.P. Natural Product Biosynthetic Diversity and Comparative Genomics of the Cyanobacteria. *Trends Microbiol.* **2015**, *23*, 642–652. [\[CrossRef\]](#)
16. Cassier-Chauvat, C.; Dive, V.; Chauvat, F. Cyanobacteria: Photosynthetic factories combining biodiversity, radiation resistance, and genetics to facilitate drug discovery. *Appl. Microbiol. Biotechnol.* **2017**, *101*, 1359–1364. [\[CrossRef\]](#)
17. Demay, J.; Bernard, C.; Reinhardt, A.; Marie, B. Natural products from cyanobacteria: Focus on beneficial activities. *Mar. Drugs* **2019**, *17*, 320. [\[CrossRef\]](#)
18. Pereira, S.B.; Sousa, A.; Santos, M.; Araújo, M.; Serôdio, F.; Granja, P.; Tamagnini, P. Strategies to obtain designer polymers based on cyanobacterial extracellular polymeric substances (EPS). *Int. J. Mol. Sci.* **2019**, *20*, 5693. [\[CrossRef\]](#)
19. Knot, C.J.; Ungerer, J.; Wangikar, P.P.; Pakrasi, H.B. Cyanobacteria: Promising biocatalysts for sustainable chemical production. *J. Biol. Chem.* **2018**, *293*, 5044–5052. [\[CrossRef\]](#)
20. Lin, P.C.; Pakrasi, H.B. Engineering cyanobacteria for production of terpenoids. *Planta* **2019**, *249*, 145–154. [\[CrossRef\]](#)
21. Schopf, J.W. The paleobiological record of photosynthesis. *Photosynth. Res.* **2011**, *107*, 87–101. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Imlay, J.A. The molecular mechanisms and physiological consequences of oxidative stress: Lessons from a model bacterium. *Nat. Rev. Microbiol.* **2013**, *11*, 443–454. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Shimakawa, G.; Matsuda, Y.; Nakajima, K.; Tamoi, M.; Shigeoka, S.; Miyake, C. Diverse strategies of O₂ usage for preventing photo-oxidative damage under CO₂ limitation during algal photosynthesis. *Sci. Rep.* **2017**, *7*. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Diaz, J.M.; Plummer, S. Production of extracellular reactive oxygen species by phytoplankton: Past and future directions. *J. Plankton Res.* **2018**, *40*, 655–666. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Narainsamy, K.; Marteyn, B.; Sakr, S.; Cassier-Chauvat, C.; Chauvat, F. Genomics of the Pleiotropic Glutathione System in Cyanobacteria. *Adv. Bot. Res.* **2013**, *65*, 157–188.
26. Picciocchi, A.; Saguez, C.; Boussac, A.; Cassier-Chauvat, C.; Chauvat, F. CGFS-type monothiol glutaredoxins from the cyanobacterium *Synechocystis* PCC6803 and other evolutionary distant model organisms possess a glutathione-ligated [2Fe-2S] cluster. *Biochemistry* **2007**, *46*, 15018–15026. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Fahey, R.C. Glutathione analogs in prokaryotes. *Biochim. Biophys. Acta Gen. Subj.* **2013**, *1830*, 3182–3198. [\[CrossRef\]](#)
28. Cassier-Chauvat, C.; Chauvat, F. Genomics of the Resistance to Metal and Oxidative Stresses in Cyanobacteria. *Stress Environ. Regul. Gene Expr. Adapt. Bact.* **2016**, *2*, 1154–1164.
29. Kammerscheit, X.; Chauvat, F.; Cassier-Chauvat, C. From cyanobacteria to human, MAPEG-type glutathione-S-transferases operate in cell tolerance to heat, cold, and lipid peroxidation. *Front. Microbiol.* **2019**, *10*, 2248. [\[CrossRef\]](#)
30. Berndt, C.; Lillig, C.H. Glutathione, Glutaredoxins, and Iron. *Antioxid. Redox Signal.* **2017**, *27*, 1235–1251. [\[CrossRef\]](#)

31. Narainsamy, K.; Farci, S.; Braun, E.; Junot, C.; Cassier-Chauvat, C.; Chauvat, F. Oxidative-stress detoxification and signalling in cyanobacteria: The crucial glutathione synthesis pathway supports the production of ergothioneine and ophthalmate. *Mol. Microbiol.* **2016**, *100*, 15–24. [[CrossRef](#)]
32. Kammerscheit, X.; Chauvat, F.; Cassier-Chauvat, C. First in vivo evidence that glutathione-s-transferase operates in photo-oxidative stress in cyanobacteria. *Front. Microbiol.* **2019**, *10*, 1899. [[CrossRef](#)]
33. Noctor, G.; Mhamdi, A.; Chaouch, S.; Han, Y.; Neukermans, J.; Marquez-Garcia, B.; Queval, G.; Foyer, C.H. Glutathione in plants: An integrated overview. *Plant Cell Environ.* **2012**, *35*, 454–484. [[CrossRef](#)] [[PubMed](#)]
34. Marteyn, B.; Sakr, S.; Farci, S.; Bedhomme, M.; Chardonnet, S.; Decottignies, P.; Lemaire, S.D.; Cassier-Chauvat, C.; Chauvat, F. The *Synechocystis* PCC6803 MerA-like enzyme operates in the reduction of both mercury and uranium under the control of the glutaredoxin 1 enzyme. *J. Bacteriol.* **2013**, *195*, 4138–4145. [[CrossRef](#)] [[PubMed](#)]
35. Cassier-Chauvat, C.; Chauvat, F. Responses to oxidative and heavy metal stresses in cyanobacteria: Recent advances. *Int. J. Mol. Sci.* **2015**, *16*, 871. [[CrossRef](#)] [[PubMed](#)]
36. Thorsen, M.; Jacobson, T.; Vooijs, R.; Navarrete, C.; Bliiek, T.; Schat, H.; Tamás, M.J. Glutathione serves an extracellular defence function to decrease arsenite accumulation and toxicity in yeast. *Mol. Microbiol.* **2012**, *84*, 1177–1188. [[CrossRef](#)] [[PubMed](#)]
37. Przybyla-Toscano, J.; Roland, M.; Gaymard, F.; Couturier, J.; Rouhier, N. Roles and maturation of iron–sulfur proteins in plastids. *J. Biol. Inorg. Chem.* **2018**, *23*, 545–566. [[CrossRef](#)]
38. Daniel, T.; Faruq, H.M.; Laura Magdalena, J.; Manuela, G.; Christopher Horst, L. Role of GSH and Iron-Sulfur Glutaredoxins in Iron Metabolism-Review. *Molecules* **2020**, *27*, 3860. [[CrossRef](#)]
39. Mailloux, R.J. Protein S–glutathionylation reactions as a global inhibitor of cell metabolism for the desensitization of hydrogen peroxide signals. *Redox Biol.* **2020**, *32*. [[CrossRef](#)]
40. Chardonnet, S.; Sakr, S.; Cassier-Chauvat, C.; Le Maréchal, P.; Chauvat, F.; Lemaire, S.D.; Decottignies, P. First proteomic study of S–glutathionylation in cyanobacteria. *J. Proteome Res.* **2015**, *14*, 59–71. [[CrossRef](#)]
41. Sakr, S.; Dutheil, J.; Saenkham, P.; Bottin, H.; Leplat, C.; Ortega-Ramos, M.; Aude, J.C.; Chapuis, V.; Guedeney, G.; Decottignies, P.; et al. The activity of the *Synechocystis* PCC6803 AbrB2 regulator of hydrogen production can be post-translationally controlled through glutathionylation. *Int. J. Hydrog. Energy* **2013**, *38*, 13547–13555. [[CrossRef](#)]
42. Gallé, Á.; Czékus, Z.; Bela, K.; Horváth, E.; Ördög, A.; Csiszár, J.; Poór, P. Plant glutathione transferases and light. *Front. Plant Sci.* **2019**, *9*. [[CrossRef](#)] [[PubMed](#)]
43. Sylvestre-Gonon, E.; Law, S.R.; Schwartz, M.; Robe, K.; Keech, O.; Didierjean, C.; Dubos, C.; Rouhier, N.; Hecker, A. Functional, structural and biochemical features of plant serinyl-glutathione transferases. *Front. Plant Sci.* **2019**, *10*. [[CrossRef](#)] [[PubMed](#)]
44. Couto, N.; Wood, J.; Barber, J. The role of glutathione reductase and related enzymes on cellular redox homeostasis network. *Free Radic. Biol. Med.* **2016**, *95*, 27–42. [[CrossRef](#)] [[PubMed](#)]
45. Marteyn, B.; Domain, F.; Legrain, P.; Chauvat, F.; Cassier-Chauvat, C. The thioredoxin reductase-glutaredoxins-ferredoxin crossroad pathway for selenate tolerance in *Synechocystis* PCC6803. *Mol. Microbiol.* **2009**, *71*, 520–532. [[CrossRef](#)] [[PubMed](#)]
46. Fleischmann, R.D.; Adams, M.D.; White, O.; Clayton, R.A.; Kirkness, E.F.; Kerlavage, A.R.; Bult, C.J.; Tomb, J.F.; Dougherty, B.A.; Merrick, J.M.; et al. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **1995**, *269*, 496–512. [[CrossRef](#)]
47. Fraser, C.M.; Gocayne, J.D.; White, O.; Adams, M.D.; Clayton, R.A.; Fleischmann, R.D.; Bult, C.J.; Kerlavage, A.R.; Sutton, G.; Kelley, J.M.; et al. The minimal gene complement of *Mycoplasma genitalium*. *Science* **1995**, *270*, 397–403. [[CrossRef](#)]
48. Kaneko, T.; Sato, S.; Kotani, H.; Tanaka, A.; Asamizu, E.; Nakamura, Y.; Miyajima, N.; Hirosawa, M.; Sugiura, M.; Sasamoto, S.; et al. Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* **1996**, *3*, 109–136. [[CrossRef](#)]
49. Cao, H.; Shimura, Y.; Steffen, M.M.; Yang, Z.; Lu, J.; Joel, A.; Jenkins, L.; Kawachi, M.; Yin, Y.; Garcia-Pichel, F. The trait repertoire enabling cyanobacteria to bloom assessed through comparative genomic complexity and metatranscriptomics. *MBio* **2020**, *11*. [[CrossRef](#)]
50. Fernandez, L.; Peura, S.; Eiler, A.; Linz, A.M.; McMahon, K.D.; Bertilsson, S. Diazotroph Genomes and Their Seasonal Dynamics in a Stratified Humic Bog Lake. *Front. Microbiol.* **2020**, *11*. [[CrossRef](#)]
51. Grettenberger, C.L.; Sumner, D.Y.; Wall, K.; Brown, C.T.; Eisen, J.A.; Mackey, T.J.; Hawes, I.; Jospin, G.; Jungblut, A.D. A phylogenetically novel cyanobacterium most closely related to *Gloeobacter*. *ISME J.* **2020**, *14*, 2142–2152. [[CrossRef](#)]
52. Salazar, V.W.; Tschoeke, D.A.; Swings, J.; Cosenza, C.A.; Mattoso, M.; Thompson, C.C.; Thompson, F.L. A new genomic taxonomy system for the *Synechococcus* collective. *Environ. Microbiol.* **2020**. [[CrossRef](#)] [[PubMed](#)]
53. Tripp, H.J.; Bench, S.R.; Turk, K.A.; Foster, R.A.; Desany, B.A.; Niazi, F.; Affourtit, J.P.; Zehr, J.P. Metabolic streamlining in an open-ocean nitrogen-fixing cyanobacterium. *Nature* **2010**, *464*, 90–94. [[CrossRef](#)] [[PubMed](#)]
54. Dagan, T.; Roettger, M.; Stucken, K.; Landan, G.; Koch, R.; Major, P.; Gould, S.B.; Goremykin, V.V.; Rippka, R.; De Marsac, N.T.; et al. Genomes of stigonematalean cyanobacteria (subsection V) and the evolution of oxygenic photosynthesis from prokaryotes to plastids. *Genome Biol. Evol.* **2013**, *5*, 31–44. [[CrossRef](#)] [[PubMed](#)]
55. Larsson, J.; Nylander, J.A.A.; Bergman, B. Genome fluctuations in cyanobacteria reflect evolutionary, developmental and adaptive traits. *BMC Evol. Biol.* **2011**, *11*. [[CrossRef](#)] [[PubMed](#)]
56. Cassier-Chauvat, C.; Poncet, M.; Chauvat, F. Three insertion sequences from the cyanobacterium *Synechocystis* PCC6803 support the occurrence of horizontal DNA transfer among bacteria. *Gene* **1997**, *195*, 257–266. [[CrossRef](#)]

57. Zhaxybayeva, O.; Gogarten, J.P.; Charlebois, R.L.; Doolittle, W.F.; Papke, R.T. Phylogenetic analyses of cyanobacterial genomes: Quantification of horizontal gene transfer events. *Genome Res.* **2006**, *16*, 1099–10108. [[CrossRef](#)]
58. Cassier-Chauvat, C.; Veaudor, T.; Chauvat, F. Comparative genomics of DNA recombination and repair in cyanobacteria: Biotechnological implications. *Front. Microbiol.* **2016**, *7*. [[CrossRef](#)] [[PubMed](#)]
59. Godde, J.S.; Baichoo, S.; Mungloo-Dilmohamud, Z.; Jaufeerally-Fakim, Y. Comparison of genomic islands in cyanobacteria: Evidence of bacteriophage-mediated horizontal gene transfer from eukaryotes. *Microbiol. Res.* **2018**, *211*, 31–46. [[CrossRef](#)]
60. Zeidner, G.; Bielawski, J.P.; Shmoish, M.; Scanlan, D.J.; Sabeji, G.; Bèjà, O. Potential photosynthesis gene recombination between *Prochlorococcus* and *Synechococcus* via viral intermediates. *Environ. Microbiol.* **2005**, *7*, 1505–1513. [[CrossRef](#)]
61. Scanlan, D.J.; Ostrowski, M.; Mazard, S.; Dufresne, A.; Garczarek, L.; Hess, W.R.; Post, A.F.; Hagemann, M.; Paulsen, I.; Partensky, F. Ecological Genomics of Marine Picocyanobacteria. *Microbiol. Mol. Biol. Rev.* **2009**, *73*, 249–299. [[CrossRef](#)]
62. Nicholson, M.L.; Gaasenbeek, M.; Laudénbach, D.E. Two enzymes together capable of cysteine biosynthesis are encoded on a cyanobacterial plasmid. *MGG Mol. Gen. Genet.* **1995**, *247*, 623–632. [[CrossRef](#)] [[PubMed](#)]
63. Kimura, A.; Hamada, T.; Morita, E.H.; Hayashi, H. A high temperature-sensitive mutant of *Synechococcus* sp. PCC 7002 with modifications in the endogenous plasmid, pAQ1. *Plant Cell Physiol.* **2002**, *43*, 217–223. [[CrossRef](#)] [[PubMed](#)]
64. Nagarajan, S.; Sherman, D.M.; Shaw, I.; Sherman, L.A. Functions of the duplicated hik31 operons in central metabolism and responses to light, dark, and carbon sources in *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **2012**, *194*, 448–459. [[CrossRef](#)] [[PubMed](#)]
65. Swingley, W.D.; Chen, M.; Cheung, P.C.; Conrad, A.L.; Dejesa, L.C.; Hao, J.; Honchak, B.M.; Karbach, L.E.; Kurdoglu, A.; Lahiri, S.; et al. Niche adaptation and genome expansion in the chlorophyll d-producing cyanobacterium *Acaryochloris marina*. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 2005–2010. [[CrossRef](#)] [[PubMed](#)]
66. Welsh, E.A.; Liberton, M.; Stöckel, J.; Loh, T.; Elvitigala, T.; Wang, C.; Wollam, A.; Fulton, R.S.; Clifton, S.W.; Jacobs, J.M.; et al. The genome of *Cyanothece* 51142, a unicellular diazotrophic cyanobacterium important in the marine nitrogen cycle. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 15094–15099. [[CrossRef](#)]
67. Bandyopadhyay, A.; Elvitigala, T.; Welsh, E.; Stöckel, J.; Liberton, M.; Min, H.; Sherman, L.A.; Pakrasi, H.B. Novel metabolic attributes of the genus *Cyanothece*, comprising a group of unicellular nitrogen-fixing cyanobacteria. *MBio* **2011**, *2*. [[CrossRef](#)]
68. Quinn, G.A.; Banat, A.M.; Abdelhameed, A.M.; Banat, I.M. Streptomyces from traditional medicine: Sources of new innovations in antibiotic discovery. *J. Med. Microbiol.* **2020**, *69*, 1040–1048. [[CrossRef](#)]
69. Ahmed, M.N.; Reyna-González, E.; Schmid, B.; Wiebach, V.; Süßmuth, R.D.; Dittmann, E.; Fewer, D.P. Phylogenomic Analysis of the Microviridin Biosynthetic Pathway Coupled with Targeted Chemo-Enzymatic Synthesis Yields Potent Protease Inhibitors. *ACS Chem. Biol.* **2017**, *12*, 1538–1546. [[CrossRef](#)]
70. Hoff, G.; Bertrand, C.; Piotrowski, E.; Thibessard, A.; Leblond, P. Genome plasticity is governed by double strand break DNA repair in *Streptomyces*. *Sci. Rep.* **2018**, *8*. [[CrossRef](#)]
71. Labella, J.I.; Llop, A.; Contreras, A. The default cyanobacterial linked genome: An interactive platform based on cyanobacterial linkage networks to assist functional genomics. *FEBS Lett.* **2020**, *594*, 1661–1674. [[CrossRef](#)]
72. Simm, S.; Keller, M.; Selymes, M.; Schleiff, E. The composition of the global and feature specific cyanobacterial core-genomes. *Front. Microbiol.* **2015**, *6*. [[CrossRef](#)]
73. Alvarenga, D.O.; Fiore, M.F.; Varani, A.M. A metagenomic approach to cyanobacterial genomics. *Front. Microbiol.* **2017**. [[CrossRef](#)]
74. Gonzalez-Esquer, C.R.; Smarda, J.; Rippka, R.; Axen, S.D.; Guglielmi, G.; Gugger, M.; Kerfeld, C.A. Cyanobacterial ultrastructure in light of genomic sequence data. *Photosynth. Res.* **2016**, *192*, 147–157. [[CrossRef](#)]
75. Shih, P.M.; Wu, D.; Latifi, A.; Axen, S.D.; Fewer, D.P.; Talla, E.; Calteau, A.; Cai, F.; Tandeau de Marsac, N.; Rippka, R.; et al. Improving the coverage of the cyanobacterial phylum using diversity-driven genome sequencing. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 1053–1058. [[CrossRef](#)] [[PubMed](#)]
76. Harel, A.; Karkar, S.; Cheng, S.; Falkowski, P.G.; Bhattacharya, D. Deciphering primordial cyanobacterial genome functions from protein network analysis. *Curr. Biol.* **2015**, *25*, 628–634. [[CrossRef](#)] [[PubMed](#)]
77. Sánchez-Baracaldo, P.; Cardona, T. On the origin of oxygenic photosynthesis and Cyanobacteria. *New Phytol.* **2020**, *225*, 1440–1446. [[CrossRef](#)]
78. Ponce-Toledo, R.I.; López-García, P.; Moreira, D. Horizontal and endosymbiotic gene transfer in early plastid evolution. *New Phytol.* **2019**, *224*, 618–624. [[CrossRef](#)] [[PubMed](#)]
79. Kopf, M.; Klähn, S.; Pade, N.; Weingärtner, C.; Hagemann, M.; Voß, B.; Hess, W.R. Comparative genome analysis of the closely related *Synechocystis* strains PCC 6714 and PCC 6803. *DNA Res.* **2014**, *21*, 255–266. [[CrossRef](#)]
80. Gale, G.A.R.; Osorio, A.A.S.; Mills, L.A.; Wang, B.; Lea-Smith, D.J.; McCormick, A.J. Emerging species and genome editing tools: Future prospects in cyanobacterial synthetic biology. *Microorganisms* **2019**, *7*, 409. [[CrossRef](#)]
81. Janasch, M.; Asplund-Samuelsson, J.; Steuer, R.; Hudson, E.P. Kinetic modeling of the Calvin cycle identifies flux control and stable metabolomes in *Synechocystis* carbon fixation. *J. Exp. Bot.* **2019**, *70*, 973–983. [[CrossRef](#)]
82. Lasry Testa, R.; Delpino, C.; Estrada, V.; Diaz, S.M. In silico strategies to couple production of bioethanol with growth in cyanobacteria. *Biotechnol. Bioeng.* **2019**, *116*, 2061–2073. [[CrossRef](#)]
83. Khan, A.Z.; Bilal, M.; Mehmood, S.; Sharma, A.; Iqbal, H.M.N. State-of-the-art genetic modalities to engineer cyanobacteria for sustainable biosynthesis of biofuel and fine-chemicals to meet bio-economy challenges. *Life* **2019**, *9*, 54. [[CrossRef](#)] [[PubMed](#)]

84. Santos-Merino, M.; Singh, A.K.; Ducat, D.C. New applications of synthetic biology tools for cyanobacterial metabolic engineering. *Front. Bioeng. Biotechnol.* **2019**, *7*. [[CrossRef](#)] [[PubMed](#)]
85. Wu, C.; Jiang, H.; Kalra, I.; Wang, X.; Cano, M.; Maness, P.C.; Yu, J.; Xiong, W. A generalized computational framework to streamline thermodynamics and kinetics analysis of metabolic pathways. *Metab. Eng.* **2020**, *57*, 140–150. [[CrossRef](#)]
86. Hendry, J.I.; Bandyopadhyay, A.; Srinivasan, S.; Pakrasi, H.B.; Maranas, C.D. Metabolic model guided strain design of cyanobacteria. *Curr. Opin. Biotechnol.* **2020**, *64*, 17–23. [[CrossRef](#)]
87. Mukherjee, B.; Madhu, S.; Wangikar, P.P. The role of systems biology in developing non-model cyanobacteria as hosts for chemical production. *Curr. Opin. Biotechnol.* **2020**, *64*, 62–69. [[CrossRef](#)] [[PubMed](#)]
88. Bender, R.A. The danger of annotation by analogy: Most “thiii” genes play no role in thiamine biosynthesis. *J. Bacteriol.* **2011**, *193*, 4574–4575. [[CrossRef](#)]
89. Fujisawa, T.; Narikawa, R.; Maeda, S.I.; Watanabe, S.; Kanesaki, Y.; Kobayashi, K.; Nomata, J.; Hanaoka, M.; Watanabe, M.; Ehira, S.; et al. CyanoBase: A large-scale update on its 20th anniversary. *Nucleic Acids Res.* **2017**, *45*, D551–D554. [[CrossRef](#)]
90. Domain, F.; Houot, L.; Chauvat, F.; Cassier-Chauvat, C. Function and regulation of the cyanobacterial genes *lexA*, *recA* and *ruvB*: LexA is critical to the survival of cells facing inorganic carbon starvation. *Mol. Microbiol.* **2004**, *53*, 65–80. [[CrossRef](#)]
91. Figge, R.M.; Cassier-Chauvat, C.; Chauvat, F.; Cerff, R. Characterization and analysis of an NAD(P)H dehydrogenase transcriptional regulator critical for the survival of cyanobacteria facing inorganic carbon starvation and osmotic stress. *Mol. Microbiol.* **2001**, *39*, 455–468. [[CrossRef](#)]
92. Jiang, Y.L.; Wang, X.P.; Sun, H.; Han, S.J.; Li, W.F.; Cui, N.; Lin, G.M.; Zhang, J.Y.; Cheng, W.; Cao, D.D.; et al. Coordinating carbon and nitrogen metabolic signaling through the cyanobacterial global repressor NdhR. *Proc. Natl. Acad. Sci. USA* **2017**, *115*, 403–408. [[CrossRef](#)]
93. Forchhammer, K.; Selim, K.A. Carbon/nitrogen homeostasis control in cyanobacteria. *FEMS Microbiol. Rev.* **2019**, *44*, 33–53. [[CrossRef](#)]
94. Cotton, C.A.R.; Kabasakal, B.V.; Miah, N.A.; Murray, J.W. Structure of the dual-function fructose-1,6/sedoheptulose-1,7-bisphosphatase from *Thermosynechococcus elongatus* bound with sedoheptulose-7-phosphate. *Acta Crystallogr. Sect. Struct. Biol. Commun.* **2015**, *71*, 1341–1345. [[CrossRef](#)]
95. Mohamed, H.E.; Vermaas, W.F.J. S110254 (CrtLdiox) is a bifunctional lycopene cyclase/dioxygenase in cyanobacteria producing myxoxanthophyll. *J. Bacteriol.* **2006**, *188*, 3337–3344. [[CrossRef](#)]
96. Yeremenko, N.; Kouřil, R.; Ihalainen, J.A.; D’Haene, S.; Van Oosterwijk, N.; Andrizhievskaya, E.G.; Keegstra, W.; Dekker, H.L.; Hagemann, M.; Boekema, E.J.; et al. Supramolecular organization and dual function of the IsiA chlorophyll-binding protein in cyanobacteria. *Biochemistry* **2004**, *43*, 10308–10313. [[CrossRef](#)] [[PubMed](#)]
97. Burnat, M.; Picossi, S.; Valladares, A.; Herrero, A.; Flores, E. Catabolic pathway of arginine in *Anabaena* involves a novel bifunctional enzyme that produces proline from arginine. *Mol. Microbiol.* **2019**, *111*, 883–897. [[CrossRef](#)] [[PubMed](#)]
98. Lee, H.; Rhee, S. Structural and mutational analyses of the bifunctional arginine dihydrolase and ornithine cyclodeaminase AgrE from the cyanobacterium *Anabaena*. *J. Biol. Chem.* **2020**, *295*, 5751–5760. [[CrossRef](#)]
99. Phogosee, S.; Hibino, T.; Kageyama, H.; Waditee-Sirisattha, R. Bifunctional alanine dehydrogenase from the halotolerant cyanobacterium *Aphanothece halophytica*: Characterization and molecular properties. *Arch. Microbiol.* **2018**, *200*, 719–727. [[CrossRef](#)] [[PubMed](#)]
100. Martínez-Noël, G.M.A.; Cumino, A.C.; De Los Angeles Kolman, M.; Salerno, G.L. First evidence of sucrose biosynthesis by single cyanobacterial bimodular proteins. *FEBS Lett.* **2013**, *587*, 1669–1674. [[CrossRef](#)]
101. Jakopitsch, C.; Regelsberger, G.; Furtmüller, P.G.; Rüter, F.; Peschek, G.A.; Obinger, C. Catalase-peroxidase from *Synechocystis* is capable of chlorination and bromination reactions. *Biochem. Biophys. Res. Commun.* **2001**, *287*, 682–687. [[CrossRef](#)]
102. Jakopitsch, C.; Auer, M.; Ivancich, A.; Rüter, F.; Furtmüller, P.G.; Obinger, C. Total conversion of bifunctional catalase-peroxidase (KatG) to monofunctional peroxidase by exchange of a conserved distal side tyrosine. *J. Biol. Chem.* **2003**, *278*, 20185–20191. [[CrossRef](#)] [[PubMed](#)]
103. Zhang, S.; Laborde, S.M.; Frankel, L.K.; Bricker, T.M. Four Novel Genes Required for Optimal Photoautotrophic Growth of the Cyanobacterium *Synechocystis* sp. Strain PCC 6803 Identified by In Vitro Transposon Mutagenesis. *J. Bacteriol.* **2004**, *186*, 875–879. [[CrossRef](#)]
104. Terauchi, K.; Sobue, R.; Furutani, Y.; Aoki, R.; Fujita, Y. Isolation of cyanobacterial mutants exhibiting growth defects under microoxic conditions by transposon tagging mutagenesis of *Synechocystis* sp. PCC 6803. *J. Gen. Appl. Microbiol.* **2017**, *63*, 131–138. [[CrossRef](#)]
105. Katayama, M.; Tsinoremas, N.F.; Kondo, T.; Golden, S.S. *cpmA*, a gene involved in an output pathway of the cyanobacterial circadian system. *J. Bacteriol.* **1999**, *181*, 3516–3524. [[CrossRef](#)] [[PubMed](#)]
106. Miyagishima, S.Y.; Wolk, P.P.; Osteryoung, K.W. Identification of cyanobacterial cell division genes by comparative and mutational analyses. *Mol. Microbiol.* **2005**, *56*, 126–143. [[CrossRef](#)] [[PubMed](#)]
107. Watabe, K.; Mimuro, M.; Tsuchiya, T. Development of a high-frequency in vivo transposon mutagenesis system for *Synechocystis* sp. PCC 6803 and *Synechococcus elongatus* PCC 7942. *Plant Cell Physiol.* **2014**, *55*, 2017–2028. [[CrossRef](#)] [[PubMed](#)]
108. Rubin, B.E.; Wetmore, K.M.; Price, M.N.; Diamond, S.; Shultzaberger, R.K.; Lowe, L.C.; Curtin, G.; Arkin, A.P.; Deutschbauer, A.; Golden, S.S. The essential gene set of a photosynthetic organism. *Proc. Natl. Acad. Sci. USA* **2015**, *112*. [[CrossRef](#)]

109. Ozaki, H.; Ikeuchi, M.; Ogawa, T.; Fukuzawa, H.; Sonoike, K. Large-scale analysis of chlorophyll fluorescence kinetics in *Synechocystis* sp. PCC 6803: Identification of the factors involved in the modulation of photosystem stoichiometry. *Plant Cell Physiol.* **2007**, *48*, 451–458. [[CrossRef](#)]
110. Kato, K.; Tanaka, R.; Sano, S.; Tanaka, A.; Hosaka, H. Identification of a gene essential for protoporphyrinogen IX oxidase activity in the cyanobacterium *Synechocystis* sp. PCC6803. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 16649–16654. [[CrossRef](#)]
111. Zhang, S.; Frankel, L.K.; Bricker, T.M. The SlI0606 protein is required for photosystem II assembly/stability in the cyanobacterium *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* **2010**, *285*, 32047–32054. [[CrossRef](#)]
112. Shibata, M.; Ohkawa, H.; Kaneko, T.; Fukuzawa, H.; Tabata, S.; Kaplan, A.; Ogawa, T. Distinct constitutive and low-CO₂-induced CO₂ uptake systems in cyanobacteria: Genes involved and their phylogenetic relationship with homologous genes in other organisms. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 11789–11794. [[CrossRef](#)]
113. Shibata, M.; Katoh, H.; Sonoda, M.; Ohkawa, H.; Shimoyama, M.; Fukuzawa, H.; Kaplan, A.; Ogawa, T. Genes essential to sodium-dependent bicarbonate transport in cyanobacteria: Function and phylogenetic analysis. *J. Biol. Chem.* **2002**, *277*, 18658–18664. [[CrossRef](#)]
114. Tyo, K.E.J.; Espinoza, F.A.; Stephanopoulos, G.; Jin, Y.S. Identification of gene disruptions for increased poly-3-hydroxybutyrate accumulation in *Synechocystis* PCC 6803. *Biotechnol. Prog.* **2009**, *25*, 1236–1243. [[CrossRef](#)] [[PubMed](#)]
115. Battchikova, N.; Wei, L.; Du, L.; Bersanini, L.; Aro, E.M.; Ma, W. Identification of novel Ssl0352 protein (NdhS), essential for efficient operation of cyclic electron transport around photosystem I, in NADPH:plastoquinone oxidoreductase (NDH-1) complexes of *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* **2011**, *286*, 36992–37001. [[CrossRef](#)]
116. Dai, H.; Zhang, L.; Zhang, J.; Mi, H.; Ogawa, T.; Ma, W. Identification of a cyanobacterial CRR6 protein, Slr1097, required for efficient assembly of NDH-1 complexes in *Synechocystis* sp. PCC 6803. *Plant J.* **2013**, *75*, 858–866. [[CrossRef](#)] [[PubMed](#)]
117. Zhang, J.; Gao, F.; Zhao, J.; Ogawa, T.; Wang, Q.; Ma, W.W. NdhP Is an exclusive subunit of large complex of NADPH dehydrogenase essential to stabilize the complex in *Synechocystis* sp. Strain PCC 6803. *J. Biol. Chem.* **2014**, *289*, 26669–26676. [[CrossRef](#)]
118. Zhao, J.; Rong, W.; Gao, F.; Ogawa, T.; Ma, W. Subunit Q is required to stabilize the large complex of NADPH dehydrogenase in *Synechocystis* sp. Strain PCC 6803. *Plant Physiol.* **2015**, *168*, 443–451. [[CrossRef](#)]
119. Gao, F.; Zhao, J.; Wang, X.; Qin, S.; Wei, L.; Ma, W. NdhV is a subunit of NADPH dehydrogenase essential for cyclic electron transport in *Synechocystis* sp. strain PCC 6803. *Plant Physiol.* **2016**, *170*, 752–760. [[CrossRef](#)] [[PubMed](#)]
120. He, Z.; Xu, M.; Wu, Y.; Lv, J.; Fu, P.; Mi, H. NdhM subunit is required for the stability and the function of NAD(P)H dehydrogenase complexes involved in CO₂ uptake in *Synechocystis* sp. strain PCC 6803. *J. Biol. Chem.* **2016**, *60*, 1374–1385. [[CrossRef](#)]
121. Wang, X.; Gao, F.; Zhang, J.; Zhao, J.; Ogawa, T.; Ma, W. A cytoplasmic protein Ssl3829 is important for NDH-1 hydrophilic arm assembly in *Synechocystis* sp. Strain PCC 6803. *Plant Physiol.* **2016**, *171*, 864–877. [[CrossRef](#)] [[PubMed](#)]
122. Voshol, G.P.; Meyer, V.; van den Hondel, C.A.M.J.J. GTP-binding protein Era: A novel gene target for biofuel production. *BMC Biotechnol.* **2015**, *15*. [[CrossRef](#)]
123. Cassier-Chauvat, C.; Chauvat, F. Function and Regulation of Ferredoxins in the Cyanobacterium, *Synechocystis* PCC6803: Recent Advances. *Life* **2014**, *4*, 666–680. [[CrossRef](#)]
124. Lelong, C.; Boekema, E.J.; Kruip, J.; Bottin, H.; Rögner, M.; Sétif, P. Characterization of a redox active cross-linked complex between cyanobacterial photosystem I and soluble ferredoxin. *EMBO J.* **1996**, *15*, 2160–2168. [[CrossRef](#)]
125. Poncelet, M.; Cassier-Chauvat, C.; Leschelle, X.; Bottin, H.; Chauvat, F. Targeted deletion and mutational analysis of the essential (2Fe-2S) plant-like ferredoxin in *Synechocystis* PCC6803 by plasmid shuffling. *Mol. Microbiol.* **1998**, *28*, 813–821. [[CrossRef](#)] [[PubMed](#)]
126. Mazouni, K.; Domain, F.; Chauvat, F.; Cassier-Chauvat, C. Expression and regulation of the crucial plant-like ferredoxin of cyanobacteria. *Mol. Microbiol.* **2003**, *49*, 1019–1029. [[CrossRef](#)] [[PubMed](#)]
127. Schorsch, M.; Kramer, M.; Goss, T.; Eisenhut, M.; Robinson, N.; Osman, D.; Wilde, A.; Sadaf, S.; Brückler, H.; Walder, L.; et al. A unique ferredoxin acts as a player in the low-iron response of photosynthetic organisms. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 12111–12120. [[CrossRef](#)]
128. Mustila, H.; Allahverdiyeva, Y.; Isojärvi, J.; Aro, E.M.; Eisenhut, M. The bacterial-type [4Fe-4S] ferredoxin 7 has a regulatory function under photooxidative stress conditions in the cyanobacterium *Synechocystis* sp. PCC 6803. *Biochim. Biophys. Acta Bioenerg.* **2014**, *1837*, 1293–1304. [[CrossRef](#)] [[PubMed](#)]
129. Hanke, G.T.; Satomi, Y.; Shinmura, K.; Takao, T.; Hase, T. A screen for potential ferredoxin electron transfer partners uncovers new, redox dependent interactions. *Biochim. Biophys. Acta Proteins Proteom.* **2011**, *1814*, 366–374. [[CrossRef](#)]
130. Angeleri, M.; Zorina, A.; Aro, E.M.; Battchikova, N. Interplay of SpkG kinase and the Slr0151 protein in the phosphorylation of ferredoxin 5 in *Synechocystis* sp. strain PCC 6803. *FEBS Lett.* **2018**, *592*, 411–421. [[CrossRef](#)]
131. Mondal, S.; Kumar, V.; Singh, S.P. Phylogenetic distribution and structural analyses of cyanobacterial glutaredoxins (Grxs). *Comput. Biol. Chem.* **2020**, *84*. [[CrossRef](#)]
132. Zaffagnini, M.; Bedhomme, M.; Marchand, C.H.; Morisse, S.; Trost, P.; Lemaire, S.D. Redox regulation in photosynthetic organisms: Focus on glutathionylation. *Antioxid. Redox Signal.* **2012**, *16*, 567–586. [[CrossRef](#)] [[PubMed](#)]
133. Sánchez-Riego, A.M.; López-Maury, L.; Florencio, F.J. Glutaredoxins are essential for stress adaptation in the cyanobacterium *Synechocystis* sp. PCC 6803. *Front. Plant Sci.* **2013**. [[CrossRef](#)] [[PubMed](#)]

134. López-Maury, L.; Sánchez-Riego, A.M.; Reyes, J.C.; Florencio, F.J. The glutathione/glutaredoxin system is essential for arsenate reduction in *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **2009**, *191*, 3534–3543. [[CrossRef](#)] [[PubMed](#)]
135. Kim, S.G.; Chung, J.S.; Sutton, R.B.; Lee, J.S.; López-Maury, L.; Lee, S.Y.; Florencio, F.J.; Lin, T.; Zabet-Moghaddam, M.; Wood, M.J.; et al. Redox, mutagenic and structural studies of the glutaredoxin/arsenate reductase couple from the cyanobacterium *Synechocystis* sp. PCC 6803. *Biochim. Biophys. Acta Proteins Proteom.* **2012**, *1824*, 392–403. [[CrossRef](#)] [[PubMed](#)]
136. Iwema, T.; Picciocchi, A.; Traore, D.A.K.; Ferrer, J.L.; Chauvat, F.; Jacquamet, L. Structural basis for delivery of the intact [Fe2S2] cluster by monothiol glutaredoxin. *Biochemistry* **2009**, *48*, 6041–6043. [[CrossRef](#)] [[PubMed](#)]
137. Deponte, M. Glutathione catalysis and the reaction mechanisms of glutathione-dependent enzymes. *Biochim. Biophys. Acta Gen. Subj.* **2013**, *1830*, 3217–3266. [[CrossRef](#)]
138. Nianiou-Obeidat, I.; Madesis, P.; Kissoudis, C.; Voulgari, G.; Chronopoulou, E.; Tsafaris, A.; Labrou, N.E. Plant glutathione transferase-mediated stress tolerance: Functions and biotechnological applications. *Plant Cell Rep.* **2017**, *36*, 791–805. [[CrossRef](#)]
139. Oakley, A. Glutathione transferases: A structural perspective. *Drug Metab. Rev.* **2011**, *43*, 138–151. [[CrossRef](#)]
140. Wiktelius, E.; Stenberg, G. Novel class of glutathione transferases from cyanobacteria exhibit high catalytic activities towards naturally occurring isothiocyanates. *Biochem. J.* **2007**, *406*, 115–123. [[CrossRef](#)]
141. Naciyar, M.S.; Karthick, L.; Prakasam, P.A.; Deviram, G.; Uma, L.; Prabakaran, D.; Saha, S.K. Diversity of glutathione S-transferases (GSTs) in cyanobacteria with reference to their structures, substrate recognition and catalytic functions. *Microorganisms* **2020**, *8*, 712. [[CrossRef](#)]
142. Pandey, T.; Chhetri, G.; Chinta, R.; Kumar, B.; Singh, D.B.; Tripathi, T.; Singh, A.K. Functional classification and biochemical characterization of a novel rho class glutathione S-transferase in *Synechocystis* PCC 6803. *FEBS Open Bio* **2015**, *5*, 1–7. [[CrossRef](#)] [[PubMed](#)]
143. Pandey, T.; Singh, S.K.; Chhetri, G.; Tripathi, T.; Singh, A.K. Characterization of a highly pH stable Chi-class glutathione S-transferase from *Synechocystis* PCC 6803. *PLoS ONE* **2015**, *10*, e0126811. [[CrossRef](#)] [[PubMed](#)]
144. Pandey, T.; Shukla, R.; Shukla, H.; Sonkar, A.; Tripathi, T.; Singh, A.K. A combined biochemical and computational studies of the rho-class glutathione s-transferase sll1545 of *Synechocystis* PCC 6803. *Int. J. Biol. Macromol.* **2017**, *94*, 378–385. [[CrossRef](#)]
145. Kammerscheit, X.; Hecker, A.; Rouhier, N.; Chauvat, F.; Cassier-Chauvat, C. Methylglyoxal Detoxification Revisited: Role of Glutathione Transferase in Model Cyanobacterium *Synechocystis* sp. Strain PCC 6803. *MBio* **2020**, *11*. [[CrossRef](#)] [[PubMed](#)]
146. Stanier, R.Y.; Kunisawa, R.; Mandel, M.; Cohen-Bazire, G. Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriol. Rev.* **1971**, *35*, 171–205. [[CrossRef](#)] [[PubMed](#)]
147. Rippka, R.; Deruelles, J.; Waterbury, J.B.; Herdman, M.; Stanier, R.Y. Generic assignments, strain histories and properties of pure cultures of cyano- bacteria. *Journal of General Microbiology* **11**: 1-61. *J. Gen. Microbiol.* **1979**, *111*, 1–61. [[CrossRef](#)]
148. Niu, T.C.; Lin, G.M.; Xie, L.R.; Wang, Z.Q.; Xing, W.Y.; Zhang, J.Y.; Zhang, C.C. Expanding the Potential of CRISPR-Cpf1-Based Genome Editing Technology in the Cyanobacterium *Anabaena* PCC 7120. *ACS Synth. Biol.* **2019**, *8*, 170–180. [[CrossRef](#)]
149. Yu, J.; Liberton, M.; Cliften, P.F.; Head, R.D.; Jacobs, J.M.; Smith, R.D.; Koppelaar, D.W.; Brand, J.J.; Pakrasi, H.B. *Synechococcus elongatus* UTEX 2973, a fast growing cyanobacterial chassis for biosynthesis using light and CO₂. *Sci. Rep.* **2015**, *5*. [[CrossRef](#)]
150. Jaiswal, D.; Sengupta, A.; Sohoni, S.; Sengupta, S.; Phadnavis, A.G.; Pakrasi, H.B.; Wangikar, P.P. Genome Features and Biochemical Characteristics of a Robust, Fast Growing and Naturally Transformable Cyanobacterium *Synechococcus elongatus* PCC 11801 Isolated from India. *Sci. Rep.* **2018**, *8*. [[CrossRef](#)]
151. Jaiswal, D.; Sengupta, A.; Sengupta, S.; Madhu, S.; Pakrasi, H.B.; Wangikar, P.P. A Novel Cyanobacterium *Synechococcus elongatus* PCC 11802 has Distinct Genomic and Metabolomic Characteristics Compared to its Neighbor PCC 11801. *Sci. Rep.* **2020**, *10*. [[CrossRef](#)]
152. Włodarczyk, A.; Selão, T.T.; Norling, B.; Nixon, P.J. Newly discovered *Synechococcus* sp. PCC 11901 is a robust cyanobacterial strain for high biomass production. *Commun. Biol.* **2020**, *3*. [[CrossRef](#)] [[PubMed](#)]
153. Tan, X.; Hou, S.; Song, K.; Georg, J.; Klähn, S.; Lu, X.; Hess, W.R. The primary transcriptome of the fast-growing cyanobacterium *Synechococcus elongatus* UTEX 2973. *Biotechnol. Biofuels* **2018**, *11*. [[CrossRef](#)] [[PubMed](#)]
154. Lou, W.; Tan, X.; Song, K.; Zhang, S.; Luan, G.; Li, C.; Lu, X. A specific single nucleotide polymorphism in the ATP synthase gene significantly improves environmental stress tolerance of *Synechococcus elongatus* PCC 7942. *Appl. Environ. Microbiol.* **2018**, *84*. [[CrossRef](#)] [[PubMed](#)]
155. Bachin, D.; Nazarenko, L.V.; Mironov, K.S.; Pisareva, T.; Allakhverdiev, S.I.; Los, D.A. Mechanosensitive ion channel MscL controls ionic fluxes during cold and heat stress in *Synechocystis*. *FEMS Microbiol. Lett.* **2015**, *362*. [[CrossRef](#)] [[PubMed](#)]
156. Li, S.; Sun, T.; Xu, C.; Chen, L.; Zhang, W. Development and optimization of genetic toolboxes for a fast-growing cyanobacterium *Synechococcus elongatus* UTEX 2973. *Metab. Eng.* **2018**, *48*, 163–174. [[CrossRef](#)]
157. Ungerer, J.; Wendt, K.E.; Hendry, J.I.; Maranas, C.D.; Pakrasi, H.B. Comparative genomics reveals the molecular determinants of rapid growth of the cyanobacterium *Synechococcus elongatus* UTEX 2973. *Proc. Natl. Acad. Sci. USA* **2018**, *115*. [[CrossRef](#)] [[PubMed](#)]
158. Zhang, M.; Qiao, C.; Luan, G.; Luo, Q.; Lu, X. Systematic Identification of Target Genes for Cellular Morphology Engineering in *Synechococcus elongatus* PCC7942. *Front. Microbiol.* **2020**, *11*. [[CrossRef](#)]
159. Golden, S.S. Principles of rhythmicity emerging from cyanobacteria. *Eur. J. Neurosci.* **2020**, *51*, 13–18. [[CrossRef](#)]
160. MacCready, J.S.; Basalla, J.L.; Vecchiarelli, A.G. Origin and Evolution of Carboxysome Positioning Systems in Cyanobacteria. *Mol. Biol. Evol.* **2020**. [[CrossRef](#)]

161. Labarre, J.; Chauvat, F.; Thuriaux, P. Insertional mutagenesis by random cloning of antibiotic resistance genes into the genome of the cyanobacterium *Synechocystis* strain PCC 6803. *J. Bacteriol.* **1989**, *171*, 3449–3457. [[CrossRef](#)]
162. Zerulla, K.; Ludt, K.; Soppa, J. The ploidy level of *Synechocystis* sp. PCC 6803 is highly variable and is influenced by growth phase and by chemical and physical external parameters. *Microbiology (UK)* **2016**, *162*, 730–739. [[CrossRef](#)] [[PubMed](#)]
163. Chauvat, F.; De Vries, L.; Van der Ende, A.; Van Arkel, G.A. A host-vector system for gene cloning in the cyanobacterium *Synechocystis* PCC 6803. *MGG Mol. Gen. Genet.* **1986**, *204*, 185–191. [[CrossRef](#)]
164. Yang, X.; McFadden, B.A. A small plasmid, pCA2.4, from the cyanobacterium *Synechocystis* sp. strain PCC 6803 encodes a Rep protein and replicates by a rolling circle mechanism. *J. Bacteriol.* **1993**, *175*, 3981–3991. [[CrossRef](#)]
165. Yang, X.; McFadden, B.A. The complete DNA sequence and replication analysis of the plasmid pCB2.4 from the cyanobacterium *Synechocystis* PCC 6803. *Plasmid* **1994**, *31*, 131–137. [[CrossRef](#)] [[PubMed](#)]
166. Xu, W.; McFadden, B.A. Sequence analysis of plasmid pCC5.2 from cyanobacterium *Synechocystis* PCC 6803 that replicates by a rolling circle mechanism. *Plasmid* **1997**, *37*, 95–104. [[CrossRef](#)] [[PubMed](#)]
167. Kaneko, T.; Nakamura, Y.; Sasamoto, S.; Watanabe, A.; Kohara, M.; Matsumoto, M.; Shimpo, S.; Yamada, M.; Tabata, S. Structural analysis of four large plasmids harboring in a unicellular cyanobacterium, *Synechocystis* sp. PCC 6803. *DNA Res.* **2003**, *10*, 221–228. [[CrossRef](#)]
168. Berla, B.M.; Pakrasi, H.B. Upregulation of plasmid genes during stationary phase in *Synechocystis* sp. strain PCC 6803, a cyanobacterium. *Appl. Environ. Microbiol.* **2012**, *18*. [[CrossRef](#)]
169. Castets, A.M.; Houmard, J.; Tandeau de Marsac, N. Is cell motility a plasmid-encoded function in the cyanobacterium *Synechocystis* 6803? *FEMS Microbiol. Lett.* **1986**, *37*, 277–281. [[CrossRef](#)]
170. Kopfmann, S.; Hess, W.R. Toxin-antitoxin systems on the large defense plasmid pSYSA of *Synechocystis* sp. pCC 6803. *J. Biol. Chem.* **2013**, *288*, 7399–7409. [[CrossRef](#)]
171. Kopfmann, S.; Roesch, S.K.; Hess, W.R. Type II toxin-antitoxin systems in the unicellular cyanobacterium *Synechocystis* sp. PCC 6803. *Toxins* **2016**, *8*, 228. [[CrossRef](#)]
172. Paynter, J.J.; Andres-Enguix, I.; Fowler, P.W.; Tottey, S.; Cheng, W.; Enkvetchakul, D.; Bavro, V.N.; Kusakabe, Y.; Sansom, M.S.P.; Robinson, N.J.; et al. Functional complementation and genetic deletion studies of KirBac channels: Activatory mutations highlight gating-sensitive domains. *J. Biol. Chem.* **2010**, *285*, 40754–40761. [[CrossRef](#)] [[PubMed](#)]
173. Giner-Lamia, J.; López-Maury, L.; Reyes, J.C.; Florencio, F.J. The CopRS two-component system is responsible for resistance to copper in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.* **2012**, *159*, 1806–1818. [[CrossRef](#)] [[PubMed](#)]
174. De Alvarenga, L.V.; Hess, W.R.; Hagemann, M. AcnSP—A Novel Small Protein Regulator of Aconitase Activity in the Cyanobacterium *Synechocystis* sp. PCC 6803. *Front. Microbiol.* **2020**, *11*. [[CrossRef](#)] [[PubMed](#)]
175. Bird, A.J.; Turner-Cavet, J.S.; Lakey, J.H.; Robinson, N.J. A carboxyl-terminal Cys2/His2-type zinc-finger motif in DNA primase influences DNA content in *Synechococcus* PCC 7942. *J. Biol. Chem.* **1998**, *273*, 21246–21252. [[CrossRef](#)]
176. Griese, M.; Lange, C.; Soppa, J. Ploidy in cyanobacteria. *FEMS Microbiol. Lett.* **2011**, *323*, 124–131. [[CrossRef](#)]
177. Watanabe, S.; Ohbayashi, R.; Kanesaki, Y.; Saito, N.; Chibazakura, T.; Soga, T.; Yoshikawa, H. Intensive DNA replication and metabolism during the lag phase in Cyanobacteria. *PLoS ONE* **2015**, *10*, e0136800. [[CrossRef](#)]
178. Van Den Hondel, C.A.M.J.J.; Verbeek, S.; van Der Ende, A.; Weisbeek, P.J.; Borrias, W.E.; van Arkel, G.A. Introduction of transposon Tn901 into a plasmid of *Anacystis nidulans*: Preparation for cloning in cyanobacteria. *Proc. Natl. Acad. Sci. USA* **1980**, *77*, 1570–1574. [[CrossRef](#)] [[PubMed](#)]
179. Chen, Y.; Kay Holtman, C.; Magnuson, R.D.; Youderian, P.A.; Golden, S.S. The complete sequence and functional analysis of pANL, the large plasmid of the unicellular freshwater cyanobacterium *Synechococcus elongatus* PCC 7942. *Plasmid* **2008**, *59*, 176–192. [[CrossRef](#)]
180. Roberts, T.M.; Koths, K.E. The blue-green alga *Agmenellum quadruplicatum* contains covalently closed DNA circles. *Cell* **1976**, *9*, 551–557. [[CrossRef](#)]
181. Akiyama, H.; Kanai, S.; Hirano, M.; Miyasaka, H. Nucleotide sequence of plasmid pAQ1 of marine cyanobacterium *Synechococcus* sp. PCC7002. *DNA Res.* **1998**, *5*, 127–129. [[CrossRef](#)]
182. Kuhlemeier, C.J.; Borrias, W.E.; van den Hondel, C.A.M.J.J.; van Arkel, G.A. Vectors for cloning in cyanobacteria: Construction and characterization of two recombinant plasmids capable of transformation to *Escherichia coli* K12 and *Anacystis nidulans* R2. *MGG Mol. Gen. Genet.* **1981**, *184*, 249–254. [[CrossRef](#)]
183. Chauvat, F.; Astier, C.; Vedel, F.; Joset-Espardellier, F. Transformation in the cyanobacterium *Synechococcus* R2: Improvement of efficiency; Role of the pUH24 plasmid. *MGG Mol. Gen. Genet.* **1983**, *191*, 39–45. [[CrossRef](#)] [[PubMed](#)]
184. Kaneko, T.; Nakamura, Y.; Wolk, C.P.; Kuritz, T.; Sasamoto, S.; Watanabe, A.; Iriguchi, M.; Ishikawa, A.; Kawashima, K.; Kimura, T.; et al. Complete genomic sequence of the filamentous nitrogen-fixing cyanobacterium *Anabaena* sp. strain PCC 7120. *DNA Res.* **2001**, *8*, 205–213. [[CrossRef](#)]
185. Ludwig, M.; Bryant, D.A. Acclimation of the global transcriptome of the cyanobacterium *Synechococcus* sp. strain PCC 7002 to nutrient limitations and different nitrogen sources. *Front. Microbiol.* **2012**, *11*. [[CrossRef](#)]
186. Labarre, J.; Thuriaux, P.; Chauvat, F. Genetic analysis of amino acid transport in the facultatively heterotrophic cyanobacterium *Synechocystis* sp. strain 6803. *J. Bacteriol.* **1987**, *169*, 4668–4673. [[CrossRef](#)] [[PubMed](#)]
187. Pérez, A.A.; Liu, Z.; Rodionov, D.A.; Li, Z.; Bryant, D.A. Complementation of cobalamin auxotrophy in *Synechococcus* sp. strain PCC 7002 and validation of a putative cobalamin riboswitch in vivo. *J. Bacteriol.* **2016**, *198*, 2743–2752. [[CrossRef](#)] [[PubMed](#)]

188. Sakamoto, T.; Delgaizo, V.B.; Bryant, D.A. Growth on urea can trigger death and peroxidation of the cyanobacterium *Synechococcus* sp. strain PCC 7002. *Appl. Environ. Microbiol.* **1998**, *64*, 2361–2366. [[CrossRef](#)]
189. Valladares, A.; Montesinos, M.L.; Herrero, A.; Flores, E. An ABC-type, high-affinity urea permease identified in cyanobacteria. *Mol. Microbiol.* **2002**, *43*, 703–715. [[CrossRef](#)]
190. Veaudor, T.; Cassier-Chauvat, C.; Chauvat, F. Genomics of Urea Transport and Catabolism in Cyanobacteria: Biotechnological Implications. *Front. Microbiol.* **2019**, *10*. [[CrossRef](#)]
191. Kirsch, F.; Klähn, S.; Hagemann, M. Salt-Regulated Accumulation of the Compatible Solutes Sucrose and Glucosylglycerol in Cyanobacteria and Its Biotechnological Potential. *Front. Microbiol.* **2019**, *10*. [[CrossRef](#)]
192. Årstøl, E.; Hohmann-Marriott, M.F. Cyanobacterial siderophores—physiology, structure, biosynthesis, and applications. *Mar. Drugs* **2019**, *17*, 281. [[CrossRef](#)] [[PubMed](#)]
193. Onizuka, T.; Endo, S.; Akiyama, H.; Kanai, S.; Hirano, M.; Yokota, A.; Tanaka, S.; Miyasaka, H. The *rbcX* gene product promotes the production and assembly of ribulose-1,5-bisphosphate carboxylase/oxygenase of *Synechococcus* sp. PCC7002 in *Escherichia coli*. *Plant Cell Physiol.* **2004**, *45*, 1390–1395. [[CrossRef](#)]
194. Emlyn-Jones, D.; Woodger, F.J.; Price, G.D.; Whitney, S.M. RbcX can function as a Rubisco chaperonin, but is non-essential in *Synechococcus* PCC7942. *Plant Cell Physiol.* **2006**, *47*, 1630–1640. [[CrossRef](#)] [[PubMed](#)]
195. Bracher, A.; Whitney, S.M.; Hartl, F.U.; Hayer-Hartl, M. Biogenesis and Metabolic Maintenance of Rubisco. *Annu. Rev. Plant Biol.* **2017**, *68*, 29–60. [[CrossRef](#)]
196. Helman, Y.; Tchernov, D.; Reinhold, L.; Shibata, M.; Ogawa, T.; Schwarz, R.; Ohad, I.; Kaplan, A. Genes encoding A-type flavoproteins are essential for photoreduction of O₂ in cyanobacteria. *Curr. Biol.* **2003**, *13*, 230–235. [[CrossRef](#)]
197. Vicente, J.B.; Gomes, C.M.; Wasserfallen, A.; Teixeira, M. Module fusion in an A-type flavoprotein from the cyanobacterium *Synechocystis* condenses a multiple-component pathway in a single polypeptide chain. *Biochem. Biophys. Res. Commun.* **2002**, *294*, 82–87. [[CrossRef](#)]
198. Allahverdiyeva, Y.; Mustila, H.; Ermakova, M.; Bersanini, L.; Richaud, P.; Ajlani, G.; Battchikova, N.; Cournac, L.; Aro, E.M. Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and photosynthesis under fluctuating light. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 4111–4116. [[CrossRef](#)]
199. Zhang, P.; Allahverdiyeva, Y.; Eisenhut, M.; Aro, E.M. Flavodiiron proteins in oxygenic photosynthetic organisms: Photoprotection of photosystem II by Flv2 and Flv4 in *Synechocystis* sp. PCC 6803. *PLoS ONE* **2009**, *4*, e5331. [[CrossRef](#)] [[PubMed](#)]
200. Shimakawa, G.; Shaku, K.; Nishi, A.; Hayashi, R.; Yamamoto, H.; Sakamoto, K.; Makino, A.; Miyake, C. FLAVODIIRON2 and FLAVODIIRON4 proteins mediate an oxygen-dependent alternative electron flow in *Synechocystis* sp. pcc 6803 under CO₂-limited conditions. *Plant Physiol.* **2015**, *167*, 472–480. [[CrossRef](#)]
201. Fujisawa, T.; Okamoto, S.; Katayama, T.; Nakao, M.; Yoshimura, H.; Kajiya-Kanegae, H.; Yamamoto, S.; Yano, C.; Yanaka, Y.; Maita, H.; et al. CyanoBase and RhizoBase: Databases of manually curated annotations for cyanobacterial and rhizobial genomes. *Nucleic Acids Res.* **2014**, *42*. [[CrossRef](#)]
202. Jittawuttipoka, T.; Planchon, M.; Spalla, O.; Benzerara, K.; Guyot, F.; Cassier-Chauvat, C.; Chauvat, F. Multidisciplinary Evidences that *Synechocystis* PCC6803 Exopolysaccharides Operate in Cell Sedimentation and Protection against Salt and Metal Stresses. *PLoS ONE* **2013**, *8*, e55564. [[CrossRef](#)]
203. Planchon, M.; Jittawuttipoka, T.; Cassier-Chauvat, C.; Guyot, F.; Gelabert, A.; Benedetti, M.F.; Chauvat, F.; Spalla, O. Exopolysaccharides protect *Synechocystis* against the deleterious effects of Titanium dioxide nanoparticles in natural and artificial waters. *J. Colloid Interface Sci.* **2013**, *405*, 35–43. [[CrossRef](#)] [[PubMed](#)]
204. Naveed, S.; Yu, Q.; Zhang, C.; Ge, Y. Extracellular polymeric substances alter cell surface properties, toxicity, and accumulation of arsenic in *Synechocystis* PCC6803. *Environ. Pollut.* **2020**, *261*. [[CrossRef](#)] [[PubMed](#)]
205. Yu, R.; Chai, H.; Yu, Z.; Wu, X.; Liu, Y.; Shen, L.; Li, J.; Ye, J.; Liu, D.; Ma, T.; et al. Behavior and mechanism of cesium biosorption from aqueous solution by living *Synechococcus* PCC7002. *Microorganisms* **2020**, *8*, 491. [[CrossRef](#)] [[PubMed](#)]
206. Schatz, D.; Nagar, E.; Sendersky, E.; Parnasa, R.; Zilberman, S.; Carmeli, S.; Mastai, Y.; Shimoni, E.; Klein, E.; Yeger, O.; et al. Self-suppression of biofilm formation in the cyanobacterium *Synechococcus elongatus*. *Environ. Microbiol.* **2013**, *15*, 1786–1794. [[CrossRef](#)]
207. Muzzopappa, F.; Kirilovsky, D. Changing Color for Photoprotection: The Orange Carotenoid Protein. *Trends Plant Sci.* **2020**, *25*, 92–104. [[CrossRef](#)]
208. Kämäräinen, J.; Knoop, H.; Stanford, N.J.; Guerrero, F.; Akhtar, M.K.; Aro, E.M.; Steuer, R.; Jones, P.R. Physiological tolerance and stoichiometric potential of cyanobacteria for hydrocarbon fuel production. *J. Biotechnol.* **2012**, *162*, 67–74. [[CrossRef](#)]
209. Gupta, A.; Bhagwat, S.G.; Sainis, J.K. *Synechococcus elongatus* PCC 7942 is more tolerant to chromate as compared to *Synechocystis* sp. PCC 6803. *BioMetals* **2013**, *26*, 309–319. [[CrossRef](#)] [[PubMed](#)]
210. Gupta, A.; Ballal, A. Unraveling the mechanism responsible for the contrasting tolerance of *Synechocystis* and *Synechococcus* to Cr(VI): Enzymatic and non-enzymatic antioxidants. *Aquat. Toxicol.* **2015**, *164*, 118–125. [[CrossRef](#)] [[PubMed](#)]
211. Kämäräinen, J.; Nylund, M.; Aro, E.M.; Kallio, P. Comparison of ethanol tolerance between potential cyanobacterial production hosts. *J. Biotechnol.* **2018**, *283*, 140–145. [[CrossRef](#)]
212. Ikeuchi, M.; Tabata, S. *Synechocystis* sp. PCC 6803—A useful tool in the study of the genetics of cyanobacteria. *Photosynth. Res.* **2001**, *70*, 73–83. [[CrossRef](#)] [[PubMed](#)]

213. Tajima, N.; Sato, S.; Maruyama, F.; Kaneko, T.; Sasaki, N.V.; Kurokawa, K.; Ohta, H.; Kanesaki, Y.; Yoshikawa, H.; Tabata, S.; et al. Genomic structure of the cyanobacterium *Synechocystis* sp. PCC 6803 strain GT-S. *DNA Res.* **2011**, *18*, 393–399. [[CrossRef](#)]
214. Morris, J.N.; Crawford, T.S.; Jeffs, A.; Stockwell, P.A.; Eaton-Rye, J.J.; Summerfield, T.C. Whole genome re-sequencing of two “wild-type” strains of the model cyanobacterium *Synechocystis* sp. PCC 6803. *N. Z. J. Bot.* **2014**, *52*, 36–47. [[CrossRef](#)]
215. Trautmann, D.; Voß, B.; Wilde, A.; Al-Babili, S.; Hess, W.R. Microevolution in cyanobacteria: Re-sequencing a motile substrain of *Synechocystis* sp. PCC 6803. *DNA Res.* **2012**, *19*, 435–448. [[CrossRef](#)]
216. Kanesaki, Y.; Shiwa, Y.; Tajima, N.; Suzuki, M.; Watanabe, S.; Sato, N.; Ikeuchi, M.; Yoshikawa, H. Identification of substrain-specific mutations by massively parallel whole-genome resequencing of *Synechocystis* sp. PCC 6803. *DNA Res.* **2012**, *19*, 67–79. [[CrossRef](#)] [[PubMed](#)]
217. Tichý, M.; Bečková, M.; Kopečná, J.; Noda, J.; Sobotka, R.; Komenda, J. Strain of *Synechocystis* PCC 6803 with aberrant assembly of photosystem II contains tandem duplication of a large chromosomal region. *Front. Plant Sci.* **2016**, *7*. [[CrossRef](#)]
218. Ding, Q.; Chen, G.; Wang, Y.; Wei, D. Identification of specific variations in a non-motile strain of cyanobacterium *Synechocystis* sp. PCC 6803 originated from ATCC 27184 by whole genome resequencing. *Int. J. Mol. Sci.* **2015**, *16*, 24081–24093. [[CrossRef](#)]
219. Morris, J.N.; Eaton-Rye, J.J.; Summerfield, T.C. Phenotypic variation in wild-type substrains of the model cyanobacterium *Synechocystis* sp. PCC 6803. *N. Z. J. Bot.* **2017**, 25–35. [[CrossRef](#)]
220. Williams, J.G.K. Construction of Specific Mutations in Photosystem II Photosynthetic Reaction Center by Genetic Engineering Methods in *Synechocystis* 6803. *Methods Enzymol.* **1988**, *167*, 766–778. [[CrossRef](#)]
221. Zavřel, T.; Očenášová, P.; Červený, J. Phenotypic characterization of *Synechocystis* sp. PCC 6803 substrains reveals differences in sensitivity to abiotic stress. *PLoS ONE* **2017**, *12*, e0189130. [[CrossRef](#)]
222. Shestakov, S.V.; Khyen, N.T. Evidence for genetic transformation in blue-green alga *Anacystis nidulans*. *MGG Mol. Gen. Genet.* **1970**, *107*. [[CrossRef](#)] [[PubMed](#)]
223. Stevens, S.E.; Porter, R.D. Transformation in *Agmenellum quadruplicatum*. *Proc. Natl. Acad. Sci. USA* **1980**, *77*, 6052–6056. [[CrossRef](#)]
224. Lambert, D.H.; Stevens, S.E. Photoheterotrophic growth of *Agmenellum quadruplicatum* PR-6. *J. Bacteriol.* **1986**, *165*. [[CrossRef](#)] [[PubMed](#)]
225. Grigorieva, G.; Shestakov, S. Transformation in the cyanobacterium *Synechocystis* sp. 6803. *FEMS Microbiol. Lett.* **1982**, *13*, 367–370. [[CrossRef](#)]
226. Orkwiszewski, K.G.; Kaney, A.R. Genetic transformation of the blue-green bacterium, *Anacystis nidulans*. *Arch. Microbiol.* **1974**, *98*, 31–37. [[CrossRef](#)] [[PubMed](#)]
227. Kufryk, G.I.; Sachet, M.; Schmetterer, G.; Vermaas, W.F.J. Transformation of the cyanobacterium *Synechocystis* sp. PCC 6803 as a tool for genetic mapping: Optimization of efficiency. *FEMS Microbiol. Lett.* **2002**, *206*, 215–219. [[CrossRef](#)] [[PubMed](#)]
228. Wendt, K.E.; Pakrasi, H.B. Genomics approaches to deciphering natural transformation in cyanobacteria. *Front. Microbiol.* **2019**, *10*. [[CrossRef](#)] [[PubMed](#)]
229. Taton, A.; Erikson, C.; Yang, Y.; Rubin, B.E.; Rifkin, S.A.; Golden, J.W.; Golden, S.S. The circadian clock and darkness control natural competence in cyanobacteria. *Nat. Commun.* **2020**, *11*. [[CrossRef](#)]
230. Essich, E.; Stevens, E.; Porter, R.D. Chromosomal transformation in the cyanobacterium *Agmenellum quadruplicatum*. *J. Bacteriol.* **1990**, *172*, 1916–1922. [[CrossRef](#)]
231. Williams, J.G.K.; Szalay, A.A. Stable integration of foreign DNA into the chromosome of the cyanobacterium *Synechococcus* R2. *Gene* **1983**, *24*, 37–51. [[CrossRef](#)]
232. Buzby, J.S.; Porter, R.D.; Stevens, S.E. Expression of the *Escherichia coli* lacZ gene on a plasmid vector in a cyanobacterium. *Science* **1985**, *230*, 805–807. [[CrossRef](#)] [[PubMed](#)]
233. Debus, R.J.; Barry, B.A.; Babcock, G.T.; McIntosh, L. Site-directed mutagenesis identifies a tyrosine radical involved in the photosynthetic oxygen-evolving system. *Proc. Natl. Acad. Sci. USA* **1988**, *85*. [[CrossRef](#)]
234. De Vos, W.M.; Venema, G.; Canosi, U.; Trautner, T.A. Plasmid transformation in *Bacillus subtilis*: Fate of plasmid DNA. *MGG Mol. Gen. Genet.* **1981**, *181*, 424–433. [[CrossRef](#)] [[PubMed](#)]
235. Jansson, C.; Debus, R.J.; Osiewacz, H.D.; Gurevitz, M.; McIntosh, L. Construction of an Obligate Photoheterotrophic Mutant of the Cyanobacterium *Synechocystis* 6803. *Plant Physiol.* **1987**, *85*, 1021–1025. [[CrossRef](#)]
236. Pakrasi, H.B.; Williams, J.G.; Arntzen, C.J. Targeted mutagenesis of the psbE and psbF genes blocks photosynthetic electron transport: Evidence for a functional role of cytochrome b559 in photosystem II. *EMBO J.* **1988**, *7*, 325–332. [[CrossRef](#)] [[PubMed](#)]
237. Vermaas, W.F.J.; Ikeuchi, M.; Inoue, Y. Protein composition of the photosystem II core complex in genetically engineered mutants of the cyanobacterium *Synechocystis* sp. PCC 6803. *Photosynth. Res.* **1988**, *17*, 97–113. [[CrossRef](#)]
238. Chitnis, P.R.; Reilly, P.A.; Miedel, M.C.; Nelson, N. Structure and targeted mutagenesis of the gene encoding 8-kDa subunit of photosystem I from the cyanobacterium *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* **1989**, *264*, 18374–18380. [[CrossRef](#)]
239. Chauvat, F.; Rouet, P.; Bottin, H.; Boussac, A. Mutagenesis by random cloning of an *Escherichia coli* kanamycin resistance gene into the genome of the cyanobacterium *Synechocystis* PCC 6803: Selection of mutants defective in photosynthesis. *MGG Mol. Gen. Genet.* **1989**, *216*, 51–59. [[CrossRef](#)]
240. Muller, E.G.; Buchanan, B.B. Thioredoxin is essential for photosynthetic growth. The thioredoxin m gene of *Anacystis nidulans*. *J. Biol. Chem.* **1989**, *264*, 4008–4014. [[CrossRef](#)]
241. Golden, S.S.; Brusslan, J.; Haselkorn, R. Expression of a family of psbA genes encoding a photosystem II polypeptide in the cyanobacterium *Anacystis nidulans* R2. *EMBO J.* **1986**, *5*, 2789–2798. [[CrossRef](#)] [[PubMed](#)]

242. Golden, S.S.; Cho, D.S.; Nalty, M.S. Two functional psbD genes in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *J. Bacteriol.* **1989**, *171*, 4707–4713. [[CrossRef](#)]
243. Omata, T.; Carlson, T.J.; Ogawa, T.; Pierce, J. Sequencing and modification of the gene encoding the 42-kilodalton protein in the cytoplasmic membrane of *Synechococcus* PCC 7942. *Plant Physiol.* **1990**, *93*, 305–312. [[CrossRef](#)]
244. Laudenbach, D.E.; Herbert, S.K.; McDowell, C.; Fork, D.C.; Grossman, A.R.; Straus, N.A. Cytochrome c-553 is not required for photosynthetic activity in the cyanobacterium *Synechococcus*. *Plant Cell* **1990**, *2*, 913–924. [[CrossRef](#)]
245. Bustos, S.A.; Golden, S.S. Light-regulated expression of the psbD gene family in *Synechococcus* sp. strain PCC 7942: Evidence for the role of duplicated psbD genes in cyanobacteria. *MGG Mol. Gen. Genet.* **1992**. [[CrossRef](#)]
246. Swanson, R.V.; Zhou, J.; Leary, J.A.; Williams, T.; De Lorimier, R.; Bryant, D.A.; Glazer, A.N. Characterization of phycocyanin produced by cpcE and cpcF mutants and identification of an intergenic suppressor of the defect in bilin attachment. *J. Biol. Chem.* **1992**, *267*, 16146–16154. [[CrossRef](#)]
247. Zhou, J.; Gasparich, G.E.; Stirewalt, V.L.; De Lorimier, R.; Bryant, D.A. The cpcE and cpcF genes of *Synechococcus* sp. PCC 7002. Construction and phenotypic characterization of interposon mutants. *J. Biol. Chem.* **1992**, *267*, 16138–16145. [[CrossRef](#)]
248. Schirmacher, A.M.; Hanamghar, S.S. Function and Benefits of Natural Competence in Cyanobacteria: From Ecology to Targeted Manipulation. *Life* **2020**, *10*, 249. [[CrossRef](#)] [[PubMed](#)]
249. Armshaw, P.; Carey, D.; Sheahan, C.; Pembroke, J.T. Utilising the native plasmid, pCA2.4, from the cyanobacterium *Synechocystis* sp. strain PCC6803 as a cloning site for enhanced product production. *Biotechnol. Biofuels* **2015**, *8*. [[CrossRef](#)] [[PubMed](#)]
250. Kopf, M.; Hess, W.R. Regulatory RNAs in photosynthetic cyanobacteria. *FEMS Microbiol. Rev.* **2015**, *39*, 301–315. [[CrossRef](#)] [[PubMed](#)]
251. Ng, A.H.; Berla, B.M.; Pakrasi, H.B. Fine-tuning of photoautotrophic protein production by combining promoters and neutral sites in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Appl. Environ. Microbiol.* **2015**, *81*, 6857–6863. [[CrossRef](#)] [[PubMed](#)]
252. Pinto, F.; Pacheco, C.C.; Oliveira, P.; Montagud, A.; Landels, A.; Couto, N.; Wright, P.C.; Urchueguía, J.F.; Tamagnini, P. Improving a *Synechocystis*-based photoautotrophic chassis through systematic genome mapping and validation of neutral sites. *DNA Res.* **2015**, *22*, 425–437. [[CrossRef](#)]
253. Andersson, C.R.; Tsinoremas, N.F.; Shelton, J.; Lebedeva, N.V.; Yarrow, J.; Min, H.; Golden, S.S. Application of bioluminescence to the study of circadian rhythms in cyanobacteria. *Methods Enzymol.* **2000**, *305*. [[CrossRef](#)]
254. Niederholtmeyer, H.; Wolfstädter, B.T.; Savage, D.F.; Silver, P.A.; Way, J.C. Engineering cyanobacteria to synthesize and export hydrophilic products. *Appl. Environ. Microbiol.* **2010**, *76*, 3462–3466. [[CrossRef](#)]
255. Taton, A.; Unglaub, F.; Wright, N.E.; Zeng, W.Y.; Paz-Yepes, J.; Brahmasha, B.; Palenik, B.; Peterson, T.C.; Haerizadeh, F.; Golden, S.S.; et al. Broad-host-range vector system for synthetic biology and biotechnology in cyanobacteria. *Nucleic Acids Res.* **2014**, *42*. [[CrossRef](#)]
256. Xu, Y.; Alvey, R.M.; Byrne, P.O.; Graham, J.E.; Shen, G.; Bryant, D.A. Expression of genes in cyanobacteria: Adaptation of endogenous plasmids as platforms for high-level gene expression in *Synechococcus* sp. PCC 7002. *Methods Mol. Biol.* **2011**. [[CrossRef](#)]
257. Ruffing, A.M.; Jensen, T.J.; Strickland, L.M. Genetic tools for advancement of *Synechococcus* sp. PCC 7002 as a cyanobacterial chassis. *Microb. Cell Fact.* **2016**, *15*. [[CrossRef](#)]
258. Nozzi, N.E.; Case, A.E.; Carroll, A.L.; Atsumi, S. Systematic Approaches to Efficiently Produce 2,3-Butanediol in a Marine Cyanobacterium. *ACS Synth. Biol.* **2017**, *6*, 2136–2144. [[CrossRef](#)] [[PubMed](#)]
259. Mohamed, A.; Jansson, C. Influence of light on accumulation of photosynthesis-specific transcripts in the cyanobacterium *Synechocystis* 6803. *Plant Mol. Biol.* **1989**, *13*, 693–700. [[CrossRef](#)]
260. Mohamed, A.; Eriksson, J.; Osiewacz, H.D.; Jansson, C. Differential expression of the psbA genes in the cyanobacterium *Synechocystis* 6803. *MGG Mol. Gen. Genet.* **1993**, *238*, 161–168. [[CrossRef](#)]
261. Albers, S.C.; Gallegos, V.A.; Peebles, C.A.M. Engineering of genetic control tools in *Synechocystis* sp. PCC 6803 using rational design techniques. *J. Biotechnol.* **2015**, *216*, 36–46. [[CrossRef](#)]
262. Lin, P.-C.; Saha, R.; Zhang, F.; Pakrasi, H.B. Metabolic engineering of the pentose phosphate pathway for enhanced limonene production in the cyanobacterium *Synechocystis* sp. PCC 6803. *Sci. Rep.* **2017**, *7*. [[CrossRef](#)] [[PubMed](#)]
263. Jin, H.; Wang, Y.; Idoine, A.; Bhaya, D. Construction of a shuttle vector using an endogenous plasmid from the cyanobacterium *Synechocystis* sp. PCC6803. *Front. Microbiol.* **2018**, *9*. [[CrossRef](#)] [[PubMed](#)]
264. Liu, D.; Pakrasi, H.B. Exploring native genetic elements as plug-in tools for synthetic biology in the cyanobacterium *Synechocystis* sp. PCC 6803. *Microb. Cell Fact.* **2018**, *17*. [[CrossRef](#)]
265. Kamennaya, N.A.; Post, A.F. Characterization of cyanate metabolism in marine *Synechococcus* and *Prochlorococcus* spp. *Appl. Environ. Microbiol.* **2011**, *29*, 76–85. [[CrossRef](#)]
266. Kamennaya, N.A.; Ahn, S.E.; Park, H.; Bartal, R.; Sasaki, K.A.; Holman, H.Y.; Jansson, C. Installing extra bicarbonate transporters in the cyanobacterium *Synechocystis* sp. PCC6803 enhances biomass production. *Metab. Eng.* **2015**. [[CrossRef](#)] [[PubMed](#)]
267. Varman, A.M.; Xiao, Y.; Pakrasi, H.B.; Tang, Y.J. Metabolic engineering of *Synechocystis* sp. Strain PCC 6803 for isobutanol production. *Appl. Environ. Microbiol.* **2013**, *79*, 908–914. [[CrossRef](#)] [[PubMed](#)]
268. Lagarde, D.; Beuf, L.; Vermaas, W. Increased production of zeaxanthin and other pigments by application of genetic engineering techniques to *Synechocystis* sp. strain PCC 6803. *Appl. Environ. Microbiol.* **2000**, *66*, 64–72. [[CrossRef](#)] [[PubMed](#)]

269. Horiuchi, M.; Nakamura, K.; Kojima, K.; Nishiyama, Y.; Hatakeyama, W.; Hisabori, T.; Hihara, Y. The PedR transcriptional regulator interacts with thioredoxin to connect photosynthesis with gene expression in cyanobacteria. *Biochem. J.* **2010**, *431*, 135–140. [[CrossRef](#)]
270. Reinsvold, R.E.; Jinkerson, R.E.; Radakovits, R.; Posewitz, M.C.; Basu, C. The production of the sesquiterpene β -caryophyllene in a transgenic strain of the cyanobacterium *Synechocystis*. *J. Plant Physiol.* **2011**, *168*, 848–852. [[CrossRef](#)]
271. Nagarajan, A.; Winter, R.; Eaton-Rye, J.; Burnap, R. A synthetic DNA and fusion PCR approach to the ectopic expression of high levels of the D1 protein of photosystem II in *Synechocystis* sp. PCC 6803. *J. Photochem. Photobiol. B Biol.* **2011**, *104*, 212–219. [[CrossRef](#)]
272. Khetkorn, W.; Incharoensakdi, A.; Lindblad, P.; Jantaro, S. Enhancement of poly-3-hydroxybutyrate production in *Synechocystis* sp. PCC 6803 by overexpression of its native biosynthetic genes. *Bioresour. Technol.* **2016**, *214*, 761–768. [[CrossRef](#)]
273. Lindberg, P.; Park, S.; Melis, A. Engineering a platform for photosynthetic isoprene production in cyanobacteria, using *Synechocystis* as the model organism. *Metab. Eng.* **2010**, *12*, 70–79. [[CrossRef](#)] [[PubMed](#)]
274. Bentley, F.K.; Melis, A. Diffusion-based process for carbon dioxide uptake and isoprene emission in gaseous/aqueous two-phase photobioreactors by photosynthetic microorganisms. *Biotechnol. Bioeng.* **2012**, *109*. [[CrossRef](#)]
275. Bentley, F.K.; Zurbriggen, A.; Melis, A. Heterologous expression of the mevalonic acid pathway in cyanobacteria enhances endogenous carbon partitioning to isoprene. *Mol. Plant* **2014**, *7*, 71–86. [[CrossRef](#)]
276. Kudoh, K.; Hotta, S.; Sekine, M.; Fujii, R.; Uchida, A.; Kubota, G.; Kawano, Y.; Ihara, M. Overexpression of endogenous 1-deoxy-D-xylulose 5-phosphate synthase (DXS) in cyanobacterium *Synechocystis* sp. PCC6803 accelerates protein aggregation. *J. Biosci. Bioeng.* **2017**, *123*, 590–596. [[CrossRef](#)]
277. Englund, E.; Shabestary, K.; Hudson, E.P.; Lindberg, P. Systematic overexpression study to find target enzymes enhancing production of terpenes in *Synechocystis* PCC 6803, using isoprene as a model compound. *Metab. Eng.* **2018**, *49*, 164–177. [[CrossRef](#)] [[PubMed](#)]
278. Eungrasamee, K.; Miao, R.; Incharoensakdi, A.; Lindblad, P.; Jantaro, S. Improved lipid production via fatty acid biosynthesis and free fatty acid recycling in engineered *Synechocystis* sp. PCC 6803. *Biotechnol. Biofuels* **2019**, *12*. [[CrossRef](#)] [[PubMed](#)]
279. Brey, L.F.; Włodarczyk, A.J.; Bang Thøfner, J.F.; Burow, M.; Crocoll, C.; Nielsen, I.; Zygadlo Nielsen, A.J.; Jensen, P.E. Metabolic engineering of *Synechocystis* sp. PCC 6803 for the production of aromatic amino acids and derived phenylpropanoids. *Metab. Eng.* **2020**, *57*, 129–139. [[CrossRef](#)] [[PubMed](#)]
280. Deshpande, A.; Vue, J.; Morgan, J. Combining random mutagenesis and metabolic engineering for enhanced tryptophan production in *Synechocystis* sp. strain PCC 6803. *Appl. Environ. Microbiol.* **2020**, *86*. [[CrossRef](#)]
281. Bersanini, L.; Battchikova, N.; Jokel, M.; Rehman, A.; Vass, I.; Allahverdiyeva, Y.; Aro, E.M. Flavodiiron protein Flv2/Flv4-related photoprotective mechanism dissipates excitation pressure of PSII in cooperation with phycobilisomes in cyanobacteria. *Plant Physiol.* **2014**, *164*, 805–818. [[CrossRef](#)]
282. Mustila, H.; Paananen, P.; Battchikova, N.; Santana-Sánchez, A.; Muth-Pawlak, D.; Hagemann, M.; Aro, E.M.; Allahverdiyeva, Y. The flavodiiron protein Flv3 functions as a homo-oligomer during stress acclimation and is distinct from the Flv1/Flv3 hetero-oligomer specific to the O₂ photoreduction pathway. *Plant Cell Physiol.* **2016**, *57*, 1468–1483. [[CrossRef](#)] [[PubMed](#)]
283. Chaves, J.E.; Romero, P.R.; Kirst, H.; Melis, A. Role of isopentenyl-diphosphate isomerase in heterologous cyanobacterial (*Synechocystis*) isoprene production. *Photosynth. Res.* **2016**, *130*, 517–527. [[CrossRef](#)] [[PubMed](#)]
284. Chaves, J.E.; Rueda-Romero, P.; Kirst, H.; Melis, A. Engineering Isoprene Synthase Expression and Activity in Cyanobacteria. *ACS Synth. Biol.* **2017**, *6*, 2281–2292. [[CrossRef](#)]
285. Formighieri, C.; Melis, A. Heterologous synthesis of geranylinalool, a diterpenol plant product, in the cyanobacterium *Synechocystis*. *Appl. Microbiol. Biotechnol.* **2017**, *101*, 2791–2800. [[CrossRef](#)] [[PubMed](#)]
286. Sivaramakrishnan, R.; Incharoensakdi, A. Enhancement of lipid production in *Synechocystis* sp. PCC 6803 overexpressing glycerol kinase under oxidative stress with glycerol supplementation. *Bioresour. Technol.* **2018**, *267*, 532–540. [[CrossRef](#)] [[PubMed](#)]
287. Chaves, J.E.; Melis, A. Biotechnology of cyanobacterial isoprene production. *Appl. Microbiol. Biotechnol.* **2018**, *102*, 6452–6458. [[CrossRef](#)] [[PubMed](#)]
288. Formighieri, C.; Melis, A. Regulation of β -phellandrene synthase gene expression, recombinant protein accumulation, and monoterpene hydrocarbons production in *Synechocystis* transformants. *Planta* **2014**, *240*, 309–324. [[CrossRef](#)]
289. Formighieri, C.; Melis, A. A phycocyanin-phellandrene synthase fusion enhances recombinant protein expression and β -phellandrene (monoterpene) hydrocarbons production in *Synechocystis* (cyanobacteria). *Metab. Eng.* **2015**, *32*, 116–124. [[CrossRef](#)] [[PubMed](#)]
290. Formighieri, C.; Melis, A. Sustainable heterologous production of terpene hydrocarbons in cyanobacteria. *Photosynth. Res.* **2016**, *130*, 123–135. [[CrossRef](#)]
291. Betterle, N.; Melis, A. Heterologous Leader Sequences in Fusion Constructs Enhance Expression of Geranyl Diphosphate Synthase and Yield of β -Phellandrene Production in Cyanobacteria (*Synechocystis*). *ACS Synth. Biol.* **2018**, *7*, 912–921. [[CrossRef](#)]
292. Betterle, N.; Melis, A. Photosynthetic generation of heterologous terpenoids in cyanobacteria. *Biotechnol. Bioeng.* **2019**, *116*, 2041–2051. [[CrossRef](#)]
293. Valsami, E.A.; Psychogyiou, M.E.; Pateraki, A.; Chrysoulaki, E.; Melis, A.; Ghanotakis, D.F. Fusion constructs enhance heterologous β -phellandrene production in *Synechocystis* sp. PCC 6803. *J. Appl. Phycol.* **2020**. [[CrossRef](#)]

294. Marbouty, M.; Mazouni, K.; Saguez, C.; Cassier-Chauvat, C.; Chauvat, F. Characterization of the *Synechocystis* strain PCC 6803 penicillin-binding proteins and cytokinetic proteins FtsQ and FtsW and their network of interactions with ZipN. *J. Bacteriol.* **2009**, *191*, 5123–5133. [[CrossRef](#)] [[PubMed](#)]
295. Xue, Y.; Zhang, Y.; Cheng, D.; Daddy, S.; He, Q. Genetically engineering *Synechocystis* sp. Pasteur Culture Collection 6803 for the sustainable production of the plant secondary metabolite p-coumaric acid. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 9449–9454. [[CrossRef](#)]
296. Aoki, S.; Kondo, T.; Ishiura, M. Circadian expression of the dnaK gene in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **1995**, *177*, 5606–5611. [[CrossRef](#)] [[PubMed](#)]
297. Peca, L.; Kós, P.B.; Máté, Z.; Farsang, A.; Vass, I. Construction of bioluminescent cyanobacterial reporter strains for detection of nickel, cobalt and zinc. *FEMS Microbiol. Lett.* **2008**, *1837*, 1293–1304. [[CrossRef](#)] [[PubMed](#)]
298. Muramatsu, M.; Hihara, Y. Transcriptional regulation of genes encoding subunits of photosystem I during acclimation to high-light conditions in *Synechocystis* sp. PCC 6803. *Planta* **2003**, *216*, 446–453. [[CrossRef](#)]
299. Takahashi, T.; Nakai, N.; Muramatsu, M.; Hihara, Y. Role of multiple HLR1 sequences in the regulation of the dual promoters of the psaAB genes in *Synechocystis* sp. PCC 6803. *J. Bacteriol.* **2010**, *192*, 4031–4036. [[CrossRef](#)]
300. Rutherford, J.C.; Cavet, J.S.; Robinson, N.J. Cobalt-dependent transcriptional switching by a dual-effector MerR-like protein regulates a cobalt-exporting variant CPx-type ATPase. *J. Biol. Chem.* **1999**, *274*, 25827–25832. [[CrossRef](#)]
301. Calderon, R.H.; García-Cerdán, J.G.; Malnoë, A.; Cook, R.; Russell, J.J.; Gaw, C.; Dent, R.M.; De Vitry, C.; Niyogi, K.K. A conserved rubredoxin is necessary for photosystem II accumulation in diverse oxygenic photoautotrophs. *J. Biol. Chem.* **2013**, *288*, 26688–26696. [[CrossRef](#)]
302. Ke, W.T.; Dai, G.Z.; Jiang, H.B.; Zhang, R.; Qiu, B.S. Essential roles of iron superoxide dismutase in photoautotrophic growth of *Synechocystis* sp. PCC 6803 and heterogeneous expression of marine *Synechococcus* sp. CC9311 copper/zinc superoxide dismutase within its sodB knockdown mutant. *Microbiology (UK)* **2014**, *160*, 228–241. [[CrossRef](#)] [[PubMed](#)]
303. Kunert, A.; Hagemann, M.; Erdmann, N. Construction of promoter probe vectors for *Synechocystis* sp. PCC 6803 using the light-emitting reporter systems Gfp and LuxAB. *J. Microbiol. Methods* **2000**, *41*, 185–194. [[CrossRef](#)]
304. Tan, X.; Yao, L.; Gao, Q.; Wang, W.; Qi, F.; Lu, X. Photosynthesis driven conversion of carbon dioxide to fatty alcohols and hydrocarbons in cyanobacteria. *Metab. Eng.* **2011**, *13*, 169–176. [[CrossRef](#)]
305. Ungerer, J.; Tao, L.; Davis, M.; Ghirardi, M.; Maness, P.C.; Yu, J. Sustained photosynthetic conversion of CO₂ to ethylene in recombinant cyanobacterium *Synechocystis* 6803. *Energy Environ. Sci.* **2012**, *5*, 8998–9006. [[CrossRef](#)]
306. Veetil, V.P.; Angermayr, S.A.; Hellingwerf, K.J. Ethylene production with engineered *Synechocystis* sp PCC 6803 strains. *Microb. Cell Fact.* **2017**, *16*. [[CrossRef](#)] [[PubMed](#)]
307. Angermayr, S.A.; Paszota, M.; Hellingwerf, K.J. Engineering a cyanobacterial cell factory for production of lactic acid. *Appl. Environ. Microbiol.* **2012**, *78*, 7098–7106. [[CrossRef](#)]
308. Angermayr, S.A.; Van Der Woude, A.D.; Correddu, D.; Vreugdenhil, A.; Verrone, V.; Hellingwerf, K.J. Exploring metabolic engineering design principles for the photosynthetic production of lactic acid by *Synechocystis* sp. PCC6803. *Biotechnol. Biofuels* **2014**, *7*. [[CrossRef](#)]
309. Van der Woude, A.D.; Angermayr, S.A.; Puthan Veetil, V.; Osnato, A.; Hellingwerf, K.J. Carbon sink removal: Increased photosynthetic production of lactic acid by *Synechocystis* sp. PCC6803 in a glycogen storage mutant. *J. Biotechnol.* **2014**, *184*, 100–102. [[CrossRef](#)] [[PubMed](#)]
310. Hidese, R.; Matsuda, M.; Osanai, T.; Hasunuma, T.; Kondo, A. Malic Enzyme Facilitates d-Lactate Production through Increased Pyruvate Supply during Anoxic Dark Fermentation in *Synechocystis* sp. PCC 6803. *ACS Synth. Biol.* **2020**, *9*, 260–268. [[CrossRef](#)] [[PubMed](#)]
311. Savakis, P.E.; Angermayr, S.A.; Hellingwerf, K.J. Synthesis of 2,3-butanediol by *Synechocystis* sp. PCC6803 via heterologous expression of a catabolic pathway from lactic acid- and enterobacteria. *Metab. Eng.* **2013**, *20*, 121–130. [[CrossRef](#)] [[PubMed](#)]
312. Du, W.; Liang, F.; Duan, Y.; Tan, X.; Lu, X. Exploring the photosynthetic production capacity of sucrose by cyanobacteria. *Metab. Eng.* **2013**, *19*, 17–25. [[CrossRef](#)] [[PubMed](#)]
313. Viola, S.; Rühle, T.; Leister, D. A single vector-based strategy for marker-less gene replacement in *Synechocystis* sp. PCC 6803. *Microb. Cell Fact.* **2014**, *13*. [[CrossRef](#)] [[PubMed](#)]
314. Yao, L.; Qi, F.; Tan, X.; Lu, X. Improved production of fatty alcohols in cyanobacteria by metabolic engineering. *Biotechnol. Biofuels* **2014**, *7*. [[CrossRef](#)] [[PubMed](#)]
315. Yoshikawa, K.; Hirasawa, T.; Shimizu, H. Effect of malic enzyme on ethanol production by *Synechocystis* sp. PCC 6803. *J. Biosci. Bioeng.* **2015**, *119*, 82–84. [[CrossRef](#)]
316. Namakoshi, K.; Nakajima, T.; Yoshikawa, K.; Toya, Y.; Shimizu, H. Combinatorial deletions of glgC and phaCE enhance ethanol production in *Synechocystis* sp. PCC 6803. *J. Biotechnol.* **2016**, *239*, 13–19. [[CrossRef](#)]
317. Bartasun, P.; Prandi, N.; Storch, M.; Akin, Y.; Bennett, M.; Palma, A.; Baldwin, G.; Sakuragi, Y.; Jones, P.R.; Rowland, J. The effect of modulating the quantity of enzymes in a model ethanol pathway on metabolic flux in *Synechocystis* sp. PCC 6803. *PeerJ* **2019**, *7*. [[CrossRef](#)] [[PubMed](#)]
318. Savakis, P.; Tan, X.; Du, W.; Branco Dos Santos, F.; Lu, X.; Hellingwerf, K.J. Photosynthetic production of glycerol by a recombinant cyanobacterium. *J. Biotechnol.* **2015**, *195*, 46–51. [[CrossRef](#)]

319. Anfelt, J.; Kaczmarzyk, D.; Shabestary, K.; Renberg, B.; Rockberg, J.; Nielsen, J.; Uhlén, M.; Hudson, E.P. Genetic and nutrient modulation of acetyl-CoA levels in *Synechocystis* for n-butanol production. *Microb. Cell Fact.* **2015**, *14*. [[CrossRef](#)]
320. Hu, J.; Li, T.; Xu, W.; Zhan, J.; Chen, H.; He, C.; Wang, Q. Small antisense RNA RbIR positively regulates RuBisCo in *Synechocystis* sp. PCC 6803. *Front. Microbiol.* **2017**. [[CrossRef](#)]
321. Wang, B.; Eckert, C.; Maness, P.C.; Yu, J. A Genetic Toolbox for Modulating the Expression of Heterologous Genes in the Cyanobacterium *Synechocystis* sp. PCC 6803. *ACS Synth. Biol.* **2018**, *7*, 276–286. [[CrossRef](#)]
322. David, C.; Schmid, A.; Adrian, L.; Wilde, A.; Bühler, K. Production of 1,2-propanediol in photoautotrophic *Synechocystis* is linked to glycogen turn-over. *Biotechnol. Bioeng.* **2018**, *115*, 300–311. [[CrossRef](#)] [[PubMed](#)]
323. Qiao, Y.; Wang, W.; Lu, X. High Light Induced Alka(e)ne Biodegradation for Lipid and Redox Homeostasis in Cyanobacteria. *Front. Microbiol.* **2020**, *11*. [[CrossRef](#)] [[PubMed](#)]
324. Wu, W.; Du, W.; Gallego, R.P.; Hellingwerf, K.J.; Van Der Woude, A.D.; Dos Santos, F.B. Using osmotic stress to stabilize mannitol production in *Synechocystis* sp. PCC6803. *Biotechnol. Biofuels* **2020**, *13*. [[CrossRef](#)] [[PubMed](#)]
325. Sebesta, J.; Peebles, C.A. Improving heterologous protein expression in *Synechocystis* sp. PCC 6803 for alpha-bisabolene production. *Metab. Eng. Commun.* **2020**, *10*. [[CrossRef](#)] [[PubMed](#)]
326. Gao, Z.; Zhao, H.; Li, Z.; Tan, X.; Lu, X. Photosynthetic production of ethanol from carbon dioxide in genetically engineered cyanobacteria. *Energy Environ. Sci.* **2012**, *5*, 9857–9865. [[CrossRef](#)]
327. Wang, W.; Liu, X.; Lu, X. Engineering cyanobacteria to improve photosynthetic production of alka(e)nes. *Biotechnol. Biofuels* **2013**, *6*. [[CrossRef](#)]
328. Immethun, C.M.; DeLorenzo, D.M.; Focht, C.M.; Gupta, D.; Johnson, C.B.; Moon, T.S. Physical, chemical, and metabolic state sensors expand the synthetic biology toolbox for *Synechocystis* sp. PCC 6803. *Biotechnol. Bioeng.* **2017**, *114*, 1561–1569. [[CrossRef](#)]
329. Gao, Q.; Wang, W.; Zhao, H.; Lu, X. Effects of fatty acid activation on photosynthetic production of fatty acid-based biofuels in *Synechocystis* sp. PCC6803. *Biotechnol. Biofuels* **2012**, *5*. [[CrossRef](#)] [[PubMed](#)]
330. Lee, T.C.; Xiong, W.; Paddock, T.; Carrieri, D.; Chang, I.F.; Chiu, H.F.; Ungerer, J.; Hank Juo, S.H.; Maness, P.C.; Yu, J. Engineered xylose utilization enhances bio-products productivity in the cyanobacterium *Synechocystis* sp. PCC 6803. *Metab. Eng.* **2015**, *30*, 179–189. [[CrossRef](#)]
331. Hein, S.; Tran, H.; Steinbüchel, A. *Synechocystis* sp. PCC6803 possesses a two-component polyhydroxyalkanoic acid synthase similar to that of anoxygenic purple sulfur bacteria. *Arch. Microbiol.* **1998**, *170*, 162–170. [[CrossRef](#)]
332. Zhou, J.; Zhang, H.; Zhang, Y.; Li, Y.; Ma, Y. Designing and creating a modularized synthetic pathway in cyanobacterium *Synechocystis* enables production of acetone from carbon dioxide. *Metab. Eng.* **2012**, *14*, 394–400. [[CrossRef](#)]
333. Cameron, J.C.; Pakrasi, H.B. Essential role of glutathione in acclimation to environmental and redox perturbations in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.* **2010**, *154*, 1672–1685. [[CrossRef](#)]
334. Aoki, R.; Goto, T.; Fujita, Y. A heme oxygenase isoform is essential for aerobic growth in the cyanobacterium *Synechocystis* sp. PCC 6803: Modes of differential operation of two isoforms/enzymes to adapt to low oxygen environments in cyanobacteria. *Plant Cell Physiol.* **2011**, *52*, 1744–1756. [[CrossRef](#)] [[PubMed](#)]
335. Osanai, T.; Oikawa, A.; Numata, K.; Kuwahara, A.; Iijima, H.; Doi, Y.; Saito, K.; Hirai, M.Y. Pathway-level acceleration of glycogen catabolism by a response regulator in the cyanobacterium *Synechocystis* species PCC 6803. *Plant Physiol.* **2014**, *164*, 1831–1841. [[CrossRef](#)]
336. Hasunuma, T.; Matsuda, M.; Senga, Y.; Aikawa, S.; Toyoshima, M.; Shimakawa, G.; Miyake, C.; Kondo, A. Overexpression of *flv3* improves photosynthesis in the cyanobacterium *Synechocystis* sp. PCC6803 by enhancement of alternative electron flow. *Biotechnol. Biofuels* **2014**, *7*. [[CrossRef](#)]
337. Tashiro, M.; Kiyota, H.; Kawai-Noma, S.; Saito, K.; Ikeuchi, M.; Iijima, Y.; Umeno, D. Bacterial Production of Pinene by a Laboratory-Evolved Pinene-Synthase. *ACS Synth. Biol.* **2016**, *5*. [[CrossRef](#)]
338. Englund, E.; Liang, F.; Lindberg, P. Evaluation of promoters and ribosome binding sites for biotechnological applications in the unicellular cyanobacterium *Synechocystis* sp. PCC 6803. *Sci. Rep.* **2016**, *6*. [[CrossRef](#)]
339. Behler, J.; Vijay, D.; Hess, W.R.; Akhtar, M.K. CRISPR-Based Technologies for Metabolic Engineering in Cyanobacteria. *Trends Biotechnol.* **2018**, *36*, 996–1010. [[CrossRef](#)] [[PubMed](#)]
340. Durall, C.; Lindberg, P.; Yu, J.; Lindblad, P. Increased ethylene production by overexpressing phosphoenolpyruvate carboxylase in the cyanobacterium *Synechocystis* PCC 6803. *Biotechnol. Biofuels* **2020**, *13*. [[CrossRef](#)] [[PubMed](#)]
341. Matsudaira, A.; Hoshino, Y.; Uesaka, K.; Takatani, N.; Omata, T.; Usuda, Y. Production of glutamate and stereospecific flavors, (S)-linalool and (+)-valencene, by *Synechocystis* sp. PCC6803. *J. Biosci. Bioeng.* **2020**, *130*, 464–470. [[CrossRef](#)]
342. Kiyota, H.; Okuda, Y.; Ito, M.; Hirai, M.Y.; Ikeuchi, M. Engineering of cyanobacteria for the photosynthetic production of limonene from CO₂. *J. Biotechnol.* **2014**, *185*, 1–7. [[CrossRef](#)] [[PubMed](#)]
343. Wang, Y.; Sun, T.; Gao, X.; Shi, M.; Wu, L.; Chen, L.; Zhang, W. Biosynthesis of platform chemical 3-hydroxypropionic acid (3-HP) directly from CO₂ in cyanobacterium *Synechocystis* sp. PCC 6803. *Metab. Eng.* **2016**, *34*, 60–70. [[CrossRef](#)]
344. Englund, E.; Andersen-Ranberg, J.; Miao, R.; Hamberger, B.; Lindberg, P. Metabolic Engineering of *Synechocystis* sp. PCC 6803 for Production of the Plant Diterpenoid Manoyl Oxide. *ACS Synth. Biol.* **2015**, *4*, 1270–1278. [[CrossRef](#)]
345. Shabestary, K.; Anfelt, J.; Ljungqvist, E.; Jahn, M.; Yao, L.; Hudson, E.P. Targeted Repression of Essential Genes to Arrest Growth and Increase Carbon Partitioning and Biofuel Titrers in Cyanobacteria. *ACS Synth. Biol.* **2018**, *7*, 1669–1675. [[CrossRef](#)] [[PubMed](#)]

346. Wang, B.; Pugh, S.; Nielsen, D.R.; Zhang, W.; Meldrum, D.R. Engineering cyanobacteria for photosynthetic production of 3-hydroxybutyrate directly from CO₂. *Metab. Eng.* **2013**, *16*, 68–77. [[CrossRef](#)]
347. Li, H.; Liao, J.C. Engineering a cyanobacterium as the catalyst for the photosynthetic conversion of CO₂ to 1,2-propanediol. *Microb. Cell Fact.* **2013**, *12*. [[CrossRef](#)]
348. Wang, Y.; Tao, F.; Ni, J.; Li, C.; Xu, P. Production of C₃ platform chemicals from CO₂ by genetically engineered cyanobacteria. *Green Chem.* **2015**, *17*. [[CrossRef](#)]
349. Li, C.; Tao, F.; Ni, J.; Wang, Y.; Yao, F.; Xu, P. Enhancing the light-driven production of d-lactate by engineering cyanobacterium using a combinational strategy. *Sci. Rep.* **2015**, *5*, 9777. [[CrossRef](#)]
350. Lan, E.I.; Wei, C.T. Metabolic engineering of cyanobacteria for the photosynthetic production of succinate. *Metab. Eng.* **2016**, *38*, 483–493. [[CrossRef](#)]
351. Carbonell, V.; Vuorio, E.; Aro, E.M.; Kallio, P. Enhanced stable production of ethylene in photosynthetic cyanobacterium *Synechococcus elongatus* PCC 7942. *World J. Microbiol. Biotechnol.* **2019**, *35*, 77. [[CrossRef](#)]
352. Velmurugan, R.; Incharoensakdi, A. Heterologous Expression of Ethanol Synthesis Pathway in Glycogen Deficient *Synechococcus elongatus* PCC 7942 Resulted in Enhanced Production of Ethanol and Exopolysaccharides. *Front Plant Sci* **2020**, *11*. [[CrossRef](#)] [[PubMed](#)]
353. Ruffing, A.M.; Jones, H.D.T. Physiological effects of free fatty acid production in genetically engineered *Synechococcus elongatus* PCC 7942. *Biotechnol. Bioeng.* **2012**, *109*, 2190–2199. [[CrossRef](#)]
354. Ruffing, A.M. Borrowing genes from *Chlamydomonas reinhardtii* for free fatty acid production in engineered cyanobacteria. *J. Appl. Phycol.* **2013**, *25*, 1495–1507. [[CrossRef](#)]
355. Ruffing, A.M. Improved free fatty acid production in cyanobacteria with *Synechococcus* sp. PCC 7002 as host. *Front. Bioeng. Biotechnol.* **2014**, *2*, 1–10. [[CrossRef](#)] [[PubMed](#)]
356. Kachel, B.; Mack, M. Engineering of *Synechococcus* sp. strain PCC 7002 for the photoautotrophic production of light-sensitive riboflavin (vitamin B2). *Metab. Eng.* **2020**, *62*, 275–286. [[CrossRef](#)]
357. Jazmin, L.J.; Xu, Y.; Cheah, Y.E.; Adebiyi, A.O.; Johnson, C.H.; Young, J.D. Isotopically nonstationary ¹³C flux analysis of cyanobacterial isobutyraldehyde production. *Metab. Eng.* **2017**, *42*, 9–18. [[CrossRef](#)]
358. Sommer, M.; Sutter, M.; Gupta, S.; Kirst, H.; Turmo, A.; Lechno-Yossef, S.; Burton, R.L.; Saechao, C.; Sloan, N.B.; Cheng, X.; et al. Heterohexamers formed by CcmK3 and CcmK4 increase the complexity of beta carboxysome shells. *Plant Physiol.* **2019**, *179*, 156–167. [[CrossRef](#)]
359. Clerico, E.M.; Cassone, V.M.; Golden, S.S. Stability and lability of circadian period of gene expression in the cyanobacterium *Synechococcus elongatus*. *Microbiology* **2009**, *155*, 635–641. [[CrossRef](#)]
360. Atsumi, S.; Higashide, W.; Liao, J.C. Direct photosynthetic recycling of carbon dioxide to isobutyraldehyde. *Nat. Biotechnol.* **2009**, *27*, 1177–1180. [[CrossRef](#)] [[PubMed](#)]
361. Lan, E.I.; Liao, J.C. Metabolic engineering of cyanobacteria for 1-butanol production from carbon dioxide. *Metab. Eng.* **2011**, *13*, 353–363. [[CrossRef](#)]
362. Lan, E.I.; Liao, J.C. ATP drives direct photosynthetic production of 1-butanol in cyanobacteria. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 6018–6023. [[CrossRef](#)] [[PubMed](#)]
363. Cameron, J.C.; Wilson, S.C.; Bernstein, S.L.; Kerfeld, C.A. Biogenesis of a bacterial organelle: The carboxysome assembly pathway. *Cell* **2013**, *155*, 1131–1140. [[CrossRef](#)] [[PubMed](#)]
364. Chen, A.H.; Robinson-Mosher, A.; Savage, D.F.; Silver, P.A.; Polka, J.K. The Bacterial Carbon-Fixing Organelle Is Formed by Shell Envelopment of Preassembled Cargo. *PLoS ONE* **2013**, *8*, e76127. [[CrossRef](#)] [[PubMed](#)]
365. Kusakabe, T.; Tatsuke, T.; Tsuruno, K.; Hirokawa, Y.; Atsumi, S.; Liao, J.C.; Hanai, T. Engineering a synthetic pathway in cyanobacteria for isopropanol production directly from carbon dioxide and light. *Metab. Eng.* **2013**, *20*, 101–108. [[CrossRef](#)]
366. Li, X.; Shen, C.R.; Liao, J.C. Isobutanol production as an alternative metabolic sink to rescue the growth deficiency of the glycogen mutant of *Synechococcus elongatus* PCC 7942. *Photosynth. Res.* **2014**, *120*, 301–310. [[CrossRef](#)]
367. Lan, E.I.; Chuang, D.S.; Shen, C.R.; Lee, A.M.; Ro, S.Y.; Liao, J.C. Metabolic engineering of cyanobacteria for photosynthetic 3-hydroxypropionic acid production from CO₂ using *Synechococcus elongatus* PCC 7942. *Metab. Eng.* **2015**, *31*, 163–170. [[CrossRef](#)] [[PubMed](#)]
368. Hirokawa, Y.; Maki, Y.; Tatsuke, T.; Hanai, T. Cyanobacterial production of 1,3-propanediol directly from carbon dioxide using a synthetic metabolic pathway. *Metab. Eng.* **2016**, *34*, 97–103. [[CrossRef](#)]
369. Wang, X.; Liu, W.; Xin, C.; Zheng, Y.; Cheng, Y.; Sun, S.; Li, R.; Zhu, X.-G.; Dai, S.Y.; Rentzepis, P.M.; et al. Enhanced limonene production in cyanobacteria reveals photosynthesis limitations. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 14225–14230. [[CrossRef](#)]
370. Chwa, J.W.; Kim, W.J.; Sim, S.J.; Um, Y.; Woo, H.M. Engineering of a modular and synthetic phosphoketolase pathway for photosynthetic production of acetone from CO₂ in *Synechococcus elongatus* PCC 7942 under light and aerobic condition. *Plant Biotechnol. J.* **2016**, *14*, 1768–1776. [[CrossRef](#)]
371. Gao, X.; Gao, F.; Liu, D.; Zhang, H.; Nie, X.; Yang, C. Engineering the methylerythritol phosphate pathway in cyanobacteria for photosynthetic isoprene production from CO₂. *Energy Environ. Sci.* **2016**, *7*, 71–86. [[CrossRef](#)]
372. Taton, A.; Ma, A.T.; Ota, M.; Golden, S.S.; Golden, J.W. NOT Gate Genetic Circuits to Control Gene Expression in Cyanobacteria. *ACS Synth. Biol.* **2017**, *6*, 2175. [[CrossRef](#)]

373. Kanno, M.; Atsumi, S. Engineering an Obligate photoautotrophic cyanobacterium to utilize glycerol for growth and chemical production. *ACS Synth. Biol.* **2017**, *6*, 69–75. [[CrossRef](#)] [[PubMed](#)]
374. Lee, H.J.; Lee, J.; Lee, S.-M.; Um, Y.; Kim, Y.; Sim, S.J.; Choi, J.-I.; Woo, H.M. Direct Conversion of CO₂ to α -Farnesene Using Metabolically Engineered *Synechococcus elongatus* PCC 7942. *J. Agric. Food Chem.* **2017**, *60*, 10424–10428. [[CrossRef](#)] [[PubMed](#)]
375. Pattharaprachayakul, N.; Lee, H.J.; Incharoensakdi, A.; Woo, H.M. Evolutionary Engineering of Cyanobacteria to Enhance the Production of α -Farnesene from CO₂. *J. Agric. Food Chem.* **2019**, *67*, 13658–13664. [[CrossRef](#)] [[PubMed](#)]
376. Hirokawa, Y.; Goto, R.; Umetani, Y.; Hanai, T. Construction of a novel D-lactate producing pathway from dihydroxyacetone phosphate of the Calvin cycle in cyanobacterium, *Synechococcus elongatus* PCC 7942. *J. Biosci. Bioeng.* **2017**, *124*, 54–61. [[CrossRef](#)] [[PubMed](#)]
377. Ducat, D.C.; Avelar-Rivas, J.A.; Way, J.C.; Silvera, P.A. Rerouting carbon flux to enhance photosynthetic productivity. *Appl. Environ. Microbiol.* **2012**, *78*, 2660–2668. [[CrossRef](#)] [[PubMed](#)]
378. Shih, P.M.; Zarzycki, J.; Niyogi, K.K.; Kerfeld, C.A. Introduction of a synthetic CO₂-fixing photorespiratory bypass into a cyanobacterium. *J. Biol. Chem.* **2014**, *289*, 9493–9500. [[CrossRef](#)]
379. Oliver, J.W.K.; Atsumi, S. A carbon sink pathway increases carbon productivity in cyanobacteria. *Metab. Eng.* **2015**, *29*, 106–112. [[CrossRef](#)]
380. Choi, S.Y.; Sim, S.J.; Choi, J.I.; Woo, H.M. Identification of small droplets of photosynthetic squalene in engineered *Synechococcus elongatus* PCC 7942 using TEM and selective fluorescent Nile red analysis. *Letts. Appl. Microbiol.* **2018**, *66*, 523–529. [[CrossRef](#)]
381. Kim, W.J.; Lee, S.M.; Um, Y.; Sim, S.J.; Woo, H.M. Development of synebrick vectors as a synthetic biology platform for gene expression in *Synechococcus elongatus* PCC 7942. *Front. Plant Sci.* **2017**, *8*. [[CrossRef](#)]
382. Roulet, J.; Taton, A.; Golden, J.W.; Arabolaza, A.; Burkart, M.D.; Gramajo, H. Development of a cyanobacterial heterologous polyketide production platform. *Metab. Eng.* **2018**, *49*, 94–104. [[CrossRef](#)] [[PubMed](#)]
383. Takahama, K.; Matsuoka, M.; Nagahama, K.; Ogawa, T. Construction and analysis of a recombinant cyanobacterium expressing a chromosomally inserted gene for an ethylene-forming enzyme at the psbAI locus. *J. Biosci. Bioeng.* **2003**, *15*, 302–305. [[CrossRef](#)]
384. Oliver, J.W.K.; Machado, I.M.P.; Yoneda, H.; Atsumi, S. Cyanobacterial conversion of carbon dioxide to 2,3-butanediol. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 1249–1254. [[CrossRef](#)]
385. Peterson, E.S.; McCue, L.A.; Schrimpe-Rutledge, A.C.; Jensen, J.L.; Walker, H.; Kobold, M.A.; Webb, S.R.; Payne, S.H.; Ansong, C.; Adkins, J.N.; et al. VESPA: Software to facilitate genomic annotation of prokaryotic organisms through integration of proteomic and transcriptomic data. *BMC Genom.* **2012**, *13*. [[CrossRef](#)]
386. Begemann, M.B.; Zess, E.K.; Walters, E.M.; Schmitt, E.F.; Markley, A.L.; Pflieger, B.F. An Organic Acid Based Counter Selection System for Cyanobacteria. *PLoS ONE* **2013**, *8*, e76594. [[CrossRef](#)]
387. Clark, R.L.; Gordon, G.C.; Bennett, N.R.; Lyu, H.; Root, T.W.; Pflieger, B.F. High-CO₂ Requirement as a Mechanism for the Containment of Genetically Modified Cyanobacteria. *ACS Synth. Biol.* **2018**, *7*, 384–391. [[CrossRef](#)] [[PubMed](#)]
388. Markley, A.L.; Begemann, M.B.; Clarke, R.E.; Gordon, G.C.; Pflieger, B.F. Synthetic Biology Toolbox for Controlling Gene Expression in the Cyanobacterium *Synechococcus* sp. strain PCC 7002. *ACS Synth. Biol.* **2015**, *4*, 595–603. [[CrossRef](#)] [[PubMed](#)]
389. Sakamoto, T.; Shen, G.; Higashi, S.; Murata, N.; Bryant, D.A. Alteration of low-temperature susceptibility of the cyanobacterium *Synechococcus* sp. PCC 7002 by genetic manipulation of membrane lipid unsaturation. *Arch. Microbiol.* **1997**, *169*, 20–28. [[CrossRef](#)] [[PubMed](#)]
390. Vogel, A.I.M.; Lale, R.; Hohmann-Marriott, M.F. Streamlining recombination-mediated genetic engineering by validating three neutral integration sites in *Synechococcus* sp. PCC 7002. *J. Biol. Eng.* **2017**, *11*. [[CrossRef](#)]
391. Selão, T.T.; Włodarczyk, A.; Nixon, P.J.; Norling, B. Growth and selection of the cyanobacterium *Synechococcus* sp. PCC 7002 using alternative nitrogen and phosphorus sources. *Metab. Eng.* **2019**, *54*, 2. [[CrossRef](#)]
392. Korosh, T.C.; Markley, A.L.; Clark, R.L.; McGinley, L.L.; McMahan, K.D.; Pflieger, B.F. Engineering photosynthetic production of L-lysine. *Metab. Eng.* **2017**, *44*, 273–283. [[CrossRef](#)]
393. Daniell, H.; Sarojini, G.; McFadden, B.A. Transformation of the cyanobacterium *Anacystis nidulans* 6301 with the *Escherichia coli* plasmid pBR322. *Proc. Natl. Acad. Sci. USA* **1986**, *83*, 2546–2550. [[CrossRef](#)]
394. Koksharova, O.; Wolk, C. Genetic tools for cyanobacteria. *Appl. Microbiol. Biotechnol.* **2002**, *58*, 123–137. [[CrossRef](#)]
395. Sherman, L.A.; Van de Putte, P. Construction of a hybrid plasmid capable of replication in the bacterium *Escherichia coli* and the cyanobacterium *Anacystis nidulans*. *J. Bacteriol.* **1982**, *150*, 410–413. [[CrossRef](#)]
396. Golden, S.S.; Sherman, L.A. Optimal conditions for genetic transformation of the cyanobacterium *Anacystis nidulans* R2. *J. Bacteriol.* **1984**, *158*, 36–42. [[CrossRef](#)]
397. Buzby, J.S.; Porter, R.D.; Stevens, S.E. Plasmid transformation in *Agmenellum quadruplicatum* PR-6: Construction of biphasic plasmids and characterization of their transformation properties. *J. Bacteriol.* **1983**, *154*, 1446–1450. [[CrossRef](#)] [[PubMed](#)]
398. Scharnagl, M.; Richter, S.; Hagemann, M. The cyanobacterium *Synechocystis* sp. strain PCC 6803 expresses a DNA methyltransferase specific for the recognition sequence of the restriction endonuclease PvuI. *J. Bacteriol.* **1998**, *180*, 4116–4122. [[CrossRef](#)] [[PubMed](#)]
399. Wang, B.; Yu, J.; Zhang, W.; Meldrum, D.R. Premethylation of Foreign DNA Improves Integrative Transformation. *Appl. Environ. Microbiol.* **2015**, *81*, 8500–8506. [[CrossRef](#)] [[PubMed](#)]

400. Hagemann, M.; Gartner, K.; Scharnagl, M.; Bolay, P.; Lott, S.C.; Fuss, J.; Huettel, B.; Reinhardt, R.; Klahn, S.; Hess, W.R. Identification of the DNA methyltransferases establishing the methylome of the cyanobacterium *Synechocystis* sp. PCC 6803. *DNA Res.* **2018**, *25*, 343–352. [[CrossRef](#)]
401. Wu, D.; Wang, Y.; Xu, X. Effects of a Type I RM System on Gene Expression and Glycogen Catabolism in *Synechocystis* sp. PCC 6803. *Front. Microbiol.* **2020**, *11*, 1258. [[CrossRef](#)]
402. Kuhlemeier, C.J.; Thomas, A.A.M.; van der Ende, A.; van Leen, R.W.; Borrias, W.E.; van den Hondel, C.A.M.J.J.; van Arkel, G.A. A host-vector system for gene cloning in the cyanobacterium *Anacystis nidulans* R2. *Plasmid* **1983**, *10*, 156–163. [[CrossRef](#)]
403. Golden, S.S.; Sherman, L.A. A hybrid plasmid is a stable cloning vector for the cyanobacterium *Anacystis nidulans* R2. *J. Bacteriol.* **1983**, *155*, 966–972. [[CrossRef](#)] [[PubMed](#)]
404. Gendel, S.; Straus, N.; Pulleyblank, D.; Williams, J. Shuttle cloning vectors for the cyanobacterium *Anacystis nidulans*. *J. Bacteriol.* **1983**, *156*, 148–154. [[CrossRef](#)] [[PubMed](#)]
405. Murphy, R.C.; Bryant, D.A.; Porter, R.D.; Tandeau De Marsac, N. Molecular cloning and characterization of the *recA* gene from the Cyanobacterium *Synechococcus* sp. strain PCC 7002. *J. Bacteriol.* **1987**, *172*, 967–976. [[CrossRef](#)] [[PubMed](#)]
406. Kuhlemeier, C.J.; Logtenberg, T.; Stoorvogel, W.; van Heugten, H.A.; Borrias, W.E.; van Arkel, G.A. Cloning of nitrate reductase genes from the cyanobacterium *Anacystis nidulans*. *J. Bacteriol.* **1984**, *159*, 36–41. [[CrossRef](#)] [[PubMed](#)]
407. Gruber, M.Y.; Glick, B.R.; Thompson, J.E. Cloned manganese superoxide dismutase reduces oxidative stress in *Escherichia coli* and *Anacystis nidulans*. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 2608–2612. [[CrossRef](#)] [[PubMed](#)]
408. Ferino, F.; Chauvat, F. A promoter-probe vector-host system for the cyanobacterium, *Synechocystis* PCC6803. *Gene* **1989**, *84*, 257–266. [[CrossRef](#)]
409. Marraccini, P.; Bulteau, S.; Cassier-Chauvat, C.; Mermet-Bouvier, P.; Chauvat, F. A conjugative plasmid vector for promoter analysis in several cyanobacteria of the genera *Synechococcus* and *Synechocystis*. *Plant Mol. Biol.* **1993**, *23*, 905–909. [[CrossRef](#)]
410. Chenebault, C.; Diaz-Santos, E.; Kammerscheit, X.; Görden, S.; Illoaia, C.; Streckaite, S.; Gall, A.; Robert, B.; Marcon, E.; Buisson, D.-A.; et al. A Genetic Toolbox for the New Model Cyanobacterium *Cyanothece* PCC 7425: A Case Study for the Photosynthetic Production of Limonene. *Front. Microbiol.* **2020**, *11*, 586–601. [[CrossRef](#)]
411. Murphy, R.C.; Stevens, S.E. Cloning and expression of the *cryIVD* gene of *Bacillus thuringiensis* subsp. *israelensis* in the cyanobacterium *Agmenellum quadruplicatum* PR-6 and its resulting larvicidal activity. *Appl. Environ. Microbiol.* **1992**, *58*, 1650–1655. [[CrossRef](#)] [[PubMed](#)]
412. Soltés-Rak, E.; Kushner, D.J.; Williams, D.D.; Coleman, J.R. Effect of promoter modification on mosquitocidal *cryIVB* gene expression in *Synechococcus* sp. strain PCC 7942. *Appl. Environ. Microbiol.* **1993**, *59*, 2404–2410. [[CrossRef](#)]
413. Erbe, J.L.; Taylor, K.B.; Hall, L.M. Expression of mouse metallothionein in the cyanobacterium *Synechococcus* PCC7942. *J. Ind. Microbiol.* **1996**, *17*, 41–46. [[CrossRef](#)]
414. Nomura, M.; Ishitani, M.; Takabe, T.; Rai, A.K.; Takabe, T. *Synechococcus* sp. PCC7942 transformed with *Escherichia coli* *bet* genes produces glycine betaine from choline and acquires resistance to salt stress. *Plant Physiol.* **1995**, *107*, 703–708. [[CrossRef](#)]
415. Kaku, N.; Hibino, T.; Tanaka, Y.; Ishikawa, H.; Araki, E.; Takabe, T.; Takabe, T. Effects of overexpression of *Escherichia coli* *katE* and *bet* genes on the tolerance for salt stress in a freshwater cyanobacterium *Synechococcus* sp. PCC 7942. *Plant Sci.* **2000**, *159*, 281–288. [[CrossRef](#)]
416. De Deng, M.; Coleman, J.R. Ethanol synthesis by genetic engineering in cyanobacteria. *Appl. Environ. Microbiol.* **1999**, *65*, 523–528. [[CrossRef](#)]
417. Fukuda, H.; Sakai, M.; Nagahama, K.; Fujii, T.; Matsuoka, M.; Inoue, Y.; Ogawa, T. Heterologous expression of the gene for the ethylene-forming enzyme from *Pseudomonas syringae* in the cyanobacterium *Synechococcus*. *Biotechnol. Lett.* **1994**, *16*, 1–6. [[CrossRef](#)]
418. Sakai, M.; Ogawa, T.; Matsuoka, M.; Fukuda, H. Photosynthetic conversion of carbon dioxide to ethylene by the recombinant cyanobacterium, *Synechococcus* sp. PCC 7942, which harbors a gene for the ethylene-forming enzyme of *Pseudomonas syringae*. *J. Ferment. Bioeng.* **1997**, *4*, 434–443. [[CrossRef](#)]
419. Jindou, S.; Ito, Y.; Mito, N.; Uematsu, K.; Hosoda, A.; Tamura, H. Engineered platform for bioethylene production by a cyanobacterium expressing a chimeric complex of plant enzymes. *ACS Synth. Biol.* **2014**, *3*, 487–496. [[CrossRef](#)] [[PubMed](#)]
420. Iwaki, T.; Haranoh, K.; Inoue, N.; Kojima, K.; Satoh, R.; Nishino, T.; Wada, S.; Ihara, H.; Tsuyama, S.; Kobayashi, H.; et al. Expression of foreign type I ribulose-1,5-bisphosphate carboxylase/ oxygenase (EC 4.1.1.39) stimulates photosynthesis in cyanobacterium *Synechococcus* PCC7942 cells. *Photosynth. Res.* **2006**, *88*, 287–297. [[CrossRef](#)] [[PubMed](#)]
421. Chen, Y.; Taton, A.; Go, M.; London, R.E.; Pieper, L.M.; Golden, S.S.; Golden, J.W. Self-replicating shuttle vectors based on pANS, a small endogenous plasmid of the unicellular cyanobacterium *Synechococcus elongatus* PCC 7942. *Microbiology (UK)* **2016**, *162*, 2029–2041. [[CrossRef](#)]
422. Scholz, P.; Haring, V.; Wittmann-Liebold, B.; Ashman, K.; Bagdasarian, M.; Scherzinger, E. Complete nucleotide sequence and gene organization of the broad-host-range plasmid RSF1010. *Gene* **1989**, *75*, 271–288. [[CrossRef](#)]
423. Bagdasarian, M.; Lurz, R.; Rückert, B.; Franklin, F.C.H.; Bagdasarian, M.M.; Frey, J.; Timmis, K.N. Specific-purpose plasmid cloning vectors II. Broad host range, high copy number, RSF 1010-derived vectors, and a host-vector system for gene cloning in *Pseudomonas*. *Gene* **1981**, *16*, 237–247. [[CrossRef](#)]
424. Kues, U.; Stahl, U. Replication of plasmids in gram-negative bacteria. *Microbiol. Rev.* **1989**, *53*, 491–516. [[CrossRef](#)] [[PubMed](#)]

425. Meyer, R. Replication and conjugative mobilization of broad host-range IncQ plasmids. *Plasmid* **2009**, *62*, 57–70. [[CrossRef](#)] [[PubMed](#)]
426. Kreps, S.; Ferino, F.; Mosrin, C.; Gerits, J.; Mergeay, M.; Thuriaux, P. Conjugative transfer and autonomous replication of a promiscuous IncQ plasmid in the cyanobacterium *Synechocystis* PCC 6803. *MGG Mol. Gen. Genet.* **1990**, *221*, 129–133. [[CrossRef](#)]
427. Sode, K.; Tatara, M.; Takeyama, H.; Burgess, J.G.; Matsunaga, T. Conjugative gene transfer in marine cyanobacteria: *Synechococcus* sp., *Synechocystis* sp. and *Pseudanabaena* sp. *Appl. Microbiol. Biotechnol.* **1992**, *37*, 369–373. [[CrossRef](#)] [[PubMed](#)]
428. Mermet-Bouvier, P.; Cassier-Chauvat, C.; Marraccini, P.; Chauvat, F. Transfer and replication of RSF1010-derived plasmids in several cyanobacteria of the genera *Synechocystis* and *Synechococcus*. *Curr. Microbiol.* **1993**, *27*, 323–327. [[CrossRef](#)]
429. Mühlhoff, U.; Chauvat, F. Gene transfer and manipulation in the thermophilic cyanobacterium *Synechococcus elongatus*. *Mol. Gen. Genet.* **1996**, *252*, 93–100. [[CrossRef](#)] [[PubMed](#)]
430. Brahamsha, B. A genetic manipulation system for oceanic cyanobacteria of the genus *Synechococcus*. *Appl. Environ. Microbiol.* **1996**, *62*, 1745–1751. [[CrossRef](#)]
431. Guo, H.; Xu, X. Broad host range plasmid-based gene transfer system in the cyanobacterium *Gloeobacter violaceus* which lacks thylakoids. *Prog. Nat. Sci.* **2004**, *14*, 31–35. [[CrossRef](#)]
432. Araki, M.; Shimada, Y.; Mimuro, M.; Tsuchiya, T. Establishment of the reporter system for a thylakoid-lacking cyanobacterium, *Gloeobacter violaceus* PCC 7421. *FEBS Open Bio* **2013**, *3*, 11–15. [[CrossRef](#)]
433. Tolonen, A.C.; Liszt, G.B.; Hess, W.R. Genetic manipulation of *Prochlorococcus* strain MIT9313: Green fluorescent protein expression from an RSF1010 plasmid and Tn5 transposition. *Appl. Environ. Microbiol.* **2006**, *72*, 7607–7613. [[CrossRef](#)]
434. Huang, H.H.; Camsund, D.; Lindblad, P.; Heidorn, T. Design and characterization of molecular tools for a synthetic biology approach towards developing cyanobacterial biotechnology. *Nucleic Acids Res.* **2010**, *38*, 2577–2593. [[CrossRef](#)] [[PubMed](#)]
435. Bishé, B.; Taton, A.; Golden, J.W. Modification of RSF1010-Based Broad-Host-Range Plasmids for Improved Conjugation and Cyanobacterial Bioprospecting. *iScience* **2019**, *20*, 216–228. [[CrossRef](#)] [[PubMed](#)]
436. Taton, A.; Lis, E.; Adin, D.M.; Dong, G.; Cookson, S.; Kay, S.A.; Golden, S.S.; Golden, J.W. Gene transfer in *Leptolyngbya* sp. strain BL0902, a cyanobacterium suitable for production of biomass and bioproducts. *PLoS ONE* **2012**, *7*, e30901. [[CrossRef](#)]
437. Mermet-Bouvier, P.; Chauvat, F. A conditional expression vector for the cyanobacteria *Synechocystis* sp. strains PCC6803 and PCC6714 or *Synechococcus* sp. strains PCC7942 and PCC6301. *Curr. Microbiol.* **1994**, *28*, 145–148. [[CrossRef](#)] [[PubMed](#)]
438. Mazouni, K.; Bulteau, S.; Cassier-Chauvat, C.; Chauvat, F. Promoter element spacing controls basal expression and light inducibility of the cyanobacterial *secA* gene. *Mol. Microbiol.* **1998**, *30*, 1113–1122. [[CrossRef](#)]
439. Duthel, J.; Saenkham, P.; Sakr, S.; Leplat, C.; Ortega-Ramos, M.; Bottin, H.; Cournac, L.; Cassier-Chauvat, C.; Chauvat, F. The AbrB2 autorepressor, expressed from an atypical promoter, represses the hydrogenase operon to regulate hydrogen production in *Synechocystis* strain PCC6803. *J. Bacteriol.* **2012**, *194*, 5423–5433. [[CrossRef](#)]
440. Mazouni, K.; Domain, F.; Cassier-Chauvat, C.; Chauvat, F. Molecular analysis of the key cytokinetic components of cyanobacteria: FtsZ, ZipN and MinCDE. *Mol. Microbiol.* **2004**, *52*, 1145–1158. [[CrossRef](#)]
441. Marbouty, M.; Saguez, C.; Cassier-Chauvat, C.; Chauvat, F. Characterization of the FtsZ-interacting septal proteins SepF and Ftn6 in the spherical-celled cyanobacterium *Synechocystis* strain PCC 6803. *J. Bacteriol.* **2009**, *191*, 6178–6185. [[CrossRef](#)]
442. Marbouty, M.; Saguez, C.; Cassier-Chauvat, C.; Chauvat, F. ZipN, an FtsA-like orchestrator of divisome assembly in the model cyanobacterium *Synechocystis* PCC6803. *Mol. Microbiol.* **2009**, *74*, 409–420. [[CrossRef](#)]
443. Damrow, R.; Maldener, I.; Zilliges, Y. The multiple functions of common microbial carbon polymers, glycogen and PHB, during stress responses in the non-diazotrophic cyanobacterium *Synechocystis* sp. PCC 6803. *Front. Microbiol.* **2016**, *7*. [[CrossRef](#)]
444. Ng, W.O.; Zentella, R.; Wang, Y.; Taylor, J.S.A.; Pakrasi, H.B. *phrA*, the major photoreactivating factor in the cyanobacterium *Synechocystis* sp. strain PCC 6803 codes for a cyclobutane-pyrimidine-dimer-specific DNA photolyase. *Arch. Microbiol.* **2000**, *173*, 412–416. [[CrossRef](#)]
445. De Porcellinis, A.J.; Nørgaard, H.; Brey, L.M.F.; Erstad, S.M.; Jones, P.R.; Heazlewood, J.L.; Sakuragi, Y. Overexpression of bifunctional fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase leads to enhanced photosynthesis and global reprogramming of carbon metabolism in *Synechococcus* sp. PCC 7002. *Metab. Eng.* **2018**, *47*, 170–183. [[CrossRef](#)] [[PubMed](#)]
446. Dienst, D.; Wichmann, J.; Mantovani, O.; Rodrigues, J.S.; Lindberg, P. High density cultivation for efficient sesquiterpenoid biosynthesis in *Synechocystis* sp. PCC 6803. *Sci. Rep.* **2020**, *10*, 5932. [[CrossRef](#)]
447. Liang, F.; Englund, E.; Lindberg, P.; Lindblad, P. Engineered cyanobacteria with enhanced growth show increased ethanol production and higher biofuel to biomass ratio. *Metab. Eng.* **2018**, *46*, 51–59. [[CrossRef](#)]
448. Guerrero, F.; Carbonell, V.; Cossu, M.; Correddu, D.; Jones, P.R. Ethylene Synthesis and Regulated Expression of Recombinant Protein in *Synechocystis* sp. PCC 6803. *PLoS ONE* **2012**, *7*, e50470. [[CrossRef](#)] [[PubMed](#)]
449. Kuchmina, E.; Klähn, S.; Jakob, A.; Bigott, W.; Enke, H.; Dühring, U.; Wilde, A. Ethylene production in *Synechocystis* sp. PCC 6803 promotes phototactic movement. *Microbiology (UK)* **2017**, *16*, 1937–1947. [[CrossRef](#)] [[PubMed](#)]
450. Thiel, K.; Mulaku, E.; Dandapani, H.; Nagy, C.; Aro, E.M.; Kallio, P. Translation efficiency of heterologous proteins is significantly affected by the genetic context of RBS sequences in engineered cyanobacterium *Synechocystis* sp. PCC 6803. *Microb. Cell Fact.* **2018**, *17*, 34. [[CrossRef](#)] [[PubMed](#)]
451. Van der Woude, A.D.; Perez Gallego, R.; Vreugdenhil, A.; Puthan Veetil, V.; Chroumpi, T.; Hellingwerf, K.J. Genetic engineering of *Synechocystis* PCC6803 for the photoautotrophic production of the sweetener erythritol. *Microb. Cell Fact.* **2016**, *15*. [[CrossRef](#)]

452. Ortega-Ramos, M.; Jittawuttipoka, T.; Saenkham, P.; Czarnecka-Kwasiborski, A.; Bottin, H.; Cassier-Chauvat, C.; Chauvat, F. Engineering *Synechocystis* PCC6803 for hydrogen production: Influence on the tolerance to oxidative and sugar stresses. *PLoS ONE* **2014**, *9*, e89372. [[CrossRef](#)]
453. Veaudor, T.; Cassier-Chauvat, C.; Chauvat, F. Overproduction of the cyanobacterial hydrogenase and selection of a mutant thriving on urea, as a possible step towards the future production of hydrogen coupled with water treatment. *PLoS ONE* **2018**, *13*, e0198836. [[CrossRef](#)]
454. Miao, R.; Liu, X.; Englund, E.; Lindberg, P.; Lindblad, P. Isobutanol production in *Synechocystis* PCC 6803 using heterologous and endogenous alcohol dehydrogenases. *Metab. Eng. Commun.* **2017**, *5*, 45–53. [[CrossRef](#)] [[PubMed](#)]
455. Miao, R.; Xie, H.; Ho, F.M.; Lindblad, P. Protein engineering of α -ketoisovalerate decarboxylase for improved isobutanol production in *Synechocystis* PCC 6803. *Metab. Eng.* **2018**, *47*, 42–48. [[CrossRef](#)] [[PubMed](#)]
456. Loeschcke, A.; Dienst, D.; Wewer, V.; Hage-Hülsmann, J.; Dietsch, M.; Kranz-Finger, S.; Hüren, V.; Metzger, S.; Urlacher, V.B.; Gigolashvili, T.; et al. The photosynthetic bacteria *Rhodobacter capsulatus* and *Synechocystis* sp. PCC 6803 as new hosts for cyclic plant triterpene biosynthesis. *PLoS ONE* **2017**, *12*, e189816. [[CrossRef](#)] [[PubMed](#)]
457. Carroll, A.L.; Case, A.E.; Zhang, A.; Atsumi, S. Metabolic engineering tools in model cyanobacteria. *Metab. Eng.* **2018**, *50*, 47–56. [[CrossRef](#)] [[PubMed](#)]
458. Yao, L.; Shabestary, K.; Björk, S.M.; Asplund-Samuelsson, J.; Joensson, H.N.; Jahn, M.; Hudson, E.P. Pooled CRISPRi screening of the cyanobacterium *Synechocystis* sp. PCC 6803 for enhanced industrial phenotypes. *Nat. Commun.* **2020**, *11*. [[CrossRef](#)] [[PubMed](#)]
459. Hihara, Y.; Kamei, A.; Kanehisa, M.; Kaplan, A.; Ikeuchi, M. DNA microarray analysis of cyanobacterial gene expression during acclimation to high light. *Plant Cell* **2001**, *13*, 793–806. [[CrossRef](#)] [[PubMed](#)]
460. Hihara, Y.; Sonoike, K.; Kanehisa, M.; Ikeuchi, M. DNA microarray analysis of redox-responsive genes in the genome of the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **2003**, *185*, 1719–1725. [[CrossRef](#)]
461. Kanesaki, Y.; Suzuki, I.; Allakhverdiev, S.I.; Mikami, K.; Murata, N. Salt stress and hyperosmotic stress regulate the expression of different sets of genes in *Synechocystis* sp. PCC 6803. *Biochem. Biophys. Res. Commun.* **2002**, *290*, 339–348. [[CrossRef](#)] [[PubMed](#)]
462. Marin, K.; Suzuki, I.; Yamaguchi, K.; Ribbeck, K.; Yamamoto, H.; Kanesaki, Y.; Hagemann, M.; Murata, N. Identification of histidine kinases that act as sensors in the perception of salt stress in *Synechocystis* sp. PCC 6803. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 9061–9066. [[CrossRef](#)]
463. Marin, K.; Kanesaki, Y.; Los, D.A.; Murata, N.; Suzuki, I.; Hagemann, M. Gene expression profiling reflects physiological processes in salt acclimation of *Synechocystis* sp. strain PCC 6803. *Plant Physiol.* **2004**, *13*, 3290–3300. [[CrossRef](#)]
464. Shoumskaya, M.A.; Paithoonrangarid, K.; Kanesaki, Y.; Los, D.A.; Zinchenko, V.V.; Tanticharoen, M.; Suzuki, I.; Murata, N. Identical Hik-Rre systems are involved in perception and transduction of salt signals and hyperosmotic signals but regulate the expression of individual genes to different extents in *Synechocystis*. *J. Biol. Chem.* **2005**, *280*, 21531–21538. [[CrossRef](#)]
465. Mikami, K.; Kanesaki, Y.; Suzuki, I.; Murata, N. The histidine kinase Hik33 perceives osmotic stress and cold stress in *Synechocystis* sp. PCC 6803. *Mol. Microbiol.* **2002**, *46*, 905–915. [[CrossRef](#)]
466. Ohta, H.; Shibata, Y.; Haseyama, Y.; Yoshino, Y.; Suzuki, T.; Kagasawa, T.; Kamei, A.; Ikeuchi, M.; Enami, I. Identification of genes expressed in response to acid stress in *Synechocystis* sp. PCC 6803 using DNA microarrays. *Photosynth. Res.* **2005**, *84*, 225–230. [[CrossRef](#)] [[PubMed](#)]
467. Uchiyama, J.; Asakura, R.; Kimura, M.; Moriyama, A.; Tahara, H.; Kobayashi, Y.; Kubo, Y.; Yoshihara, T.; Ohta, H. Slr0967 and Slr0939 induced by the SphR response regulator in *Synechocystis* sp. PCC 6803 are essential for growth under acid stress conditions. *Biochim. Biophys. Acta Bioenerg.* **2012**, *1817*, 1270–1276. [[CrossRef](#)]
468. Suzuki, I.; Kanesaki, Y.; Hayashi, H.; Hall, J.J.; Simon, W.J.; Slabas, A.R.; Murata, N. The histidine kinase Hik34 is involved in thermotolerance by regulating the expression of heat shock genes in *Synechocystis*. *Plant Physiol.* **2005**, *138*, 1409–1421. [[CrossRef](#)] [[PubMed](#)]
469. Suzuki, I.; Simon, W.J.; Slabas, A.R. The heat shock response of *Synechocystis* sp. PCC 6803 analysed by transcriptomics and proteomics. *J. Exp. Bot.* **2006**, *57*, 1573–1578. [[CrossRef](#)] [[PubMed](#)]
470. Rowland, J.G.; Pang, X.; Suzuki, I.; Murata, N.; Simon, W.J.; Slabas, A.R. Identification of components associated with thermal acclimation of photosystem II in *Synechocystis* sp. PCC6803. *PLoS ONE* **2010**, *5*, e10511. [[CrossRef](#)] [[PubMed](#)]
471. Paithoonrangarid, K.; Shoumskaya, M.A.; Kanesaki, Y.; Satoh, S.; Tabata, S.; Los, D.A.; Zinchenko, V.V.; Hayashi, H.; Tanticharoen, M.; Suzuki, I.; et al. Five histidine kinases perceive osmotic stress and regulate distinct sets of genes in *Synechocystis*. *J. Biol. Chem.* **2004**, *279*, 53078–53086. [[CrossRef](#)] [[PubMed](#)]
472. Shapiguzov, A.; Lyukevich, A.A.; Allakhverdiev, S.I.; Sergeyenko, T.V.; Suzuki, I.; Murata, N.; Los, D.A. Osmotic shrinkage of cells of *Synechocystis* sp. PCC 6803 by water efflux via aquaporins regulates osmostress-inducible gene expression. *Microbiology* **2005**, *151*, 447–455. [[CrossRef](#)]
473. Kobayashi, M.; Ishizuka, T.; Katayama, M.; Kanehisa, M.; Bhattacharyya-Pakrasi, M.; Pakrasi, H.B.; Ikeuchi, M. Response to oxidative stress involves a novel peroxiredoxin gene in the unicellular cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* **2004**, *45*, 290–299. [[CrossRef](#)] [[PubMed](#)]
474. Houot, L.; Floutier, M.; Marteyn, B.; Michaut, M.; Picciocchi, A.; Legrain, P.; Aude, J.C.; Cassier-Chauvat, C.; Chauvat, F. Cadmium triggers an integrated reprogramming of the metabolism of *Synechocystis* PCC6803, under the control of the Slr1738 regulator. *BMC Genom.* **2007**, *8*. [[CrossRef](#)] [[PubMed](#)]

475. Kanesaki, Y.; Yamamoto, H.; Paithoonrangsarid, K.; Shoumskaya, M.; Suzuki, I.; Hayashi, H.; Murata, N. Histidine kinases play important roles in the perception and signal transduction of hydrogen peroxide in the cyanobacterium, *Synechocystis* sp. PCC 6803. *Plant J.* **2007**, *49*, 313–324. [[CrossRef](#)] [[PubMed](#)]
476. Zhang, Z.; Pendse, N.D.; Phillips, K.N.; Cotner, J.B.; Khodursky, A. Gene expression patterns of sulfur starvation in *Synechocystis* sp. PCC 6803. *BMC Genom.* **2008**, *9*. [[CrossRef](#)]
477. Leplat, C.; Champeimont, R.; Saenkham, P.; Cassier-Chauvat, C.; Jean-Christophe, A.; Chauvat, F. Genome-wide transcriptome analysis of hydrogen production in the cyanobacterium *Synechocystis*: Towards the identification of new players. *Int. J. Hydrog. Energy* **2013**, *38*, 1866–1872. [[CrossRef](#)]
478. Mitschke, J.; Georg, J.; Scholz, I.; Sharma, C.M.; Dienst, D.; Bantscheff, J.; Voß, B.; Steglich, C.; Wilde, A.; Vogel, J.; et al. An experimentally anchored map of transcriptional start sites in the model cyanobacterium *Synechocystis* sp. PCC6803. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 2124–2129. [[CrossRef](#)]
479. Muro-Pastor, A.M.; Hess, W.R. Regulatory RNA at the crossroads of carbon and nitrogen metabolism in photosynthetic cyanobacteria. *Biochim. Biophys. Acta Gene Regul. Mech.* **2020**, *1863*, 194477. [[CrossRef](#)]
480. Baumgartner, D.; Kopf, M.; Klähn, S.; Steglich, C.; Hess, W.R. Small proteins in cyanobacteria provide a paradigm for the functional analysis of the bacterial micro-proteome. *BMC Microbiol.* **2016**, *16*. [[CrossRef](#)]
481. Huang, S.; Chen, L.; Te, R.; Qiao, J.; Wang, J.; Zhang, W. Complementary iTRAQ proteomics and RNA-seq transcriptomics reveal multiple levels of regulation in response to nitrogen starvation in *Synechocystis* sp. PCC 6803. *Mol. Biosyst.* **2013**, *9*, 2565–2574. [[CrossRef](#)]
482. Pei, G.; Sun, T.; Chen, S.; Chen, L.; Zhang, W. Systematic and functional identification of small non-coding RNAs associated with exogenous biofuel stress in cyanobacterium *Synechocystis* sp. PCC 6803. *Biotechnol. Biofuels* **2017**, *10*. [[CrossRef](#)]
483. Hernández-Prieto, M.A.; Semeniuk, T.A.; Futschik, M.E. Toward a systems-level understanding of gene regulatory, protein interaction, and metabolic networks in cyanobacteria. *Front. Genet.* **2014**, *5*. [[CrossRef](#)]
484. Nakao, M.; Okamoto, S.; Kohara, M.; Fujishiro, T.; Fujisawa, T.; Sato, S.; Tabata, S.; Kaneko, T.; Nakamura, Y. CyanoBase: The cyanobacteria genome database update 2010. *Nucleic Acids Res.* **2009**, *38*, D379–D381. [[CrossRef](#)]
485. Zhang, Y.; Fonslow, B.R.; Shan, B.; Baek, M.C.; Yates, J.R. Protein analysis by shotgun/bottom-up proteomics. *Chem. Rev.* **2013**, *113*, 2343–2394. [[CrossRef](#)] [[PubMed](#)]
486. Gao, L.; Wang, J.; Ge, H.; Fang, L.; Zhang, Y.; Huang, X.; Wang, Y. Toward the complete proteome of *Synechocystis* sp. PCC 6803. *Photosynth. Res.* **2015**, *126*, 203–219. [[CrossRef](#)] [[PubMed](#)]
487. Gao, L.; Shen, C.; Liao, L.; Huang, X.; Liu, K.; Wang, W.; Guo, L.; Jin, W.; Huang, F.; Xu, W.; et al. Functional proteomic discovery of Slr0110 as a central regulator of carbohydrate metabolism in *Synechocystis* species PCC6803. *Mol. Cell. Proteom.* **2014**, *13*, 204–219. [[CrossRef](#)]
488. Song, Z.; Chen, L.; Wang, J.; Lu, Y.; Jiang, W.; Zhang, W. A transcriptional regulator Sll0794 regulates tolerance to biofuel ethanol in photosynthetic *Synechocystis* sp. PCC 6803. *Mol. Cell. Proteom.* **2014**, *13*, 3519–3532. [[CrossRef](#)] [[PubMed](#)]
489. Chen, W.; Fang, L.; Huang, X.; Ge, H.; Wang, J.; Wang, X.; Zhang, Y.; Sui, N.; Xu, W.; Wang, Y. Systematic identification of light-regulated cold-responsive proteome in a model cyanobacterium. *J. Proteom.* **2018**, *179*, 100–109. [[CrossRef](#)]
490. Angeleri, M.; Muth-Pawlak, D.; Wilde, A.; Aro, E.M.; Battchikova, N. Global proteome response of *Synechocystis* 6803 to extreme copper environments applied to control the activity of the inducible *petJ* promoter. *J. Appl. Microbiol.* **2019**, *126*, 826–841. [[CrossRef](#)]
491. Battchikova, N.; Vainonen, J.P.; Vorontsova, N.; Keränen, M.; Carmel, D.; Aro, E.M. Dynamic changes in the proteome of *Synechocystis* 6803 in response to CO₂ limitation revealed by quantitative proteomics. *J. Proteome Res.* **2010**, *9*, 5896–5912. [[CrossRef](#)]
492. Miranda, H.; Cheregi, O.; Netotea, S.; Hvidsten, T.R.; Moritz, T.; Funk, C. Co-expression analysis, proteomic and metabolomic study on the impact of a Deg/HtrA protease triple mutant in *Synechocystis* sp. PCC 6803 exposed to temperature and high light stress. *J. Proteom.* **2013**, *78*, 294–311. [[CrossRef](#)]
493. Slabas, A.R.; Suzuki, I.; Murata, N.; Simon, W.J.; Hall, J.J. Proteomic analysis of the heat shock response in *Synechocystis* PCC6803 and a thermally tolerant knockout strain lacking the histidine kinase 34 gene. *Proteomics* **2006**, *6*, 845–864. [[CrossRef](#)] [[PubMed](#)]
494. Rowland, J.G.; Simon, W.J.; Prakash, J.S.S.; Slabas, A.R. Proteomics reveals a role for the RNA helicase *crhR* in the modulation of multiple metabolic pathways during cold acclimation of *Synechocystis* sp. PCC6803. *J. Proteome Res.* **2011**, *10*, 3674–3689. [[CrossRef](#)] [[PubMed](#)]
495. Zhang, L.F.; Yang, H.M.; Cui, S.X.; Hu, J.; Wang, J.; Kuang, T.Y.; Norling, B.; Huang, F. Proteomic analysis of plasma membranes of cyanobacterium *Synechocystis* sp. strain PCC 6803 in response to high pH stress. *J. Proteome Res.* **2009**, *8*, 2892–2902. [[CrossRef](#)] [[PubMed](#)]
496. Kurian, D.; Phadwal, K.; Mäenpää, P. Proteomic characterization of acid stress response in *Synechocystis* sp. PCC 6803. *Proteomics* **2006**, *6*, 3614–3624. [[CrossRef](#)]
497. Ren, Q.; Shi, M.; Chen, L.; Wang, J.; Zhang, W. Integrated proteomic and metabolomic characterization of a novel two-component response regulator Slr1909 involved in acid tolerance in *Synechocystis* sp. PCC 6803. *J. Proteom.* **2014**, *109*, 76–89. [[CrossRef](#)] [[PubMed](#)]

498. Wegener, K.M.; Singh, A.K.; Jacobs, J.M.; Elvitigala, T.; Welsh, E.A.; Keren, N.; Gritsenko, M.A.; Ghosh, B.K.; Camp, D.G.; Smith, R.D.; et al. Global proteomics reveal an atypical strategy for carbon/nitrogen assimilation by a cyanobacterium under diverse environmental perturbations. *Mol. Cell. Proteom.* **2010**, *12*, 2678–2689. [[CrossRef](#)]
499. Fuszard, M.A.; Ow, S.Y.; Gan, C.S.; Noirel, J.; Ternan, N.G.; McMullan, G.; Biggs, C.A.; Reardon, K.F.; Wright, P.C. The quantitative proteomic response of *Synechocystis* sp. PCC6803 to phosphate acclimation. *Aquat. Biosyst.* **2013**, *9*. [[CrossRef](#)]
500. Chen, L.; Zhu, Y.; Song, Z.; Wang, J.; Zhang, W. An orphan response regulator Sll0649 involved in cadmium tolerance and metal homeostasis in photosynthetic *Synechocystis* sp. PCC 6803. *J. Proteom.* **2014**, *103*, 87–102. [[CrossRef](#)]
501. Mehta, A.; López-Maury, L.; Florencio, F.J. Proteomic pattern alterations of the cyanobacterium *Synechocystis* sp. PCC 6803 in response to cadmium, nickel and cobalt. *J. Proteom.* **2014**, *102*, 98–112. [[CrossRef](#)]
502. Fulda, S.; Huang, F.; Nilsson, F.; Hagemann, M.; Norling, B. Proteomics of *Synechocystis* sp. strain PCC 6803: Identification of periplasmic proteins in cells grown at low and high salt concentrations. *Eur. J. Biochem.* **2000**, *267*, 5900–5907. [[CrossRef](#)]
503. Fulda, S.; Mikkat, S.; Huang, F.; Huckauf, J.; Marin, K.; Norling, B.; Hagemann, M. Proteome analysis of salt stress response in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Proteomics* **2006**, *6*, 2733–2745. [[CrossRef](#)]
504. Huang, F.; Fulda, S.; Hagemann, M.; Norling, B. Proteomic screening of salt-stress-induced changes in plasma membranes of *Synechocystis* sp. strain PCC 6803. *Proteomics* **2006**, *6*, 910–920. [[CrossRef](#)]
505. Pandhal, J.; Ow, S.Y.; Wright, P.C.; Biggs, C.A. Comparative proteomics study of salt tolerance between a nonsequenced extremely halotolerant cyanobacterium and its mildly halotolerant relative using in vivo metabolic labeling and in vitro isobaric labeling. *J. Proteome Res.* **2009**, *8*, 818–828. [[CrossRef](#)]
506. Li, T.; Yang, H.M.; Cui, S.X.; Suzuki, I.; Zhang, L.F.; Li, L.; Bo, T.T.; Wang, J.; Murata, N.; Huang, F. Proteomic study of the impact of Hik33 mutation in *Synechocystis* sp. PCC 6803 under normal and salt stress conditions. *J. Proteome Res.* **2012**, *11*, 502–514. [[CrossRef](#)]
507. Qiao, J.; Huang, S.; Te, R.; Wang, J.; Chen, L.; Zhang, W. Integrated proteomic and transcriptomic analysis reveals novel genes and regulatory mechanisms involved in salt stress responses in *Synechocystis* sp. PCC 6803. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 8253–8264. [[CrossRef](#)] [[PubMed](#)]
508. Gao, Y.; Xiong, W.; Li, X.B.; Gao, C.F.; Zhang, Y.L.; Li, H.; Wu, Q.Y. Identification of the proteomic changes in *Synechocystis* sp. PCC 6803 following prolonged UV-B irradiation. *J. Exp. Bot.* **2009**, *60*, 1141–1154. [[CrossRef](#)]
509. Liu, J.; Chen, L.; Wang, J.; Qiao, J.; Zhang, W. Proteomic analysis reveals resistance mechanism against biofuel hexane in *Synechocystis* sp. PCC 6803. *Biotechnol. Biofuels* **2012**, *5*. [[CrossRef](#)] [[PubMed](#)]
510. Qiao, J.; Wang, J.; Chen, L.; Tian, X.; Huang, S.; Ren, X.; Zhang, W. Quantitative iTRAQ LC-MS/MS proteomics reveals metabolic responses to biofuel ethanol in cyanobacterial *Synechocystis* sp. PCC 6803. *J. Proteome Res.* **2012**, *11*, 5286–5300. [[CrossRef](#)] [[PubMed](#)]
511. Tian, X.; Chen, L.; Wang, J.; Qiao, J.; Zhang, W. Quantitative proteomics reveals dynamic responses of *Synechocystis* sp. PCC 6803 to next-generation biofuel butanol. *J. Proteom.* **2013**, *78*, 326–345. [[CrossRef](#)] [[PubMed](#)]
512. Chen, L.; Wu, L.; Wang, J.; Zhang, W. Butanol tolerance regulated by a two-component response regulator Slr1037 in photosynthetic *Synechocystis* sp. PCC 6803. *Biotechnol. Biofuels* **2014**, *7*. [[CrossRef](#)] [[PubMed](#)]
513. Wang, Y.; Chen, L.; Zhang, W. Proteomic and metabolomic analyses reveal metabolic responses to 3-hydroxypropionic acid synthesized internally in cyanobacterium *Synechocystis* sp. PCC 6803. *Biotechnol. Biofuels* **2016**, *9*. [[CrossRef](#)] [[PubMed](#)]
514. Ma, Y.; Yang, M.; Lin, X.; Liu, X.; Huang, H.; Ge, F. Malonylome Analysis Reveals the Involvement of Lysine Malonylation in Metabolism and Photosynthesis in Cyanobacteria. *J. Proteome Res.* **2017**, *15*, 2030–2043. [[CrossRef](#)]
515. Lin, X.; Yang, M.; Liu, X.; Cheng, Z.; Ge, F. Characterization of Lysine Monomethylome and Methyltransferase in Model Cyanobacterium *Synechocystis* sp. PCC 6803. *Genom. Proteom. Bioinform.* **2020**, *18*, 289–304. [[CrossRef](#)] [[PubMed](#)]
516. Mikkat, S.; Fulda, S.; Hagemann, M. A 2D gel electrophoresis-based snapshot of the phosphoproteome in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Microbiology (UK)* **2014**, *160*, 296–306. [[CrossRef](#)] [[PubMed](#)]
517. Spät, P.; Macek, B.; Forchhammer, K. Phosphoproteome of the cyanobacterium *Synechocystis* sp. PCC 6803 and its dynamics during nitrogen starvation. *Front. Microbiol.* **2015**, *6*. [[CrossRef](#)] [[PubMed](#)]
518. Toyoshima, M.; Tokumaru, Y.; Matsuda, F.; Shimizu, H. Assessment of protein content and phosphorylation level in *Synechocystis* sp. PCC 6803 under various growth conditions using quantitative phosphoproteomic analysis. *Molecules* **2020**, *20*, 3582. [[CrossRef](#)]
519. Krall, L.; Huege, J.; Catchpole, G.; Steinhäuser, D.; Willmitzer, L. Assessment of sampling strategies for gas chromatography-mass spectrometry (GC-MS) based metabolomics of cyanobacteria. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* **2009**, *877*, 2952–2960. [[CrossRef](#)]
520. Narainsamy, K.; Cassier-Chauvat, C.; Junot, C.; Chauvat, F. High performance analysis of the cyanobacterial metabolism via liquid chromatography coupled to a LTQ-Orbitrap mass spectrometer: Evidence that glucose reprograms the whole carbon metabolism and triggers oxidative stress. *Metabolomics* **2013**, *9*, 21–32. [[CrossRef](#)]
521. Achyuthan, K.E.; Harper, J.C.; Manginell, R.P.; Moorman, M.W. Volatile metabolites emission by in vivo microalgae—An overlooked opportunity? *Metabolites* **2017**, *7*, 39. [[CrossRef](#)]
522. Majchrzak, T.; Wojnowski, W.; Rutkowska, M.; Wasik, A. Real-Time Volatilomics: A Novel Approach for Analyzing Biological Samples. *Trends Plant Sci.* **2020**, *25*, 302–312. [[CrossRef](#)]

523. Yoshikawa, K.; Hirasawa, T.; Ogawa, K.; Hidaka, Y.; Nakajima, T.; Furusawa, C.; Shimizu, H. Integrated transcriptomic and metabolomic analysis of the central metabolism of *Synechocystis* sp. PCC 6803 under different trophic conditions. *Biotechnol. J.* **2013**, *8*, 571–580. [[CrossRef](#)]
524. Nakajima, T.; Kajihata, S.; Yoshikawa, K.; Matsuda, F.; Furusawa, C.; Hirasawa, T.; Shimizu, H. Integrated metabolic flux and omics analysis of *Synechocystis* sp. PCC 6803 under mixotrophic and photoheterotrophic conditions. *Plant Cell Physiol.* **2014**, *55*, 1605–1612. [[CrossRef](#)] [[PubMed](#)]
525. Cheregi, O.; Miranda, H.; Gröbner, G.; Funk, C. Inactivation of the Deg protease family in the cyanobacterium *Synechocystis* sp. PCC 6803 has impact on the outer cell layers. *J. Photochem. Photobiol. B Biol.* **2015**, *152*, 383–394. [[CrossRef](#)] [[PubMed](#)]
526. Klähn, S.; Orf, I.; Schwarz, D.; Matthiessen, J.K.F.; Kopka, J.; Hess, W.R.; Hagemann, M. Integrated transcriptomic and metabolomic characterization of the low-carbon response using an *ndhR* mutant of *Synechocystis* sp. PCC 6803. *Plant Physiol.* **2015**, *169*, 1540–1556. [[CrossRef](#)]
527. Jablonsky, J.; Papacek, S.; Hagemann, M. Different strategies of metabolic regulation in cyanobacteria: From transcriptional to biochemical control. *Sci. Rep.* **2016**, *6*. [[CrossRef](#)] [[PubMed](#)]
528. Saha, R.; Liu, D.; Hoynes-O'Connor, A.; Liberton, M.; Yu, J.; Bhattacharyya-Pakrasi, M.; Balassy, A.; Zhang, F.; Moon, T.S.; Maranas, C.D.; et al. Diurnal regulation of cellular processes in the cyanobacterium *Synechocystis* sp. strain PCC 6803: Insights from transcriptomic, fluxomic, and physiological analyses. *MBio* **2016**, *7*, e00464–16. [[CrossRef](#)] [[PubMed](#)]
529. Tibiletti, T.; Hernández-Prieto, M.A.; Matthijs, H.C.P.; Niyogi, K.K.; Funk, C. Deletion of the gene family of small chlorophyll-binding proteins (ScpABCDE) offsets C/N homeostasis in *Synechocystis* PCC 6803. *Biochim. Biophys. Acta Bioenerg.* **2016**, *1857*, 396–407. [[CrossRef](#)] [[PubMed](#)]
530. Hasunuma, T.; Matsuda, M.; Kondo, A. Improved sugar-free succinate production by *Synechocystis* sp. PCC 6803 following identification of the limiting steps in glycogen catabolism. *Metab. Eng. Commun.* **2016**, *3*, 130–141. [[CrossRef](#)]
531. Zhu, H.; Ren, X.; Wang, J.; Song, Z.; Shi, M.; Qiao, J.; Tian, X.; Liu, J.; Chen, L.; Zhang, W. Integrated OMICS guided engineering of biofuel butanol-tolerance in photosynthetic *Synechocystis* sp. PCC 6803. *Biotechnol. Biofuels* **2013**, *6*. [[CrossRef](#)] [[PubMed](#)]
532. Liu, Z.X.; Li, H.C.; Wei, Y.P.; Chu, W.Y.; Chong, Y.L.; Long, X.H.; Liu, Z.P.; Qin, S.; Shao, H.B. Signal transduction pathways in *Synechocystis* sp. PCC 6803 and biotechnological implications under abiotic stress. *Crit. Rev. Biotechnol.* **2015**, *35*, 269–280. [[CrossRef](#)] [[PubMed](#)]
533. Gärtner, K.; Klähn, S.; Watanabe, S.; Mikkat, S.; Scholz, I.; Hess, W.R.; Hagemann, M. Cytosine N4-methylation via M.Ssp6803II is involved in the regulation of transcription, fine-tuning of DNA replication and DNA repair in the cyanobacterium *Synechocystis* sp. PCC 6803. *Front. Microbiol.* **2019**, *10*. [[CrossRef](#)]
534. Xu, W.; Wang, Y. Post-translational Modifications of Serine/Threonine and Histidine Kinases and Their Roles in Signal Transductions in *Synechocystis* Sp. PCC 6803. *Appl. Biochem. Biotechnol.* **2020**. [[CrossRef](#)] [[PubMed](#)]
535. Broddrick, J.T.; Rubin, B.E.; Welkie, D.G.; Du, N.; Mih, N.; Diamond, S.; Lee, J.J.; Golden, S.S.; Palsson, B.O. Unique attributes of cyanobacterial metabolism revealed by improved genome-scale metabolic modeling and essential gene analysis. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E8344–E8353. [[CrossRef](#)]
536. Broddrick, J.T.; Welkie, D.G.; Jallet, D.; Golden, S.S.; Peers, G.; Palsson, B.O. Predicting the metabolic capabilities of *Synechococcus elongatus* PCC 7942 adapted to different light regimes. *Metab. Eng.* **2019**, *52*, 42–56. [[CrossRef](#)] [[PubMed](#)]